

THE INTERNATIONAL RHODODENDRON REGISTER AND CHECKLIST (2004) *Second Supplement*

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first Supplement included names registered from 1 January 2003 to 31 December 2004. This second Supplement adds those names registered between 1 January 2005 and 31 December 2005. A list of further corrections to the 2004 *Register and Checklist* is also included.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7-10 of the Introduction. It may be helpful to repeat here that the following abbreviations are employed:

- (a) = azalea
- (az) = azaleodendron
- cv.= cultivar
- G = grown to first flower by...
- gp = Group
- H = hybridised by...
- I = introduced by...
- N = named by...
- (r) = rhododendron
- R = raised by...
- REG = registered by...
- S = selected by...
- (s) = seed parent
- (v) = vireya

Where possible colour references are given to the *RHS Colour Chart* (1966, 1986, 1995 & 2001) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS web site (www.rhs.org.uk/plants/registerpages/Rhodoform.PDF) or from national or regional registrars as follows:

North America	Mrs J.W. Murray, 21 Squire Terrace, Colts Neck, New Jersey 07722-1021, USA
Australia	Mr K. Gillanders, Woodbank, 2040 Huon Road, Longley, Tasmania 7150, Australia
New Zealand	Mr B.J. Coker, 42 Massey Crescent, Christchurch 1, New Zealand
Japan	Mr K. Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

**Dr A.C. Leslie, International Rhododendron Registrar, 109 York Street, Cambridge CB12PZ, England
(Email: alanleslie@rhs.org.uk)**

INTERNATIONAL RHODODENDRON REGISTER & CHECKLIST

(r) 'Above All'

cv. Elepidote rhododendron: 'Award' (s) × 'Ayers Rock'. H: H. O'Rourke (1987), G: H. O'Rourke (1999), N: H. O'Rourke (2005), REG: H. O'Rourke (2005). Fls 9/domed truss, broadly funnel-shaped, 65 × 100mm, with 7 very wavy-edged lobes, orange-pink in bud, opening inside yellow in the throat and pale pink on the lobes, outside pale pink. Calyx 20mm long, yellow and pink. Lvs lanceolate, 110 × 40mm, margins flat, matt green, obtuse at base, acute at apex, hairless. Shrub 1.5 × 1.8m in 16 years. November (Australia).

(r) 'Albarelo'

cv. Elepidote rhododendron: 'Holger' (s) × 'Schneebukett'. H: H. Hachmann (1990), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 15-17/fairly compact truss, broadly funnel-shaped, 35-45 × 65-75mm, with 5 slightly wavy-edged lobes, buds light to pale purplish pink (62C/D), opening inside and out white, delicately edged pale purplish pink (62D); conspicuous blotch on the dorsal lobe with mixed markings of deep greenish yellow to strong yellow (153A-D) and deep to strong orange-yellow (163A/B); stamens 10, conspicuous, filaments white, anthers dark pink (182D); styles pale yellow-green (149D), stigma dark red (187B); ovary moderate yellow-green (138B), white hairy. Calyx 3-5mm long, light yellow-green (145B/C). Lvs elliptic to ovate, 110 × 40-50mm, margins slightly wavy-edged, dark matt green above, rounded at base, apiculate at apex, with sparse brownish hairs below. Shrub c.1.3 × c.1.5m in 10 years. Late May to early June. (First published in Hachmann cat. 2004-2008 (2004), p.26. Nomenclatural standard: colour print provided by the registrant (WSY0070129)).

(a) 'Alba Rosa Cavadini'

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 32 × 24mm, with 5 flat-edged lobes, buds strong pink (52D), opening deep pink (52C) throughout; three lobes with deep purplish pink (61D) markings; stamens 7-10, exerted from corolla, anthers light olive brown (199B), filaments light purplish pink (55C); pistil equalling or just exceeding the stamens, style light purplish pink (55C), shaded strong purplish pink (55B), stigma light yellow (21D). Lvs lanceolate, 29 × 12 mm, margins flat, glossy above, obtuse at base, acute at apex, hairy, moderate olive green (146A) above and moderate yellow-green (146D) below when mature; petioles 6-7mm long. Shrub 2.0 × 3.0m in 60 years (regularly pruned). May. (Etymology: colour of the flowers resembles the pink colour of the sky at daybreak ("alba" in Italian)).

(a) 'Alsides Ferrari da Barbè'

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 60 × 44mm, with 5 flat or rarely wavy-edged lobes, vivid reddish orange (43A) in bud and throughout on opening; three lobes with strong red (46A) markings, shaded with deep red (53A); stamens 6-7, not exerted from the corolla, anthers dark red (59A), filaments moderate red (47A); pistil shorter than stamens, style vivid red (52A), stigma deep red (53A). Lvs lanceolate, 49 × 25mm, margins wavy, glossy above, acute at base and apex, hairy, strong yellow-green (144A) above and moderate olive green (146A) below when mature; petioles 4mm long. Shrub 1.3 × 1.4m in 50 years. April to May. (Etymology: named for a friend of Mario Carmine who lives in Barbè, Italy).

(r) 'Amarlie Crowden'

cv. Elepidote rhododendron: 'Miss B.L. Jones' (s) × 'D.J. Dosser's Choice'. H: D.J. Dosser (1995), G: D.J. Dosser (2003), N: D.J. Dosser (2004), REG: D.J. Dosser (2005). Fls 18/conical truss, funnel-shaped, 40 × 100mm, with 5 wavy-edged lobes, moderate purplish pink (66D) in bud, opening inside pale purplish pink (65D), edged deep purplish pink (66C), outside deep purplish pink (66C); blotch of reddish spots over dorsal lobe and parts of the adjacent lobes. Lvs oblong, 190 × 75mm, glossy green, obtuse at base, mucronate at apex, margins curved down, hairless. Shrub 1.0 × 0.6m in 8 years. November (Australia). (Nomenclatural standard: colour print provided by the registrant (WSY0070185)).

(r) 'Anatevka'

cv. Elepidote rhododendron: 'Galaxis' (s) × 'Violette Funken'. H: H. Hachmann (1995), G: H. Hachmann (2000), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 11/lax ball truss, open funnel-shaped, 50-60 × 70-100mm, with 5 very wavy-edged lobes (some 6-lobed), inside strong purple (83D) with the centre of lobes unevenly marked whitish, outside entirely strong purple (83D) with the midveins strong purple (77B); dorsal lobe with a large, conspicuous blotch and markings of dark red (187A); stamens 10-12, filaments strong reddish purple (72C), some filaments occasionally petaloid, anthers pale yellow (158A-B); styles strong purplish pink (73B), stigma dark red (187A/B), ovary reddish green, hairless. Calyx 1mm long, light yellow-green (144D). Lvs narrowly elliptic, 80-110 × 30-38mm, margins slightly decurved, matt mid green above, cuneate at base, acute at apex, with a scarcely noticeable indumentum. Shrub c.1.3 × c.1.5m in 10 years. Late May to mid June. (First published in

Hachmann cat. 2004-2008 (2004), p.27.
Nomenclatural standard: colour print provided by the registrant (WSY0070130).

(r) **'Ao-Chin'**

cv. Elepidote rhododendron: 'Roza Stevenson' (s) × Albatross Group. H: M.L.A. Robinson (1990), G: M.L.A. Robinson, N: M.L.A. Robinson, REG: M.L.A. Robinson (2005). Fls 10-12/rounded truss, funnel-campanulate, 50-65 × 110-120mm, with 7 very wavy-edged lobes, moderate purplish pink (74D) in bud - or close to this colour - opening at darkest to moderate purplish pink (74D) to light purple (75A) on lobes, fading at edges to yellowish white (158D) and to pale yellow (158B) in the throat; outside with lobes moderate purplish pink (74D) to light purple (75A), with a small area of yellowish white (158D) near the calyx; unmarked; stamens 14-16, filaments white, anthers brown; style pale yellow-green, stigma moderate yellow-green (146D); ovary green, dotted with short-stalked white glands; slightly scented. Calyx 2-4mm long, green. Lvs lanceolate to oblong, 120-180 × 50-75mm, margins flat, glossy above, slightly cordate or obtuse at base, acute at apex, with a slight fawn indumentum; petioles red. Shrub 2.4 × 2.0m in 24 years. Mid April. PC 2005. (Nomenclatural standard: herbarium specimen provided by the registrant (WSY0060731). Etymology: named for a Chinese deity).

(r) **'Aprilmorgen'**

cv. Elepidote rhododendron: probably *yakushimanum* (s) × *adenogynum*. S: H. Hachmann (1998), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 9-10/compact truss, broadly funnel-shaped, 50 × 66-75mm, with 5 wavy-edged lobes, opening inside pure white (155D), outside pure white (155D); unspotted; stamens 10, filaments white; style brilliant yellow-green (150C). Calyx 3.5mm long, whitish green, edged red. Lvs elliptic, 120 × 36mm, convex, margins decurved, slightly glossy and dark green above, cuneate at base, acute at apex, with a dense, cinnamon brown, hairy indumentum below when mature. Shrub c.0.6 × c.1.1m in 10 years; habit flat, dense and broadly rounded. Late April to early May. (Early for a *yakushimanum* hybrid. Nomenclatural standard: colour illustration provided by registrant (WSY0070173). First published in Hachmann cat. 2004-2008 (2004), p.90).

(a) **'Arneson Cameo'**

cv. Deciduous azalea: (F3 selection from ('Gallipoli' × 'Cecile')) (s) × (F3 selection from ('Gallipoli' × 'Cecile')). H: I. & R. Arneson (1980), G: I. & R. Arneson (1986), N: Briggs Nursery (2005), I: Briggs Nursery (2005), REG: I. & R. Arneson (2005). Fls 12-14/ball truss, funnel-shaped, 50 × 70mm, with 6+6 very wavy-edged lobes, deep red (53A) in bud, opening inside strong pink (48C) with a brilliant orange-yellow (21B) dorsal blotch, outside deep pink (48A). Calyx c 1.5mm long, moderate yellow-green (139C). Lvs elliptic, 45-70 × 20-40mm, broadly acute

at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Dense shrub, 0.5 × 0.7m in 24 years. Mid to late May.

(a) **'Arneson Fairy Jewel'**

cv. Deciduous azalea: (('Gallipoli' × 'Cecile') × 'Pink Delight') (s) × ('Strawberry Ice' × 'Sunset Pink'). H: I. & R. Arneson (1984), G: I. & R. Arneson (1989), N: Briggs Nursery (2005), I: Briggs Nursery (2005), REG: I. & R. Arneson (2005). Fls 5-7/ball truss, open funnel-shaped, 50 × 75mm, with 5+5 wavy-edged lobes, strong purplish red (63A) in bud, opening inside moderate purplish pink (68C), outside strong purplish pink (68B). Calyx c.1.5mm long, moderate yellow-green (138B). Lvs elliptic, 40-65 × 20-25mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above. Dense shrub 0.5 × 0.6m in 20 years. Mid May.

(a) **'Arneson Flame'**

cv. Deciduous azalea: un-named red Exbury seedling (s) × un-named red Exbury seedling. H: I. & R. Arneson (1987), G: I. & R. Arneson (1992), N: Briggs Nursery (2005), I: Briggs Nursery (2005), REG: I. & R. Arneson (2005). Fls 8-12/ball truss, tubular funnel-shaped, 55 × 70mm, with 5 very wavy-edged lobes, strong red (46A) in bud, opening inside vivid red (46B), outside vivid red (46C); slightly scented. Calyx c.3mm long, light yellow-green (139D), broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Shrub with a dense habit, 0.4 × 0.6m in 17 years. Mid to late May.

(a) **'Arneson Frilly Orange'**

cv. Deciduous azalea: parentage unknown. H: I. & R. Arneson (1986), G: I. & R. Arneson (1992), N: I. & R. Arneson (1993), I: Briggs Nursery (2005), REG: I. & R. Arneson (2005). Fls 4-8/ball truss, tubular funnel-shaped, 55 × 75mm, with 6-8 very wavy-edged lobes, vivid red (46B) in bud, opening vivid reddish orange (32A) throughout; moderately scented. Calyx c.1.5mm long, moderate yellow-green (139D). Lvs elliptic, 50-90 × 25-35mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and strong yellow-green (147A) above, hairless. Dense shrub, 0.5 × 0.7m in 18 years. Mid to late May.

(a) **'Arneson Golden Solitaire'**

cv. Deciduous azalea: un-named, yellow, Exbury seedling (s) × un-named, yellow, Exbury seedling. H: I. & R. Arneson (1986), G: I. & R. Arneson (1991), N: Briggs Nursery (2005), I: Briggs Nursery (2005), REG: I. & R. Arneson (2005). Fls 4-6/ball truss, tubular funnel-shaped, 65 × 75mm, with 5 lobes, strong orange (25A) in bud, opening inside vivid yellow (16A) with a strong orange-yellow (17A) dorsal flare, outside vivid yellow (16A). Calyx c.3mm long, moderate yellow-green (139D). Lvs elliptic, 25-50 × 20-25mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Dense shrub 0.4 × 0.7m in 18 years. Early to mid May.

(a) **'Arneson Medallion'**

cv. Deciduous azalea: parentage unknown. H: I. & R. Arneson (1986), G: I. & R. Arneson (1992), N: Briggs Nursery (2005), I: Briggs Nursery (2005), REG: I. & R. Arneson (2005). Fls 11-15/ball truss, tubular funnel-shaped, 50 × 70mm, with 5 very wavy-edged lobes, vivid reddish orange (42A) in bud, opening vivid reddish orange (33A) throughout; slightly scented. Calyx c.3mm long, moderate yellow-green (139D). Lvs elliptic, 45-70 × 20mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Dense shrub 0.4 × 0.8m in 18 years. Early May.

(a) **'Arneson Ruby Princess'**

cv. Deciduous azalea: (('Gibraltar' × 'Favor Major') × 'Corringe') (s) × 'Wallowa Red'. H: I. & R. Arneson (1986), G: I. & R. Arneson (1992), N: Briggs Nursery (2005), I: Briggs Nursery (2005), REG: I. & R. Arneson (2005). Fls 6-8/ball truss, tubular funnel-shaped, 55 × 75mm, with 5 very wavy-edged lobes, strong red (46A) in bud, opening inside vivid red (46B), outside strong red (46A); slightly scented. Calyx 3mm long, moderate yellow-green (138B). Lvs elliptic, 40-65 × 20-25mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Dense shrub, 0.5 × 0.7m in 18 years. Mid May.

(r) **'Avram'**

cv. Elepidote rhododendron: 'Girard Joshua' (s) × 'Caroline' (Gable). H: J.D. Hooper (1992), G: J.D. Hooper (c 1997), N: J.D. Hooper (2005), REG: J.D. Hooper (2005). Fls 9-12/domed truss, broadly funnel-shaped, 60 × 95mm, with 5 wavy-edged lobes, strong purplish red (72A) in bud, opening inside and out strong reddish purple (72B) at margins, blending to light purple (87D) at centre; faint dorsal flare of moderate orange-yellow (164B), with greenish speckling. Calyx c.3mm long, strong purplish red (60D). Pedicels strong purplish red (60D). Lvs elliptic, 130 × 55mm, broadly acute at apex, oblique at base, with wavy margins, dull and dark olive green above, hairless. Shrub 1.8 × 2.0m in c.13 years. Late May.

(r) **'Baronne Nadine de Rothschild'**

cv. Elepidote rhododendron: 'Mrs T.H. Lowinsky' × 'One Thousand Butterflies'. H: M. Colombel (1999), G: M. Colombel (2003), N: M. Colombel (2005), REG: M. Colombel (2005). Fls 12-19/more-or-less ball (or slightly conical) truss, funnel-shaped, 60 × 80mm, with 5 wavy-edged lobes, strong reddish purple (78B) in bud, opening inside light purple (80C), outside moderate purplish pink (78D); red blotch all around base of throat, with a solid, blackish mark and blackish spots on dorsal lobe. Lvs elliptic, 140-170 × 50-70mm, convex but with flat margins, rounded at base, broadly acute at apex, hairless, very dark green above, somewhat bullate. Shrub 1.1 × 1.0m in 5 years. Early to mid May. (Nomenclatural standard: colour slide provided by the registrant (WSY0070243).

(r) **'B.C.B.G.'**

cv. Elepidote rhododendron: 'Party Pink' (s) × 'September Song'. H: M. Colombel (1998), G: M. Colombel (2003), N: M. Colombel (2005), REG: M. Colombel (2005). Fls 8-12/domed truss, funnel-shaped, 70 × 75mm, with 6 wavy-edged lobes, strong purplish red (63A) in bud, opening inside and out deep purplish pink (64D), with a large yellow blotch (most extensive on dorsal lobes but also on parts of the ventral lobes and in the throat), the dorsal lobes with red spots, base of the throat nearly black; no stamens; scented. Calyx 2mm long, light green. Lvs elliptic, 130-150 × 60-70mm, margins flat, matt green above, cuneate at base, broadly acute at apex, hairless; petioles red. Shrub 0.9 × 1.0m in 6 years. Early to mid May. (Nomenclatural standard: colour slide provided by the registrant (WSY0070245). Etymology: epithet is a French acronym meaning chic and conservative ("Bon Chic Bon Genre").

(a) **'Bèe'**

cv. Evergreen azalea: parentage unknown. R: Fratelli Savioli (1970), N: Fratelli Savioli (1985), I: Fratelli Savioli (1985), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 40 × 30mm, with 5 wavy-edged lobes, buds moderate purplish pink (62B), shaded vivid reddish orange (43B), opening white, shaded strong purplish pink (62A) inside and out; three lobes with moderate red (180B) markings; stamens 5, exerted from corolla, anthers moderate orange (167B), filaments white; pistil distinctly exceeding stamens, style pale yellow-green (149D), stigma brilliant yellow-green (149A). Lvs obovate, 28 × 15mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (137A) above and moderate yellowish green (138B) below when mature; petioles 5mm long. Shrub 1.0 × 1.0m in 10 years. April. (Etymology: named for a village near the city of Verbania (Italy). The latter is where the Savioli nursery is situated).

(r) **'Beguiling'**

cv. Elepidote rhododendron: *nuttallii* (s) × 'Countess of Haddington'. H: H. O'Rourke (1995), G: H. O'Rourke (2003), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 3/loose truss, tubular funnel-shaped, 70 × 30mm, with 5 wavy-edged lobes, pink and white in bud, opening inside white with a yellow eye, outside white with pink stripes down each midvein. Calyx 15mm long, red. Lvs elliptic, 100 × 40 mm, margins wavy, matt green above, cuneate at base, acute at apex, hairless. Shrub 1.4 × 1.1m in 8 years. October (Australia).

(r) **'Bethany Jade'**

cv. Elepidote rhododendron: 'Arthur Bedford' (s) × 'Jonathan Shaw'. H: unknown, S: J. Doppel (1992), G: J. Doppel (1999), N: J. Doppel (2004), REG: J. Doppel (2005). Fls 10/domed truss, saucer-shaped, 50 × 75mm, with 5 slightly wavy-edged lobes, deep purple in bud, opening inside moderate purple on dorsal lobes and margins of ventral lobes, the ventral lobes shade to very light purple towards the centre; outside purple; prominent dorsal markings of

discrete, dark spots. Lvs oblong, 150 × 45mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and mid green above, hairless. Shrub 1.5 × 1.5m in 10 years. Late May. (Reported to be seed and pollen fertile; does not sprout from the base.)

(r) **'Betty J. Miller'**

cv. Elepidote rhododendron: 'Autumn Gold' (s) × *fortunei* (selection from Gable). H: G.T. Miller, G: G.T. Miller, N: G.T. Miller (2005), REG: G.T. Miller (2005). Fls 8/lax truss, broadly funnel-shaped, c.75 × 100mm, with 7 wavy-edged lobes, light yellowish pink (27A) in bud, with darker veins, opening pale yellow-green (4D) changing to light greenish yellow (4C) in the throat and green at the base; outside pale yellowish pink (27C); no markings. Calyx c.20mm long, light greenish yellow (4C) with light yellowish pink (29C) veins. Lvs elliptic, 140 × 65mm, broadly acute at apex, rounded at base, margins flat, dull and moderate yellow-green (137C) above, hairless. Shrub 1.8 × 1.2m in 10 years. Midseason.

(r) **'Bisque'**

cv. Elepidote rhododendron: 'Rio' (s) × 'Tan'. H: T. Ahern (1996), G: J. Doppel (2001), N: J. Doppel (2004), REG: J. Doppel (2005). Fls 10/ball truss, saucer-shaped, 50 × 95mm, with 6 flat-edged lobes, apricot in bud, opening inside light peach shading slightly darker in the centre, outside peach; sparse greenish brown dorsal spotting. Lvs oblong, 140 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and mid green above, hairless; withstanding winter wind and sun well. Shrub 1.2 × 0.9m in 10 years. Early May. (Reported to be pollen sterile but seed fertile.)

(r) **'Blithe Spirit'**

cv. Elepidote rhododendron: 'Rhonda' (s) × 'Boulter's Robyn'. H: H. O'Rourke (1987), G: H. O'Rourke (1996), N: H. O'Rourke (2005), REG: H. O'Rourke (2005). Fls 15/loose truss, broadly funnel-shaped, 50 × 70mm, with 5 very wavy-edged lobes, red in bud, opening inside strong purplish pink (55B) with the tube fading to white with age, outside pale pink. Calyx 2mm long, yellowish green. Lvs lanceolate, 110 × 30mm, margins decurved, matt green above, obtuse at base, acute at apex, hairless. Shrub 1.0 × 1.0m in 15 years. September to October (Australia).

(r) **'Bordüre'**

cv. Elepidote rhododendron: 'Hachmann's Charmant' (s) × 'Schneespiegel'. H: H. Hachmann (1990), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 10-14/dense, ball truss, broadly funnel-shaped, 45 × 65-80mm, with 5(-6) wavy-edged lobes, buds strong purplish red (63A/B), opening inside pure white, edged strong purplish red (60D), outside strong purplish red (60D), shaded strong purplish red (60C), with slightly darker (60B) veins; conspicuous blotch of dark red (187A/B) in dorsal throat with markings of deep purplish red (187D) spreading onto

dorsal lobe; stamens 10(-12), filaments white (155C), anthers moderate yellow (161B); style white (155D), stigma deep purplish red (187D); ovary strong yellow-green (144B), silvery hairy. Calyx 1-2mm long, light greenish/red. Lvs ovate, 110-130 × 60-70mm, with flat or slightly wavy margins, light green and slightly glossy above, rounded at base, apiculate at apex, with some hardly noticeable brownish hairs below. Shrub c.1.2 × c.1.1m in 10 years; habit erect, rather loose, with long-lasting flowers. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070132). Published in Hachmann cat. 2004-2008 (2004), p.33.)

(a) **'Brenda Marie'**

cv. Evergreen azalea: un-named seedling (no. CB-1 from Bob Stewart) (s) × ('Elsie Lee' × 'Dancing Butterfly'). H: J.F. Klimavicz (1997), G: J.F. Klimavicz (1999), N: J.F. Klimavicz (2005), I: J.F. Klimavicz (2005), REG: J.F. Klimavicz (2005). Fls 2-3/ball truss, saucer-shaped, 55 × 100mm, held erect, with 5 wavy-edged lobes, white in bud, opening inside white, the throat tinted light yellow-green (154D) and with a very large, strong to light yellow-green (145A-B), speckled, dorsal blotch; outside white; pistil 50mm long; stigma 2mm wide, light yellow-green (145B). Calyx 10mm long, moderate yellowish green (138B). Lvs elliptic, 55 × 15mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above. Dense shrub, 0.6 × 0.6m in 5 years. Mid April to May.

(a) **'Briana Murin'**

cv. Evergreen azalea: ('Desiree' × unknown (a semi-double, evergreen/semi-evergreen azalea known as CB-1)) (s) × 'Nami'. H: Robert Stewart (1994), G: Robert Stewart (1997), N: Robert Stewart (2004), I: Robert Stewart (2004), REG: Robert Stewart (2005). Fls semi-double, 2/truss, saucer-shaped, 45 × 70mm, with 5 wavy-edged lobes, lighter than deep purplish pink (68A) in bud, opening inside slightly lighter than moderate purplish pink (78D) with brownish yellow dorsal spotting; outside similar to bud colour. Calyx c. 5mm long, brilliant yellow-green (142B). Lvs elliptic, 45 × 20mm, broadly acute at apex, cuneate at base, margins upcurved, semi-glossy and mid green above. Shrub 0.6 × 0.75m in 8 years. Midseason.

(r) **'Brookhaven'**

cv. Elepidote rhododendron: 'Hotei' (s) × 'Janet Blair'. H: W. Brack (1987), G: W. Brack (1995), N: W. Brack (2004), I: Roslyn Nursery (2004), REG: W. Brack (2005). Fls 15/domed truss, open funnel-shaped, 60 × 95mm, with 6 wavy-edged lobes, light yellowish pink (26D) in bud, opening light greenish yellow (4C) inside with two small dorsal flares composed of discrete, vivid reddish orange (43A) spots; outside pale yellow-green (4D). Calyx c.5mm long, light yellow-green (154D). Lvs elliptic, 90 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137A) above, hairless. Shrub 1.5 × 1.5m in 17 years. Midseason.

(r) 'Bush's Bush'

cv. *Elepidote rhododendron*: 'Murray's Pink' (s) × 'Roy Stockman'. H: M. Stockman (1993), G: M. & H. Stockman (2001), N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 12-15/ball truss, open funnel-shaped, 55 × 90mm, with 6 wavy-edged lobes, deep red (53A) in bud, opening inside with a distinct, strong red (53D) band, c.15-20mm wide at lobe margins, changing abruptly to pale yellowish pink (36D); outside strong red (53B); dorsal throat with two freckled markings of deep red (53A), c.20mm long, on a pale greenish yellow (10D) ground. Calyx c.10mm long, dark greenish yellow (152D). Lvs oblong, 190 × 55mm, acute at apex, cuneate at base, margins flat, downcurved, semi-glossy and moderate olive green (137A) above, moderate yellow-green (146C) below, hairless. Shrub 1.0 × 0.9m in 11 years. Late May.

(r) 'Bzzz Bee'

cv. *Elepidote rhododendron*: 'Blimey Limey' (s) × 'Gallacci's Gold'. H: M. Stockman (1993), G: M. & H. Stockman (2001), N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 5-7/lax truss, open funnel-shaped, 45 × 65mm, with 5-6 wavy-edged lobes, buds light orange-yellow (19A) at base, shading to strong yellowish pink (31C) at tips, opening inside light yellow (20B) shading to strong yellowish pink (31C) on outer rim of lobes; outside similarly coloured but with more strong yellowish pink (31C) along side of lobes; dark greenish yellow (152C) freckles on inside of dorsal lobe and on portions of adjacent lobes. Calyx lobes c.20-25mm long with one separated, shorter lobe c.10-12mm long, light yellow (20B) with lines of dark greenish yellow (152C) freckles on basal portion of tube. Lvs oblong, 100 × 40mm, broadly acute at apex, oblique at base, margins flat, upcurved, semi-glossy and moderate olive green (137B) above, moderate yellow-green (147D) below, hairless. Sprawling shrub, 0.75 × 0.75m in 11 years. Late April.

(a) 'Cagle Nocturne'

cv. *Deciduous azalea*: *arborescens* (s) × *cumberlandense*. H: J.E. Schild (1986), G: J.E. Schild (1994), N: J.E. Schild (2003), I: Schild Azalea Gardens & Nursery (2006), REG: J.E. Schild (2005). Fls 15-28/ball truss, tubular funnel-shaped, 45 × 50mm, with 5 flat-edged lobes, buds strong purplish pink (68B), with many, white, glandular setae, opening inside moderate purplish pink (68C) with a strong orange (25B) dorsal blotch; outside light purplish pink (68D) with many white hairs and glandular setae along centre rib of tube and on corolla lobes; filaments light purplish pink (68D); style deep purplish pink (68A); stigma strong purplish red (71C); strongly scented. Calyx c. 3mm long, brilliant greenish yellow (1B). Lvs ovate, 75 × 40mm, broadly acute at apex, cuneate at base, margins upcurved, semi-glossy and moderate yellow-green (137C) above, light yellow-green (138D) below; hairs along margins and also sparsely on midveins below (hairs white at first, aging to green-white).

Shrub 1.5 × 1.2m in 10 years. June. (Published in *The Azalean* 27(2):33 (2005)).

(a) 'Caldè'

cv. *Evergreen azalea*: parentage unknown. R: Fratelli Savioli (1970), N: Fratelli Savioli (1985), I: Fratelli Savioli (1985), REG: Prof. E. Accati (2005). Fls single, 2-5/truss, 50 × 38mm, with 5-7 flat-edged lobes, vivid red (45A) in bud and on opening, both inside and out; three lobes with strong red (46A) markings; stamens 5-7, exerted from the corolla, anthers deep purplish red (59B), filaments pale purplish pink (56D); pistil distinctly exceeding stamens, style strong red (53B), stigma vivid reddish orange (42A). Lvs lanceolate, 39 × 16mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (146A) above and moderate yellow-green (146D) below when mature; petioles 4-8mm long. Shrub 1.0 × 1.0m in 10 years. April. (Etymology: named after a village near the city of Verbania, the latter being where the Savioli nursery is situated).

(r) 'Camparina'

cv. *Elepidote rhododendron*: 'Busuki' (s) × 'Phyllis Ballard'. H: H. Hachmann (1993), G: H. Hachmann (1999), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 7-11/rather loose truss, broadly funnel-shaped, 50-55 × 70-80mm, with 5 wavy-edged lobes, buds strong red (45D), opening inside deep pink (50B), shaded light yellow (19B) on dorsal lobe and throat; outside strong red (45D), shaded with vivid red (45C), with vivid red/deep pink (52A/B) nerves; no blotch or markings; stamens 10, filaments strong purplish red (58C), anthers pale yellow (164D); style dark pink (182D), stigma moderate red (185B); ovary greenish, white hairy. Calyx 1mm long, strong to light yellow-green (144C/D). Lvs narrowly lanceolate, 90-110 × 35-45mm, margins flat, matt mid green above, rounded at base, apiculate at apex. Shrub c.0.8 × c.1.1m in 10 years; habit broadly rounded, slow-growing. Late May to mid June. (Nomenclatural standard: colour print provided by registrant (WSY0070131). Published in Hachmann cat. 2004-2008 (2004), p.36).

(a) 'Cannobbio'

cv. *Evergreen azalea*: parentage unknown. R: Fratelli Savioli (1970), N: Fratelli Savioli (1985), I: Fratelli Savioli (1985), REG: Prof. E. Accati (2005). Fls single, 1-3/truss, 32 × 90mm, with 5 very wavy-edged lobes, buds white shaded brilliant yellow-green (149A), opening white throughout; three lobes with brilliant yellow-green (149A) markings; stamens 7-8, exerted from the corolla, anthers orange-yellow (164B), filaments white; pistil distinctly exceeding stamens, style pale yellow-green (149D), stigma brilliant yellow-green (150B). Lvs oblanceolate, 45 × 21mm, margins flat, glossy above, acute at apex, obtuse at base, hairy, moderate olive green (146A) above and moderate yellow-green (147C) below when mature; petioles 8mm long. Shrub 1.0 × 1.0m in 10 years. May. (Etymology: named after a village

near the city of Verbania, the latter being where the Savioli nursery is situated).

(a) **'Carillon Melody'**

cv. Evergreen azalea: 'Reinhold Ambrosius' (s) × 'Orchiphillum'. H: C. Carroll (1985), G: C. Carroll (1988), N: C. Carroll (2004), I: Arborglen Pty Ltd, REG: Carroll's Country Gardens Nursery (2005). Fls single, 2/truss, broadly funnel-shaped, with 5 very broadly overlapping, wavy-edged lobes, vivid red (57A) in bud, opening vivid purplish red (57B) inside with vivid red (57A) dorsal spotting. Calyx 7mm long, green. Lvs elliptic, 20 × 10mm, margins flat, matt green, hairy. Shrub 1.0 × 1.0m in 5 years. Spring. (Nomenclatural standard: colour print provided by the registrant (WSY0070183)).

(a) **'Carla Ferrero'**

cv. Evergreen azalea: parentage unknown. N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single, 1-2/truss, 66 × 45mm, with 5 wavy or some very wavy-edged lobes, white in bud and when open both inside and out; three lobes with light yellow-green (150D) markings; stamens 10, not exerted from the corolla, anthers moderate orange-yellow (165C), filaments white; style distinctly exceeding corolla, white (155C), stigma brilliant yellow-green (154C). Lvs lanceolate, 37 × 14mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (146C) below when mature; petioles 3-5mm long. Shrub 1.5 × 2.5m in 100 years (but regularly pruned). May. (Probably raised (c.1890) by Fratelli Rovelli and Fratelli Contini, gardeners at the Villa del Pascià in the 19th century, the villa where the King of Egypt stayed during his Italian holidays on Lake Maggiore. Etymology: named for the wife of the Director of the International Book Exhibition in Turin).

(a) **'Carmine Emilia'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 2/truss, 80 × 47mm, with 5 wavy to rarely very wavy-edged lobes, buds red-purple (N57B), opening red-purple (N57C) throughout; three lobes with strong purplish red (61B) markings shaded strong purplish red (60B); stamens 8-9, some exerted from the corolla, anthers deep red (60A), filaments vivid purplish red (61C) shaded deep purplish pink (61A) and white; pistil equalling or shorter than stamens, style vivid purplish red (61C), shaded deep purplish pink (61D), stigma strong greenish yellow (151A). Lvs lanceolate, 42 × 19mm, margins flat, glossy above, obtuse at apex, hairy, moderate olive green (137A) above and strong yellow-green (144A) below when mature; petioles 3-5mm long. Shrub 1.5 × 2.8m in 60 years. April to May. (Etymology: named for a sister of Mario Carmine).

(a) **'Carmine Paola'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M.

Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 3-8/hemispherical truss, 52 × 36mm, with 5 flat-edged lobes, buds strong red (47B), opening deep yellowish pink (47C) inside and out; three lobes with strong red (46A) markings shaded deep yellowish pink (43C); stamens 5-6, exerted from the corolla, anthers brilliant yellow (20A), filaments white, shaded strong red (47B); pistil equalling or exceeding stamens, style deep yellowish pink (46D), stigma moderate pink (49B) shaded strong red (46A). Lvs ovate, 35 × 18mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (147B) below when mature; petioles 3mm long. Shrub 1.0 × 1.7m in 50 years. April to May. (Etymology: named for a sister of Mario Carmine).

(a) **'Carmine Pierina'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls hose-in-hose, 2/truss, 58 × 30mm, with 5+5 usually wavy-edged lobes (rarely very wavy and lobed), buds red-purple (N66B), opening red-purple (N66C) throughout; three lobes with strong purplish red (67A) markings; stamens 5-6, exerted from the corolla, anthers brilliant yellow (13C), filaments vivid purplish red (67B) shaded white; pistil equalling or shorter than the stamens, style strong red (53D), stigma moderate pink (49B). Lvs oblanceolate, 23 × 12mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (137A) above and moderate yellow-green (138B) below when mature; petioles 4mm long. Shrub 1.3 × 1.5m in 50 years. April to May. (Etymology: named for a sister of Mario Carmine).

(r) **'Carol's Choice'**

cv. Elepidote rhododendron: *fortunei* (s) × 'Burgundy'. H: C. McNew, G: E.T. Anderson, N: Mrs E.T. Anderson, REG: Mrs E.T. Anderson (2005). Fls 10/ball truss, broadly funnel-shaped, 75 × 125mm, with 5 very wavy-edged lobes, dark red (59A) in bud, opening strong purplish red (61B) inside and out; dark purple (79A) basal blotch extending upwards c.15mm on all the lobes and covering about a quarter of the dorsal lobe. Lvs elliptic, 75-125 × 25-50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and dark yellowish green (139A) above, hairless. Shrub 3.0 × 1.8m in 20 years (in crowded, shaded conditions). Mid May.

(a) **'Carroll's Brocaded Pillow'**

cv. Evergreen azalea: 'Princesse Maud' (s) × 'Kirin'. H: C. Carroll (1987), G: C. Carroll (1990), N: C. Carroll (2004), I: Arborglen Pty Ltd, REG: Carroll's Country Gardens Nursery (2005). Fls hose-in-hose, 2/truss, funnel-shaped, 10 × 20mm, with 11 lobes in all, lobes smooth-edged, vivid purplish red (57B) in bud and on inside when open, unmarked; filaments and style also purplish red. No normal calyx present. Lvs ovate, 25 × 12mm, margins flat, obtuse at apex, hairy. Shrub to 1.0m wide in 6 years. Spring.

(Nomenclatural standard: colour print provided by the registrant (WSY0070186)).

(r) **'Charis'**

cv. Elepidote rhododendron: 'Duke of York' (s) × 'Lem's Monarch'. H: H. Hachmann (1990), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: H. Hachmann (2005). Fls 9-12/rounded, upright truss, broadly funnel-shaped, 70-75 × 100-120mm, with 5 slightly wavy-edged lobes, buds strong red (53C), opening inside strong/light purplish pink (63C/D), edges of lobes on inside strong purplish red (63A-B); outside strong purplish red (58B), veins vivid purplish red (61C); dorsal lobe markings of dark red (187C); stamens 10-12, filaments pale purplish pink (62D), anthers strong purplish red/strong reddish purple (72A/B); style pale greenish yellow (1D), stigma strong red (46A); ovary greenish, silvery hairy. Calyx 1-2mm long, whitish green, edged with red-purple. Lvs elliptic, 120-180 × 48-55mm, margins slightly wavy, matt light green above with yellowish green venation, rounded to slightly oblique at base, apiculate at apex, with sparse brownish hairs below. Shrub c.1.7 × c.2.0m in 10 years; habit broadly upright. Mid to late May. Fls long-lasting. (Nomenclatural standard: colour print provided by registrant (WSY0070133)). Published in Hachmann cat. 2004-2008 (2004), p.38).

(r) **'Charismatic'**

cv. Elepidote rhododendron: 'Lem's Cameo' (s) × 'Golden Star'. H: H. O'Rourke (1990), G: H. O'Rourke (1999), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 17/ball truss, funnel-shaped, 50 × 80mm, with 7 wavy-edged lobes, strong purplish red (54A) in bud, opening with a yellow tube, deep purplish (54B) lobes and a red eye, outside strong purplish red (54A); a few red spots on the dorsal lobe. Calyx 12mm long, yellow and pink. Lvs oblong, 140 × 65mm, margins decurved, glossy green above, cuneate at base, acute at apex, hairless. Shrub 1.4 × 1.3m in 13 years. October (Australia).

(r) **'Charlotte Rose Newman'**

cv. Elepidote rhododendron: 'Lem's Tangerine' (s) × *lacteum*, Lem's form. H: L.L. Newcomb (1985), G: L.L. Newcomb (1992), N: L.L. Newcomb (2004), I: Hammond Nursery (2006), REG: E.F. Newcomb (2005). Fls 19-27/tall, domed truss, funnel-campanulate, 55 × 75mm, with 5 wavy-edged lobes, strong red (53B) in bud, opening inside strong red (53D), changing to deep pink (52B) in centre and light yellowish pink (36A) in the throat, with two small flares at base of dorsal lobe, the flares consisting of deep red (185A) and strong red (50A); outside strong red (51A) with strong red (53C) stripes on centre of lobes and pale yellowish pink (36D) at base; slightly scented. Calyx with 5 irregular lobes, c.13-50mm long, inside strong red (51A) edged deep pink (52C), the throat light yellowish pink (36); outside similar but slightly lighter. Lvs elliptic, 90-160 × 30-75mm, acute at apex, rounded at base, margins downcurved, dull and moderate

olive green (147A) above, moderate yellow-green (146D) below, hairless. Shrub 2.4 × 2.1m in 19 years. Mid April.

(r) **'Christina Dee'**

cv. Elepidote rhododendron: 'Casanova' (s) × 'Wyandanch Pink'. H: J. Doppel, S: J. Doppel (1995), G: J. Doppel (2002), N: J. Doppel (2004), REG: J. Doppel (2005). Fls 15/ball truss, broadly funnel-shaped, 40 × 50mm, with 5 wavy-edged lobes, coral pink in bud, opening pinkish cream and fading to cream; outside slightly darker than inside; large area of discrete dorsal spotting - spots light maroon at first, fading to tan. Lvs oblong, 120 × 40mm, broadly acute at apex, oblique at base, margins downcurved, semi-glossy and mid green above, hairless. Shrub 1.1 × 0.9m in 10 years. Early May. (Seed and pollen fertile; tolerates winter sun and wind rather well).

(r) **'Christopher Loder'**

cv. Elepidote rhododendron: *fortunei* (s) × *griffithianum*. G: M.L.A. Robinson (2002), N: M.L.A. Robinson, I: Loder Plants (2006), REG: M.L.A. Robinson (2005). Fls 12/truss, funnel-campanulate to tubular funnel-shaped, 65-75 × 105-115mm, light purple (76A) in bud, mature flowers very light purple (76B) on lobes inside, shading through very pale purple (76C) to very pale purple (76D) in the throat, with two small areas of pale greenish yellow (2D) on dorsal ribs (c.15 × 3mm); outside with lobes very light purple (76B) shading to very pale purple (76C) near calyx, unmarked; strongly scented. Calyx 2mm long, greenish purple. Lvs oblanceolate/oblong, 150-200 × 40-60mm, margins flat, glossy above, obtuse at base, acute at apex, hairless; petioles deep purplish red (61A) at darkest. Shrub 1.4 × 2.0m in 9 years. Mid April. Differs from other Loderi Group plants in its clear unfading pink corollas. (Loderi Group. Nomenclatural standard: colour print provided by the registrant (WSY0060738)). Etymology: named for the owner of Loder Plants nursery who is a descendant of the original family after which the Loderi Group is named. The original plant was found in Leonardslee Garden (Sussex, England), but has now been lost).

(a) **'Cinzia Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls double, 2-5/hemispherical truss, 51 × 37mm, with 5 wavy to very wavy-edged outer lobes, light yellow-green (144D) in bud, opening white throughout; three of outer lobes with light yellow-green (144D) markings; stamens absent; pistil exceeding corolla, style light yellow-green (145D), stigma light yellow-green (144D). Lvs elliptic, 28 × 14mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (137B) above and strong yellow-green (144B) below when mature; petioles 5mm long. Shrub 1.7 × 2.3m in 60 years (regularly pruned). May. (Etymology: named for the nominant's wife).

(r) **'City of Burnie'**

cv. Elepidote rhododendron: 'Noyo Chief' (s) × Matador Group. H: H. O'Rourke (1980), G: H. O'Rourke (1988), N: H. O'Rourke (1988), REG: H. O'Rourke (2005). Fls 20-21/flat-topped truss, funnel-campanulate, 50 × 60mm, with 5 wavy-edged lobes, red in bud, opening strong red (53C) inside and out. Calyx 3mm long, green. Lvs lanceolate, 150 × 45mm, matt green above, cuneate at base, acute at apex, hairy (hairs fawn, lost with age). Shrub 3.0 × 2.5m in 24 years. September to October (Australia).

(r) **'Clara Cluck'**

cv. Elepidote rhododendron: (*makinoi*, white-flowered × *luciferum*) (s) × *lanatum*. H: C.A. Lehmann, G: T. Stein (2005), N: T. Stein, REG: T. Stein (2005). Fls 10-12/ball to domed truss, funnel-campanulate, 70mm wide, with 5 slightly wavy-edged lobes, buds strong purplish pink (62A), opening inside pale purplish pink (62D), outside light purplish pink (62C); numerous, fine, strong purplish red (59D) spots on dorsal lobe (showing through on outside). Calyx 2-3mm long, pale yellowish pink (36D). Lvs oblong, 100 × 30mm, margins decurved, glossy green above, cuneate at base, broadly acute at apex, upper surface with an indumentum at first but later lost, the indumentum below pale yellow (161C) at first, aging to brownish orange (165B). Shrub 0.7 × 0.8m in 8 years. April to May. (Nomenclatural standard: colour print provided by the registrant (WSY0070084). Etymology: Clara Cluck is the name of the registrant's grand-daughter: "cluck" being an anglicised version of the Danish pet name for her ("kluk").

(r) **'Cocheco Big George'**

cv. Elepidote rhododendron: 'Mrs Charles S. Sargent' (s) × 'Dexter's Purple'. H: J.B. Parks (1985), G: J.B. Parks (1991), N: J.B. Parks (1995), REG: J.B. Parks (2005). Fls 13-15/ball truss, broadly funnel-shaped, 65 × 80mm, with 6-7 wavy-edged lobes, light purple (75A) in bud, opening inside light purple (75B) with the centre of the lobes slightly paler than the edges, the base light yellow (163D); outside light purple (75A-B). Calyx 3mm long, pale yellow-green (193A). Lvs elliptic, 225 × 85mm, broadly acute at apex, oblique at base, margins flat, glossy and moderate olive green (137A) above, hairless. Shrub 2.4 × 2.4m in 10 years. Late May. (First published in *The Rosebay* 30(2), Fall 2002/Winter 2003).

(r) **'Cocheco Honey Glacé'**

cv. Elepidote rhododendron: 'Ice Cube' (s) × white-flowered Consolini hybrid. H: J.B. Parks (1987), G: J.B. Parks (1992), N: J.B. Parks (1998), REG: J.B. Parks (2005). Fls 15/domed truss, broadly funnel-shaped, 45 × 65mm, with 5 wavy-edged lobes, moderate purplish pink (186D) in bud, opening inside near white (paler than pale orange-yellow (159C)) or pale yellowish pink (159D); outside between yellowish white (155D) and white; the central dorsal lobe is so heavily spotted as to appear almost a solid colour ranging from light yellow (163D) at margin to greyish reddish orange (174B) at the base; slightly scented.

Lvs elliptic, 160 × 65mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137B) to moderate yellow-green (137C) above, hairless. Shrub 1.5 × 1.8m in 13 years. Early June. (First published in *J. Am. Rb. Soc.* 57(4):195-201 (2003)).

(r) **'Cocheco Purple Butterfly'**

cv. Elepidote rhododendron: 'Roseum Elegans' (s) × 'Olin O. Dobbs'. H: J.B. Parks (1983), G: J.B. Parks (1990), N: J.B. Parks (1993), REG: J.B. Parks (2005). Fls 12/domed truss, broadly funnel-shaped, 75 × 55mm, with 5 flat-edged lobes, strong reddish purple (78B) in bud, opening inside between strong reddish purple (78B) and light purple (78C), with a very large, dark greenish yellow (152C) dorsal blotch speckled with greyish reddish orange (177C) dots; outside light purple (75B). Lvs elliptic, 145 × 55mm, semi-glossy and dark to moderate yellowish green (139A-B) above, hairless. Shrub 1.8 × 1.5m in 14 years. Late May. (First published in *The Rosebay* 30(2), Fall 2002/Winter 2003).

(r) **'Cody'**

cv. Elepidote rhododendron: 'Wind River' (s) × 'Janet Blair'. H: J. Barlup (1996), G: J. Barlup (2000), N: J. Barlup (2004), REG: J. Barlup (2005). Fls 15/ball truss, broadly funnel-shaped, 55 × 80mm, with 5 very wavy-edged lobes, strong purplish red (60C) in bud, opening inside pale yellow (20D), overlaid with moderate purplish pink (62B), deepening at edges to strong purplish pink (62A), the dorsal lobe tinged light yellow (11B) c.40mm from the base; outside pale yellow (20D) blending with moderate purplish pink (62B), edges and midribs strong purplish pink (62A); dark greenish yellow (152D) dorsal flare, extending c.45mm from the base, speckled with discrete, dark red spots. Calyx 25mm long, a blend of pale yellow (20D) and moderate purplish pink (62B). Lvs elliptic, 120 × 45mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.2 × 0.9m in 8 years. Late April.

(a) **'Colombè'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 1-2/truss, 53 × 47mm, with 5 wavy-edged lobes, vivid reddish orange (42A); three lobes with strong red (46A) markings; stamens 5, not exerted from the corolla, anthers dark red (59A), filaments white shaded vivid red (52A); pistil shorter than the stamens, style light yellowish pink (26D) shaded strong red (50A), stigma moderate reddish orange (34A). Lvs elliptic, 39 × 16mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (146C) below when mature; petioles 4mm long. Shrub 1.2 × 2.1m in 50 years. May to June. (Etymology: local name for a place near Cannero Riviera (where the Carmine nursery is located)).

(r) 'Cool Customer'

cv. Elepidote rhododendron: 'Janet Blair' (s) × 'Golden Belle'. H: H. O'Rourke (1990), G: H. O'Rourke (1999), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 10-12/loose truss, funnel-shaped, 55 × 80mm, with 5 flat-edged lobes, cream in bud, opening yellow in the throat paling to cream, the lobes tinged mauve, outside cream; large flare of green streaks. Calyx 4mm long, green. Lvs elliptic, 110 × 50mm, margins decurved, matt green above, truncate at base, acute at apex, hairless. Shrub 1.0 × 1.2m in 14 years. October to November (Australia).

(r) 'Crème Brûlée'

cv. Elepidote rhododendron: 'Mrs J.G. Millais' (s) × Inamorata Group. H: E.G. Millais (1963), G: E.G. Millais, N: Millais Nurseries (2005), REG: D.G. Millais (2005). Fls 12/lax, domed truss, open funnel-shaped, 30 × 80mm, with 5 wavy-edged lobes, pinkish in bud, opening perhaps creamier than white 155D, outside creamy white (155D); prominent speckling on dorsal lobes of deep greenish yellow (153A) fading to strong yellow (153D). Lvs elliptic, 120 × 50mm, margins flat, matt green above when mature (glossy when young), hairless. Shrub 1.5 × 1.5m in 10 years. Late May to early June. AM 2005. (Nomenclatural standard: herbarium specimen provided by the registrant (WSY0066123)).

(a) 'Cumberland Cheer'

cv. Deciduous azalea: *arborescens* (s) × *cumberlandense*. H: J.E. Schild (1986), G: J.E. Schild (1993), N: J.E. Schild (2003), I: Schild Azalea Gardens & Nursery (2006), REG: J.E. Schild (2005). Fls 6-10/flat truss, tubular funnel-shaped, 65 × 65mm, with 5 flat-edged lobes, buds vivid red (45B) with white, glandular setae along midribs and tube, opening inside moderate red (47A) with a vivid reddish orange (N30A) dorsal blotch; outside strong red (47B) with white, glandular setae along tube and lobes; style moderate red (47A); filaments deep pink (48B); stigma moderate red (N34A); anthers moderate orange-yellow (164B); strongly scented. Calyx c.3mm long, light green. Lvs obovate, 65 × 25mm, broadly acute at apex, cuneate at base, margins flat, glossy and strong yellow-green (144B) above, strong yellow-green (N144C) below, hairs on margins and midvein below (hairs white, aging to light green). Open shrub, 1.8 × 1.2m in 10 years. June. (Published in *The Azalean* 27(2):31 (2005)).

(a) 'Cumberland Coral'

cv. Deciduous azalea: *arborescens* (s) × *cumberlandense*. H: J.E. Schild (1986), G: J.E. Schild (1993), N: J.E. Schild (2003), I: Schild Azalea Gardens & Nursery (2006), REG: J.E. Schild (2005). Fls 7-10/domed truss, tubular funnel-shaped, 55 × 50mm, with 5 flat-edged lobes, buds strong reddish orange (40C), the midveins pale yellow (11D), opening inside moderate yellowish pink (39C) with pale yellow (11D) down the centre of each lobe and a vivid yellow (16A) blotch in the centre of the dorsal lobe; outside strong yellowish pink (43D) along margin, pale yellow (11D)

along midrib to calyx, with many glandular setae along the tube and lobes; filaments pale greenish yellow (2D); style strong red (47B); stigma strong red (46A); strongly scented. Calyx 3mm long, light yellow-green (142C). Lvs obovate, 65 × 40mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and moderate olive green (137B) above, moderate yellowish green (N138C) below, hairs on margins and on midvein below (hairs light greenish white aging to greenish white). Dense shrub, 1.8 × 1.2m in 10 years. June. (Published in *The Azalean* 27(2):32 (2005)).

(a) 'Cumberland Fire'

cv. Deciduous azalea: *cumberlandense* (s) × *arborescens*. H: J.E. Schild (1986), G: J.E. Schild (1994), N: J.E. Schild (1996), I: Schild Azalea Gardens & Nursery (2006), REG: J.E. Schild (2005). Fls 6-8/domed truss, narrowly tubular funnel-shaped, vivid reddish orange (43B) in bud, opening strong red (47B), with a vivid reddish orange (N30B) dorsal blotch; outside deep pink (48A), the tube with white hairs and eglandular deep pink (48B) hairs; filaments and style deep yellowish pink (43C); stigma dark greyish reddish brown (200A); not scented. Calyx 5mm long, yellow-green. Lvs obovate, 50 × 30mm, broadly acute at apex, cuneate at base, margins upcurved, semi-glossy and moderate olive green (137A) above, moderate yellow-green (137C) below, hairs on margins and on midvein below (hairs white, aging to light green). Shrub 0.6 × 0.6m in 10 years, slow-growing, with many horizontal branches. June. (Published in *The Azalean* 27(2):31 (2005)).

(a) 'Cumberland Gold'

cv. Deciduous azalea: *cumberlandense* (s) × *arborescens*. H: J.E. Schild (1986), G: J.E. Schild (1994), N: J.E. Schild (2003), I: Schild Azalea Gardens & Nursery (2006), REG: J.E. Schild (2005). Fls 6-8/domed truss, narrowly tubular funnel-shaped, 45 × 60mm, with 5 flat-edged lobes, brilliant greenish yellow (4A) tipped strong orange (N25A) in bud, opening inside brilliant yellow (13B), tinged vivid reddish orange (N30B) at margins, the dorsal lobe covered almost entirely with a vivid yellow (13A) blotch; outside brilliant yellow (12B), tinged deep yellowish pink (43C) on the lobes, with light greenish yellow (1C) hairs and a few glandular setae; filaments strong reddish orange (N34B); style strong red (39A); stigma strong red (46A); not scented. Calyx 5mm long, light green. Lvs elliptic, 60 × 25mm, acute at apex, cuneate at base, margins flat, glossy and dark yellowish green (136B) above, dull and moderate yellowish green (137D) below, hairs on margins and midvein below (hairs white, aging to yellow-white). Dense, compact shrub, 0.9 × 0.9m in 10 years. June. (Published in *The Azalean* 27(2):31 (2005)).

(r) 'Daddy Dan'

cv. Elepidote rhododendron: (Fabia Group × *bureauvii*) (s) × 'The Honourable Jean Marie de Montague'. H: D.A. Griswold (1970), G: Griswold Nursery (1980), N: Griswold Nursery, I: Griswold Nursery (2005), REG: L. Malland (2005). Fls 9/lax truss, open funnel-

campanulate, 50 × 65mm, with 6 wavy-edged lobes, deep purplish pink (61D) in bud, opening very pale purple (69C) throughout, with sparse spotting (closest to 61D) on lower portion of dorsal lobe. Lvs narrowly elliptic, 125 × 25mm, moderate olive green (137A) above, acute at apex, hairless. Shrub 1.8 × 1.8m in 35 years. Mid April.

(r) **'Dalton's Sunrise'**

cv. Elepidote rhododendron: 'Blimey Limey' (s) × 'Gallacci's Gold'. H: M. Stockman (1992), G: M. & H. Stockman (2000), N: S. Zachman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 11-13/domed truss, open funnel-shaped, 65 × 80mm, with (6-)7 wavy-edged lobes, buds strong yellowish pink (31C) shading to deep yellowish pink (47C) at tips, opening inside and out pale yellow (20C) with moderate yellowish pink (38B) along lobe sides and edges; with an inconspicuous light orange-yellow dorsal blotch. Calyx lobes irregular, c.5-20mm long, light orange-yellow (22C), edged deep yellowish pink (38B). Lvs elliptic, 120 × 40mm, acute at apex, cuneate at base, margins flat and downcurved, dull and moderate olive green (137B) above, moderate yellow-green (147C) below, hairless. Shrub 1.5 × 1.5m in 12 years. Mid May.

(r) **'Darren Arblaster'**

cv. Elepidote rhododendron: 'Goldsworth Orange' (s) × 'Bergie Larson'. H: G. Arblaster (1983-1984), G: G. Arblaster (1988), N: Mrs J. Arblaster (2005), REG: Mrs J. Arblaster (2005). Fls 8-10/fairly loose, open-topped truss, broadly tubular-campanulate, 68 × 78mm, with 5 slightly wavy-edged lobes, deep yellowish pink (47C) in bud, opening inside deep pink (47D) flushed in the centre with light orange (29B); outside strong yellowish pink (33C) shading to deep pink (47C) at lobe edges; inconspicuous spotting of strong reddish orange (42B) on all but ventral lobe; stamens 10, short (held well within the corolla); style pale, suffused green just below the stigma. Calyx 15-35mm long (very unevenly lobed), yellow-orange with strong yellowish pink (33C) streaks. Lvs elliptic/oblanceolate, up to 150 × 54mm, margins flat to slightly wavy, matt green above, cuneate at base, acute and mucronate at apex, with a few minute, glandular hairs on the petiole, otherwise hairless. Slow-growing shrub, 1.4 × 1.35m in 17 years, often with reddish linear bud scales. Late June to early July. (Nomenclatural standard: colour print provided by the registrant (WSY0071000). Etymology: named for a son of the registrant).

(r) **'Dartmoor Rose'**

cv. Lepidote rhododendron: *glaucophyllum* 'Len Beer' (s) × *charitopes* subsp. *tsangpoense*. H: B.N. Starling (1983), G: B.N. Starling (1988), N: B.N. Starling (1988), I: B.N. Starling (1994), REG: B.N. Starling (2005). Fls 4/lax truss, out to down-facing, 15-20 × 18-20mm, campanulate, with 5 flat-edged lobes, buds moderate pink (49B), opening light pink (50D) inside and out; unmarked; filaments white, anthers dark brown; style 20mm long. Calyx 7-9mm long,

green, covered with white scales; pedicels to 35mm long, reddish. Lvs elliptic to oblong, 23-40 × 8-15mm, margins flat, matt green above, broadly tapered at base, pointed (or sometimes rounded) at apex, with small grey white and large black scales below. Compact shrub, 0.7 × 1.0m in 20 years. Mid May. (Nomenclatural standard: colour slide provided by the registrant (WSY0070088). Etymology: named for where it was raised (the north-east corner of Dartmoor, UK) and for its flower colour).

(r) **'D.D. Noyce'**

cv. Elepidote rhododendron: Cornubia Group (s) × 'Coronation Day'. H: D.J. Dosser (1990), G: D.J. Dosser (2000), N: D.J. Dosser (2005), REG: D.J. Dosser (2005). Fls 14/domed truss, funnel-shaped, 60 × 90mm, with 6 wavy-edged lobes, deep purplish pink (68A) in bud, opening inside light purplish pink (68D), the centre of each lobe and the upper part of the throat paler, outside moderate purplish pink (68C); with rays of deep purplish pink (68A) spots in the dorsal throat. Lvs oblong, 225 × 65mm, flat, matt green above, attenuate at base, mucronate at apex, hairless. Shrub 3.0 × 2.0m in 14 years. November to December (Australia). (Nomenclatural standard: colour print provided by the registrant (WSY0060824)).

(r) **'Dean Barber'**

cv. Lepidote rhododendron: *keiskei* 'Yaku Fairy' (s) × *keiskei*. H: D. Barber (c.1976), G: D. Barber (c.1981), N: J.K. Weagle, I: Rhodoland Nursery, REG: J.K. Weagle (2005). Fls 4-9/lax truss (2-3 fls/bud, 2-3 buds/shoot), open funnel-shaped, 35mm long, with slightly wavy-edged lobes, brilliant to light yellow-green (154C-D) in bud, opening throughout between pale greenish yellow (2D) and light yellow-green (150D), with a slight flare of brilliant yellow-green (150B) spots - about 7 × 7mm and emanating from the centre onto the three dorsal lobes; stigma c.30mm long, mostly pale yellow-green, some tinged red. Calyx 2.5mm long, greenish yellow, some tinged red. Lvs narrowly elliptic to lanceolate, acute at apex, rounded at base, lamina flat; in winter dark red (between 187A & 187B), in summer close to moderate olive green (146A), with large brown scales typical of *keiskei* below. Compact, mounded shrub, 0.5 × 1.4m in 23 years. Late to mid May.

(r) **'Diane's Delight'**

cv. Elepidote rhododendron: 'Peter Alan' (s) × 'Anah Kruschke'. H: J. Doppel (1992), G: J. Doppel (1998), N: J. Doppel (2004), REG: J. Doppel (2005). Fls 20/conical truss, open funnel-shaped, 50 × 65mm, with 5 slightly wavy-edged lobes, dark reddish purple in bud, opening inside reddish purple, outside slightly darker than inside; prominent, black, dorsal spotting. Lvs obovate, 115 × 40mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and dark green above, hairless. Shrub 1.4 × 1.2m in 10 years. Mid May. (Both pollen and seed fertile).

(r) 'Doreen Johnson'

cv. Elepidote rhododendron: 'Rigginslinger' (s) × (*yakushimanum* × 'One Thousand Butterflies'). H: J.D. Winberg (1993), G: J.D. Winberg (1999), N: J.D. Winberg, REG: J.D. Winberg (2005). Fls 13/truss, open campanulate, 55 × 40mm, with 5 wavy-edged lobes, deep purplish pink (66C) in bud, opening strong purplish pink (63C). Lvs oblong, 125 × 40mm, margins upcurved, dull and moderate olive green (137A) above, with discrete hairs below. Shrub 0.4 × 0.5m in 6 years. Late April to May.

(a) 'Drapa'

cv. (Evergreen azalea: 'Squirrel' (s) × 'Chippewa'). H: Holger Hachmann (1990), G: Holger Hachmann (1994), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls single, 2-3/truss, broadly funnel-shaped, 30-35 × 40-45(-55)mm, with a longish tube and with 5 slightly wavy-edged lobes, buds strong purplish red (58B/C), opening deep purplish pink (61D) inside and out; 3 dorsal lobes with sparse markings of strong purplish red (60D); stamens 5(-6), filaments strong purplish pink (to light purplish pink (62A(-C))), anthers dark red (187B); style strong purplish pink (62A), stigma moderate red (183C); ovary greenish, silvery hairy. Calyx 3-4mm long, brilliant yellow-green (149C), silvery hairy. Lvs ovate, 25-35 × 13-17mm, margins flat, mid green and glossy above, cuneate at base, apiculate at apex, with brownish hairs; lasting well in winter. Shrub c.0.3 × c.0.7m in 10 years, of compact, flat-rounded habit. Early to late June (i.e. late). (Nomenclatural standard: colour print provided by the registrant (WSY0070168). Published in Hachmann cat. 2004-2008 (2004), p.164.

(r) 'Dreambaby'

cv. Elepidote rhododendron: 'Dreamland' (Waterer) (s) × 'Pretty Baby'. H: S. Imrie (1998), G: S. Imrie (2002), N: S. Imrie (2005), I: Lake Tapps Rhododendron Nursery (2007), REG: S. Imrie (2005). Fls 9/ball truss, campanulate, 40 × 65mm, with 5 very wavy-edged lobes, vivid red (52A) in bud, opening inside light pink (50D), outside strong pink (52D); no markings. Calyx lobes to 40mm long, many with recurved margins, deep pink (52B) streaked with pale yellowish pink (36D). Lvs elliptic, 90 × 40mm, broadly acute at apex, cuneate at base, margins downcurved, dull and dark yellowish green (139A) above, hairless. Shrub 1.2 × 1.2m in 7 years. Late May.

(r) 'Early Light'

cv. Elepidote rhododendron: 'Vinecrest' (s) × ('Janet Blair' × ('Donna Hardgrove' × 'Weston's Hardy Yellow')). H: A. Glassman (1996), G: J. Doppel (2001), N: J. Doppel (2004), REG: J. Doppel (2005). Fls 8/ball truss, saucer-shaped, 50 × 90mm, with 7 slightly wavy-edged lobes, light brick red in bud, opening inside light yellow, shading slightly darker yellow in the throat; outside similarly coloured; unmarked; slightly scented. Lvs oblong, 115 × 45mm, broadly acute at apex, oblique at base, margins flat, semi-glossy and mid green (slightly olive) above, hairless. Shrub 0.9 × 1.4m in 10 years. Early May. Syn:

'Yellow Fellow'. (Pollen fertile but usually seed sterile).

(r) 'Eastern Ring of Fire'

cv. Elepidote rhododendron: 'Vulcan' (s) × ('Ted's Yellow' × *fortunei*). H: D.B. Johnson (1988), G: D.B. Johnson (1995), N: D.B. Johnson (2005), REG: D.B. Johnson (2005). Fls 10/domed truss, open funnel-shaped, 50 × 75mm, with 5 very wavy-edged lobes, vivid reddish orange (41A) in bud, opening pale yellow (20C) inside and out, with margins of deep yellowish pink (39B) - 5-6mm wide inside and 25mm wide outside; brilliant yellow (11A) dorsal spotting. Lvs elliptic, 115 × 45mm, broadly acute at apex, rounded at base, with wavy margins, bronze at first, slowly becoming moderate yellowish green (138B) above, hairless. Dense shrub, 1.1 × 1.8m in 10 years. Mid May.

(r) 'Ed Brown'

cv. Elepidote rhododendron: 'Scintillation' (s) × ('Golden Star' × ((Dido Group × ('Chlorops' × *lacteum*)) × 'Golden Star')). H: P. Brown (1994), G: P. Brown (2000), N: P. Brown (2000), I: Whispering Pines Nursery (2005), REG: P. Brown (2005). Fls 10-12/domed truss, broadly funnel-shaped, 50 × 90mm, with 7 flat-edged lobes, light yellowish pink (26D) in bud, opening inside light yellow-green (150D); outside pale yellow (8D). Calyx 3mm long, light yellow-green (145C). Lvs elliptic, 150 × 65mm, dull and moderate olive green (137A) above, hairless. Shrub 1.2 × 1.2m in 10 years. Mid May.

(r) 'Elegant Air'

cv. Elepidote rhododendron: 'Lem's Cameo' (s) × 'Roza Stevenson'. H: H. O'Rourke (1994), G: H. O'Rourke (2002), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 7/loose truss, funnel-shaped, 60 × 90mm, with 7 wavy-edged lobes, cream in bud, opening cream, deeper in the throat which has a small red flare; outside cream. Calyx 35mm long, cream. Lvs elliptic, 100 × 40mm, margins wavy, glossy green above, rounded at base, obtuse at apex, hairless. Shrub 1.3 × 1.2m in 10 years. October (Australia).

(a) 'Elfie'

cv. Evergreen azalea: 'Squirrel' (s) × 'Chippewa'. H: Holger Hachmann (1990), G: Holger Hachmann (1994), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls single, 2-3/truss, funnel-shaped, 30 × 35-40(-45)mm, with 5 slightly wavy-edged lobes, buds deep yellowish pink (46D), opening deep yellowish pink (46D) shaded vivid red (46C) inside and out; dorsal lobes slightly spotted with vivid red (46C), but the marks very inconspicuous; stamens 6, filaments strong red (46A), anthers dark red (187B); style deep/strong pink (52C/D), stigma deep red (53A); ovary greenish, silvery hairy. Calyx 2-4mm long. Lvs narrowly elliptic, 24-30 × 5-10mm, margins flat, matt to mid light green above, cuneate at base, pointed to slightly apiculate at apex, with whitish hairs. Shrub c.0.3 × 0.8m in 10 years, with a cushion-shaped, prostrate habit. Early to mid June (i.e. late). (Nomenclatural standard:

colour print provided by the registrant (WSY0070162). Published in Hachmann cat. 2004-2008 (2004), p.165).

(r) **'Eliska'**

cv. Elepidote rhododendron: 'Rotgold' (s) × 'Goldkrone'. H: H. Hachmann (1994), G: H. Hachmann (1999), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls up to 15/rather loose, flattened ball truss, almost rotate, the 5(-6) flat to slightly wavy-edged lobes reaching almost to the base of the corolla, 30-35 × 40-45(-50)mm, buds brilliant yellow (7B), tinged orange-red, opening inside and out light greenish yellow (4B); dark red (59A) marks in the throat; stamens 10, filaments light yellow-green (150D), anthers moderate orange-yellow (165C); style brilliant yellow-green (150B/C), stigma brownish orange (165B), ovary greenish, silvery hairy. Calyx apparently absent. Lvs ovate, 40-65 × 25-42mm, margins flat, matt fresh green above, rounded at base, apiculate at apex, hairless. Shrub c.0.6 × c.0.9m in 10 years, with a compact, rounded habit. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070152). Published in Hachmann cat. 2004-2008 (2004), p.151).

(r) **'Ella Carpenter'**

cv. Elepidote rhododendron: parentage unknown. H: unknown, G: E. Carpenter (1970), N: Griswold Nursery, I: Griswold Nursery (2005), REG: L. Malland (2005). Fls 5/lax truss, broadly funnel-shaped, c.50 × c.65mm, with 5-6 wavy-edged lobes, vivid reddish orange (33A) in bud, opening light greenish yellow (4B) throughout, with vivid orange (28B) spotting on the three dorsal lobes. Lvs elliptic, 100 × 50mm, broadly acute at apex, margins flat, moderate olive green (137B) above, hairless. Upright shrub, 1.5 × 1.5m. May.

(r) **'Enbu'**

cv. Elepidote rhododendron: *arboreum* (s) × ((Carmen Group × (*williamsianum* × *fortunei*)) × *simiarum*). H: K. Terai (1997), G: K. Terai (2002), N: K. Terai (2005), REG: K. Terai (2005). Fls 11/hemispherical truss, open funnel-shaped, 35 × 60mm, with 5 flat-edged lobes, deep red (60A) in bud and both inside and out when open; unmarked; colour lasting well in the sun. Lvs elliptic, 80-100 × 25-35mm, margins slightly decurved, glossy green above, obtuse at base, acute at apex, hairy below (indumentum light brown when young and when mature). Shrub 15 × 50cm in 7 years. Mid May. (Nomenclatural standard: colour print provided by registrant (WSY0070571)).

(r) **'Enrico'**

cv. Lepidote rhododendron: 'Sacko' (s) × *calostrotum* subsp. *riparioides*. H: H. Hachmann (1990), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 3-4/individual truss, gathered in a dense ball inflorescence at the shoot tips, broadly funnel-shaped, c.20 × 32-36mm, with 5 wavy-edged lobes,

inside light violet (88C), slightly tinged purple, outside similarly coloured but midveins tinged light purple (80C); dorsal lobe with sparse markings of moderate purple (79D) near the throat; stamens 10, filaments light purple (85A/B), anthers dark red (187A); style strong purplish red (71B), stigma dark red (187B). Calyx 5-7mm long, strong/light yellow-green (145A/B). Lvs narrowly elliptic, 18-22 × 6-10mm, margins flat, glossy above, rounded at base, apiculate at apex, scaly (young scales light yellow-green (142C/D), mature scales moderate reddish brown (174A)). Shrub 25 × 40cm in 10 years, with a dense, flat to rounded habit. Early to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070154). Published in Hachmann cat. 2004-2008 (2004), p.118).

(r) **'Enumclaw Chinook'**

cv. Elepidote rhododendron: 'Anna' (Lem) (s) × 'Elizabeth de Rothschild'. H: W.B. Wilkes (1982), S: R. Wieder (1987), G: R. Wieder (1990), N: J.D. Anderson (2004), I: Anderson Garden (2005), REG: J.D. Anderson (2005). Fls 16-20/ball truss, broadly funnel-shaped, 55 × 75mm, with 5 wavy-edged lobes, dark pink in bud, opening inside white with a small pink marking in the throat below one lobe; outside pink, becoming paler with age. Calyx c.3mm long, light green. Pedicel c.20mm long, green to dark pink. Lvs oblong, 125 × 50mm, broadly acute at apex, rounded at base, margins upcurved, dark green, variegated with yellowish green and light green streaks in the centre of the blade; hairless. Shrub 1.5 × 0.9m in 10 years. Mid May.

(r) **'Erica J.'**

cv. Elepidote rhododendron: *wardii* (Rock form), open-pollinated (s). H: E. Jago (1996), G: E. Jago (2001), N: E. Jago (2004), REG: E. Jago (2005). Fls 19/very dense ball truss, flat saucer-shaped, 40 × 70mm, with 5 wavy-edged lobes, deep pink in bud, opening porcelain pink and becoming white both inside and out, with a light green blotch on the dorsal lobe; slightly scented. Calyx 2mm long, pale green. Lvs elliptic, 100 × 50mm, margins decurved, obtuse at base, acute at apex, hairless. Shrub 1.36 × 1.05m in 8 years. Late October to mid November (New Zealand).

(a) **'Ester Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls hose-in-hose, 3-13/hemispherical truss, 33 × 23mm, with 5+5 flat or wavy-edged lobes, buds white shaded vivid purplish red (57B), opening throughout white shaded vivid purplish red (57C); three lobes with deep pink (48A) markings shaded with strong purplish red (63B); stamens 5, exerted from corolla, anthers moderate orange (167C), filaments white; pistil equalling or distinctly exceeding stamens, style white, stigma brilliant greenish yellow (2B). Lvs elliptic, 22 × 9mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (147A) above and

greyish yellow-green (148C) below when mature; petioles 4–8mm long. Shrub 1.7 × 1.0m in 60 years (regularly pruned). April. (Etymology: named for the aunt of the nominant).

(a) **'Ester della Valle di Casanova'**

cv. Evergreen azalea: parentage unknown. R: Gardeners of the Villa San Remigio (pre 1920), N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single, 1-3/truss, 64 × 48mm, with 5 flat-edged lobes, buds and open flowers strong red (53B) shaded red (N57A); three lobes with deep red (53A) markings; stamens 10, not exerted from the corolla, anthers moderate red (183C), filaments strong purplish red (67A); pistil distinctly exceeding stamens and exerted from the corolla, style strong purplish red (58B), stigma light greenish yellow (8C). Lvs lanceolate, 44 × 17mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (146D) below when mature; petioles 4mm long. Shrub 1.8 × 3.0m in c.100 years. May. (Etymology: named for the individual who donated the Villa San Remigio to the Piedmont Region as a public park).

(a) **'Eugenio Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-18/hemispherical truss, 49 × 28mm, with 5 very wavy-edged lobes, white or white shaded light purple (75B) in bud, opening a similar colour throughout; three lobes with deep purplish red (59B) markings; stamens 7-8, exerted from corolla, anthers greyish reddish brown (200B), filaments light purple (75B); pistil exceeding stamens, style white; stigma brilliant yellow (8A). Lvs lanceolate, 27 × 11mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (137A) above and moderate yellow-green (146B) below when mature; petioles 5mm long. Shrub 1.5 × 3.0m in 60 years (regularly pruned). April. (Etymology: named for the nominant's father).

(r) **'Eveline Pilkington'**

cv. Elepidote rhododendron: un-named seedling (Consolini no.279) (s) × un-named seedling (Consolini no.303). H: Mrs E. Pilkington (1976), G: Mrs E. Pilkington (1989), N: N. Beaudry (2005), REG: N. Beaudry (2005). Fls 13/conical truss, open funnel-shaped, 65 × 115mm, with 5-6 wavy-edged lobes, vivid purplish red (67B) in bud, opening inside vivid purplish red (67B) fading to very pale purple (69A) at edge, with a large dorsal blotch of strong greenish yellow (151A) speckles; outside strong purplish pink (68B); filaments yellowish white (155D); anthers light reddish brown (177B); pistil c.50mm long. Calyx c.5mm long, strong purplish pink (68B). Lvs oblong, 150 × 65mm, acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, moderate yellow-green (147C) below, hairless. Shrub 3.0 × 2.8m in 14 years. Early June.

(r) **'Exhibitionist'**

cv. Elepidote rhododendron: 'Lem's Cameo' (s) × 'Whitney's Orange'. H: H. O'Rourke (1990), G: H. O'Rourke (2000), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 12/loose truss, broadly funnel-shaped, 50 × 65mm, with 6 wavy-edged lobes, orange-yellow in bud, opening inside yellow, with lighter coloured lobes, outside orange-yellow, with yellow lobes; a small, red, central blotch in the throat. Calyx 2mm long, green. Lvs lanceolate, 100 × 40mm, margins wavy, matt green above, rounded at base, acute at apex, hairless. Shrub 2.0 × 1.0m in 15 years. Early November (Australia).

(a) **'Fiore di Pesco Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-5/truss, 27 × 33mm, with 5 flat-edged lobes, light purplish pink (55C) in bud and on opening, both inside and out, unmarked; stamens 5-6, exerted from corolla, anthers greyed brown (N199C), filaments pale purplish pink (55D); pistil exceeding stamens, style strong pink (54C), stigma light orange-yellow (24C). Lvs oblanceolate 20 × 8mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (146A) above and moderate yellow-green (146D) below when mature; petioles 3-4mm long. Shrub 0.5 × 1.5m in 60 years (regularly pruned). May. (Etymology: the flower resembles peach blossom ("fiore di pesco" in Italian)).

(r) **'Florteppich'**

cv. Elepidote rhododendron: 'August' (s) × *forrestii* Repens Group. H: H. Hachmann (1990), G: H. Hachmann (1995), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 5-6/rather loose truss, funnel-campanulate, 45 × 55mm, with 5 wavy-edged lobes, opening inside and out vivid red/deep pink (52A/B), fading with age to strong purplish red/deep purplish pink (54A/B); unmarked; stamens 10, filaments strong pink (52D), anthers brownish orange/moderate orange-yellow (164A/B); style light yellowish pink (29C), stigma dark red (187B). Calyx 1mm long, vivid red (52A). Lvs obovate, 55-60 × 30-35mm, convex, margins decurved, dark green and slightly glossy above, leathery, cuneate at base, apiculate at apex, hairless. Almost prostrate shrub, c.0.3 × c.1.1m in 10 years. Late April to early May (needing protection from spring frost). (Nomenclatural standard: colour print provided by the registrant (WSY0070153). Published in Hachmann cat. 2004-2008 (2004), p.126.).

(a) **'Franca Ciampi'**

cv. Evergreen azalea: parentage unknown. R: B. Salvatore (c.1920), N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single, 2-4/hemispherical truss, 68 × 43mm, with 5 lobes varying from flat to very wavy-edged, buds red-purple (N57B), opening inside and out red-purple (N57D); three lobes with red-purple (N57B) markings; stamens 9, not exerted, anthers greyish reddish orange (174B),

shaded moderate purple (79D), filaments strong red (53D); pistil distinctly exceeding stamens, style and stigma strong red (53C). Lvs lanceolate, 48 × 14mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (147B) below when mature; petioles 7-11mm long. Shrub 2.0 × 3.0m in 40 years. April to May. (Etymology: named for the wife of the President of Italy in 2004).

(a) **'Fredonia Surprise'**

cv. Deciduous azalea: *arborescens* (s) × *cumberlandensis*. H: J.E. Schild (1986), G: J.E. Schild (1994), N: J.E. Schild (2003), I: Schild Azalea Gardens and Nursery (2006), REG: J.E. Schild (2005). Fls 25-40/ball truss, tubular funnel-shaped, 50 × 50mm, with 5 wavy-edged lobes, strong purplish pink (62A) in bud, with some sparse, glandular setae on the midribs, opening moderate purplish pink (62B) with pinkish white at tips of lobe margins and a strong orange (24A) blotch in centre of dorsal lobe; outside with lobes light purplish pink (62C) and tube moderate purplish pink (62B) with many hairs and a few glandular setae; filaments strong yellowish pink (33C); styles vivid reddish orange (N30A); stigma deep red (60A); strongly scented. Calyx c.3mm long, light green. Lvs obovate, 75 × 40mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and strong green (132B) above, light bluish green (133C) below, with white hairs on leaf margins and on midvein below. Shrub 1.2 × 1.2m in 10 years. June. (Published in *The Azalean* 27(2):33 (2005)).

(r) **'Frentano'**

cv. Elepidote rhododendron: 'Scintillation' (s) × 'Germania'. H: H. Hachmann (1989), G: H. Hachmann (1995), N: H. Hachmann (2000), I: H. Hachmann (2001), REG: G. Stück (2005). Fls 12-18/rounded, rather loose truss, broadly open funnel-shaped, 65 × 85-105mm, with 5 slightly wavy-edged lobes, buds deep purplish pink (73A), opening inside and out light purplish pink (73C) with strong purplish pink (73B) edges to lobes, outside with deep purplish pink (73A) ribs and veins; dorsal lobe with deep greenish yellow (to strong greenish yellow) (153A(-C)) markings; stamens 10, filaments very pale purple (73D), anthers light purple (80D); styles light yellow-green (150D), stigma dark red (187C). Calyx 1-2mm long, strong yellow-green (144C). Lvs ovate, 130-180 × 60-75mm, partly auriculate, margins flat, mid green and slightly glossy above, rounded at base, cuspidate at apex, hairless. Shrub c.1.5 × c.1.7m in 10 years; habit broadly upright. May. A good foliage plant. (Nomenclatural standard: colour print provided by registrant (WSY0070134). Published in Hachmann cat. 2004-2008 (2004), p.45).

(a) **'Gardanina'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 1-2/truss, 43 × 26mm, with 5 flat to rarely wavy-edged lobes, buds red-purple (N66A), opening

red-purple (N66B) throughout; three lobes with deep red (60A) markings; stamens 5-6, exceeding corolla, anthers red-purple (N66B), filaments red-purple (N66B) shaded white near the ovary; pistil exceeding the corolla, style strong yellow-green (144B) shaded strong purplish pink (55B), stigma pale yellow (161C). Lvs elliptic, 41 × 18mm, margins flat, glossy above, acute at base and apex, hairy, strong yellow-green (144A) above and moderate yellow-green (146D) below when mature; petioles 4mm long. Shrub 1.8 × 2.4m in 50 years. April to May. (Etymology: local name for a place near Cannero Riviera (where the Carmine nursery is situated)).

(a) **'Ghetta Bruno'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls double, 2-3/truss, 72 × 48mm, with 5 outer, wavy-edged lobes, vivid purplish red (61C) in bud and throughout on opening, three of the outer lobes with deep red (60A) markings; stamens 5-10, not exceeding the corolla, anthers dark red (59A) shaded deep purplish red (59B), filaments red-purple (N57A); pistil longer than the stamens, style strong purplish red (58B), stigma dark red (59A). Lvs oblanceolate, 45 × 18mm, margins flat, glossy above, obtuse at base, acute at apex, hairy, moderate olive green (146A) above and moderate yellow-green (146C) below when mature; petioles 4mm long. Shrub 1.9 × 2.0m in 50 years. May. (Etymology: named for the best friend of the raiser).

(a) **'Gian Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 51 × 31mm, with 5 flat-edged lobes, deep purplish pink (67C) in bud and throughout on opening, with dark red (59A) markings shaded with deep red (60A) on three lobes; stamens 6-7, exerted from the corolla, filaments red-purple (N66D), anthers light olive grey (197B); pistil exerted from the corolla but shorter than the stamens, style strong purplish red (58C) shaded white, stigma moderate yellow-green (139C). Lvs elliptic, 32 × 16mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (146D) below when mature; petioles 5-6mm long. Shrub 1.6 × 2.0m in 60 years (regularly pruned). April. (Etymology: nickname of the nominant).

(a) **'Gianluigi Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-4/truss, 40 × 24mm, with 5 wavy to very wavy-edged lobes, white (155C) shaded deep purplish pink (68A) in bud, opening inside and out white (155C) shaded strong purplish pink (68B); unmarked; stamens 5, exerted from the corolla, anthers moderate orange-yellow (164B) shaded moderate reddish brown

(166B), filaments white; pistil exceeding the stamens, style light yellow-green (150D), stigma brilliant yellow-green (149B). Lvs obovate, 22 x 10mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (147C) below when mature; petioles 3-4mm long. Shrub 1.0 x 1.5m in 60 years (regularly pruned). May. (Etymology: Gianluigi Cavadini is the owner of the Cavadini nursery and grandson of Luigi Cavadini, founder of the nursery).

(r) **'Gingerine'**

cv. Elepidote rhododendron: 'Apricot Fantasy' (s) X 'Brandt's Tropicana'. H: C. Weeks (1991), G: C. Weeks (1998), N: C. Weeks, I: Westgate Gardens Nursery (2000), REG: C. Weeks (2005). Fls 15/slightly lax truss, campanulate, 65 x 80mm, with 7 wavy-edged lobes, vivid reddish orange (32A) in bud, opening inside strong reddish orange (32B), the three dorsal lobes marked with vivid reddish orange (30A) and vivid orange-yellow (23A); outside strong reddish orange (32B). Lvs elliptic, 140 x 45mm, acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 x 1.1m in 14 years. Late April.

(a) **'Gioia Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 1-3/truss, 67 x 42mm, with 5 wavy to very wavy-edged lobes, buds deep purplish pink (68A), opening inside strong purplish pink (68B), with a little sprinkled deep purplish pink (68A) and margins of 68A; outside strong purplish pink (68B) with margins of deep purplish pink (68A); three lobes with strong red (53B) markings; stamens 5-8, exerted from the corolla, filaments deep purplish pink (68A), anthers moderate orange-yellow (164B) shaded pale orange-yellow (16D); pistil equalling or shorter than the stamens, style strong purplish pink (67D), stigma light yellowish pink (26D). Lvs oblanceolate, 28 x 9mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (147A) above and moderate yellow-green (146D) below when mature; petioles 5-7mm long. Shrub 2.0 x 3.0m in 50 years (regularly pruned). May. (Etymology: "gioia" means "joy" in Italian).

(a) **'Giulia Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-10/hemispherical truss, 35 x 25mm, with 5 very wavy-edged lobes, deep pink (52C) in bud and the same colour both inside and out on opening; three lobes with deep yellowish pink (46D) markings; stamens 5, exerted from the corolla, filaments pale purplish pink (55D), anthers moderate red (183C); pistil just exceeding the stamens, style white, stigma brilliant yellow-green (149A). Lvs ovate, 24 x 16mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (137B) above and

moderate yellow-green (147B) below when mature; petioles 4mm long. Shrub 1.8 x 2.0m in 60 years (regularly pruned). April. (Etymology: named for the daughter of the nominant).

(v) **'Glenn Sahara'**

cv. Lepidote rhododendron: (*zoelleri* x *leucogigas*) (s) X 'Doctor Hermann Sleumer'. H: R.A. Mitchell (1995), G: G. Sahara (2003), N: R.A. Mitchell (2004), REG: R.A. Mitchell (2005). Fls 7/domed truss, tubular funnel-shaped, 85 x 25mm, with 5 wavy-edged lobes, margins deep yellowish pink (43C) inside and out, inside blending to brilliant yellow (11A) at base, outside blending into light yellow (11B) over its entire length; unmarked. Lvs elliptic, 100 x 65mm, broadly acute at apex, rounded at base, semi-glossy and dark yellowish green (139A) above, no scales visible to the naked eye. Shrub 1.2 x 0.9m in 5 years. December to January.

(a) **'Gloria Louis'**

cv. Evergreen azalea: 'Carol Kittel' (s) X 'Girard's Fuchsia'. H: J.F. Klimavicz (1998), G: J.F. Klimavicz (2000), N: J.F. Klimavicz (2005), I: J.F. Klimavicz (2005), REG: J.F. Klimavicz (2005). Fls 3/ball truss, saucer-shaped, 40 x 70mm, with 5 wavy-edged lobes, deep purplish pink (73A) in bud, opening inside moderate purplish pink (66D) with 5 yellowish white bands originating c.20mm from the base where each band is c.6mm wide and decreasing in width to c.1mm as it approaches the lobe tip; outside deep purplish pink (73A); two V-shaped, spotted, dorsal blotches of moderate to strong purplish red (59C-D). Calyx 7mm long, moderate yellow-green (138B). Lvs obovate, 50 x 20mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above. Dense shrub, 0.45 x 0.45m in 5 years, flowering from an early age. Mid April to May.

(r) **'Glowing'**

cv. Elepidote rhododendron: parentage unknown. H: H. O'Rourke (1994), G: H. O'Rourke (2001), N: H. O'Rourke (2005), REG: H. O'Rourke (2005). Fls in a flat-topped truss, 70 x 100mm, with 7 wavy-edged lobes, buds pink and yellow, opening yellow inside and out with pink lobes. Calyx 3mm long, red and green. Lvs lanceolate, 120 x 45mm, glossy green, obtuse at base, acute at apex, hairless. Shrub 1.25 x 1.0m in 10 years. November (Australia).

(r) **'Golden Mystery'**

cv. Elepidote rhododendron: parentage unknown. S: M. & H. Stockman, N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 14-16/domed truss, broadly funnel-shaped, 45 x 75mm, with 7 very wavy-edged lobes, strong red (53B) in bud, opening inside light yellow (11B), with an irregular, very dark blotch (almost black) in the base which is edged with strong red (46A) linear markings; corolla also with strong red (46A) dorsal spotting on a reddish ground; outside strong red (53B). Calyx either with 2 lobes c.12mm long and the rest c.1.5mm long, or all lobes c.1.5mm

long, strong red (53B). Lvs oblong, 95 × 40mm, broadly acute at apex, cordate at base, margins flat and downcurved, moderate olive green (137B) above, greyish yellow (148D) below, hairless. Shrub 1.0 × 1.5m. Late April. (Purchased by the Stockmans from a plant sale in 1997).

(r) **'Golden Promise'**

cv. Elepidote rhododendron: 'Golden Wit' (s) × *irroratum*. H: H. O'Rourke (1991), G: H. O'Rourke (1999), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 14/ball truss, funnel-shaped, 35 × 65mm, with 5 wavy-edged lobes, brilliant yellow-green (154C) in bud and throughout on opening; with a ray of red spots and dashes on three dorsal lobes. Calyx dark green. Lvs oblong, 80 × 35mm, margins flat, matt green above, cuneate at base, acute at apex, hairless. Shrub 0.75 × 0.6m in 12 years. October (Australia).

(r) **'Golden Sovereign'**

cv. Elepidote rhododendron: 'Prelude' (s) × 'Ostbo's Low Yellow'. H: H. O'Rourke (1988), G: H. O'Rourke (1997), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 14/rounded truss, funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, orange-yellow in bud, opening inside yellow, with the lobes edged in cream, outside pale yellow. Calyx 2mm long, green. Lvs elliptic, 115 × 55mm, margins wavy, glossy green above, rounded at base, acute at apex, hairless. Shrub 1.2 × 1.5m in 15 years. October (Australia).

(r) **'Goldkollier'**

cv. Elepidote rhododendron: 'Kupferberg' (s) × (('Goldrausch' × late-flowering *wardii* hybrid) × 'Goldrausch'). H: H. Hachmann (1994), G: H. Hachmann (1999), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 12/dense, rounded truss, broadly funnel-shaped, 40-45 × 65-75mm, with 5 strongly wavy-edged lobes, light orange in bud, opening inside light greenish yellow (8B), edges at first tinged a delicate orange-red, throat brilliant yellow (8A), outside light greenish yellow (8B) with delicate orange-red margins at first; dorsal lobe marked dark red/deep purplish red (59A/B); stamens 10, filaments light yellow-green (154D), anthers moderate reddish brown/greyish reddish orange (174A/B); style brilliant yellow-green (150A/B), stigma strong yellow-green (144A). Calyx 2-4mm long, greenish yellow. Lvs ovate, somewhat twisted, 100 × 40-55mm, margins slightly wavy, slightly glossy and light green above, with yellowish green petioles, midveins and edges, rounded to cordate at base, apiculate at apex, hairless. Shrub c.0.8 × c.1.1m in 10 years, with a broadly compact habit. Mid to late May. (Nomenclatural standard: colour print provided by registrant (WSY0070155). Published in Hachmann cat. 2004-2008 (2004), p.152).

(r) **'Goldsam'**

cv. Elepidote rhododendron: parentage unknown, a chance seedling, one parent of which is clearly

yakushimanum. S: G. Nouws, I: G. Coners (2006), N: W. Schmalscheidt (2005), REG: W. Schmalscheidt (2005). Fls 11-19/full, spherical truss, open funnel-shaped, 32-46 × 40-60mm, with 5 wavy-edged lobes, strong pink lilac in bud, opening inside light pink lilac, shading to nearly white in centre; outside purplish pink; green markings on the dorsal lobe. Calyx 0.5-1.0mm long, light green. Lvs elliptic to narrowly elliptic, 55-107 × 19-39mm, margins flat or decurved, matt green above and with striking yellow variegation (not only at the margin), obtuse at base, acute at apex. Shrub 0.4 × 0.4m in 11 years. Early to mid May. (Etymology: epithet refers to the gold margin of the leaves).

(r) **'Goldsprenkel'**

Syn. of 'Hachmann's Goldsprenkel'.

(r) **'Goldzwerg'**

cv. Elepidote rhododendron: ('Brinny' × ('Goldrausch' × late-flowering *wardii* hybrid)) (s) × 'Karibia'. H: Holger Hachmann (1993), G: Holger Hachmann (1998), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 12-14/lax ball truss, broadly funnel-shaped, 45-50 × 70-72mm, with 7 wavy-edged lobes, opening light greenish yellow (4C) inside and out; on the dorsal lobes, in the centre and near the throat the colour intensifies to light greenish yellow (7D); dorsal lobe with dark red (187B) markings; stamens 10, filaments light yellow-green (150D), anthers brownish orange (164A); style brilliant/light greenish yellow (1B/C), stigma strong yellow-green (144C). Calyx 2-3mm long, greenish yellow, narrowly rosy-edged. Lvs elliptic, slightly wavy, 70-85 × 35-40mm, margins flat, matt light green above, rounded at base, more-or-less apiculate at apex, hairless. Shrub c.0.55 × c.0.9m in 10 years, with a broadly compact habit. Early to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070157). Published in Hachmann cat. 2004-2008 (2004), p.152).

(a) **'Grace Katherine'**

cv. Evergreen azalea: 'Saybrook Glory' (s) × 'Dorothy Clark'. H: Robert Stewart (1996), G: Robert Stewart (1999), N: Robert Stewart (2004), I: Robert Stewart (2004), REG: Robert Stewart (2005). Fls solitary, broadly funnel-shaped, 50 × 80mm, with 5 wavy-edged lobes, somewhat brighter than vivid red (52A) in bud, opening inside light purplish pink (62C) with faint dorsal spotting; outside similar to bud colour. Calyx c.5mm long, close to brilliant yellow-green (142B). Lvs elliptic, 45 × 20mm, broadly acute at apex, cuneate at base, margins downcurved, semi-glossy and mid green above, hairless. Shrub 0.6 × 0.75m in 8 years. Midseason.

(a) **'Gracie Girl'**

cv. Evergreen azalea: ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'. H: W.L. Clagett (1996), G: W.L. Clagett (2000), N: W.L. Clagett (2005), I: White's Nursery (2006), REG: W.L. Clagett (2005). Fls hose-in-hose, 3/flat truss, open funnel-shaped, 30 × 40mm, with

5+5 wavy-edged lobes, strong red (53D) in bud, opening inside narrowly edged strong red (53D) shading to white in the centre and yellow-green at the base of the tube; outside strong red (53D) shading to white at the base; sparse salmon to ochre spotting on the dorsal lobe. Lvs elliptic, 45 × 25mm, broadly acute at apex, cuneate at base, margins flat, glossy above, hairless. Shrub 0.9 × 0.45m in 9 years. Midseason.

(r) **'Guyens's Djingis Khan'**

cv. Elepidote rhododendron: ('Souvenir of Anthony Waterer' × 'Direktor E. Hjelm') (s) × 'Goldzauber'. H: H.J.A. Guyens (1994), G: H.J.A. Guyens (1998), N: H.J.A. Guyens (1998), I: H.J.A. Guyens, REG: H.J.A. Guyens (2005). Fls 10-14/truss, funnel-shaped, 65 × 70mm, with 5 wavy-edged lobes, inside a mixture of orange, salmon and yellow, paling with age, margins salmon pink aging to yellow; outside a mixture of orange, salmon and yellow; three dorsal lobes with ochre to green markings and spots. Lvs ovate, 75-120 × 35-55mm, margins not wavy but slightly upcurved, matt green above, broadly acute at apex, rounded at base, hairless, veins conspicuous both above and below. Compact shrub, 0.65 × 0.75m in 7 years. May.

(r) **'Guyens's Paradiesvogel'**

cv. Elepidote rhododendron: 'Guyens's Fasching' (s) × 'Marrakesch'. H: H.J.A. Guyens (1994), G: H.J.A. Guyens (1998), N: H.J.A. Guyens (2002), I: H.J.A. Guyens, REG: H.J.A. Guyens (2005). Fls 11-15/dense truss, funnel-shaped, 75 × 90mm, with 5 wavy-edged lobes, inside pink, marbled with white, margins pinkish red, outside pinkish red; dorsal lobes with ochre yellow markings and brown spots, chiefly on the main dorsal lobe; stamens and style white. Lvs oblong, 115-130 × 40-60mm, margins not wavy but decurved, broadly acute at base and apex, hairless. Compact shrub, 1.1 × 1.3m in 10 years. Early to mid May. (Fls larger and fleshier than in 'Professor Horst Robenek'. Nomenclatural standard: colour print provided by the registrant (WSY0052452)).

(r) **'Hachmann's Goldsprenkel'**

cv. Elepidote rhododendron: (*dichroanthum* subsp. *scythocalyx* hybrid × 'Allegretto') (s) × (('Golddrausch' × late-flowering *wardii* hybrid) × 'Golddrausch'). R: H. Hachmann (1993), H: H. Hachmann (1993), G: H. Hachmann (1999), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 16-21/dense ball truss, broadly open funnel-shaped, 40-45 × 65-75mm, with 5 wavy to strongly wavy-edged lobes, opening inside and out light greenish yellow (7D), tinged brilliant yellow/brilliant greenish yellow (7B/C); three dorsal lobes with numerous chocolate markings; stamens 10 (with 1-2 tending to be slightly petaloid), filaments pale greenish yellow (10D), anthers greyish brown/moderate reddish brown (166A/B); style brilliant yellow-green (149A/B), stigma strong yellow-green (144B); ovary moderate yellow-green/light yellow-green (138C/D), silvery hairy. Calyx 3-5mm long, green, edged red. Lvs ovate,

c.55-100 × c.35-50mm, margins flat or slightly wavy-edged, matt mid green above, rounded at base, apiculate at apex, hairless. Shrub c.0.7 × c.0.9m in 10 years. Mid to late May, with long-lasting fls. Syn: 'Goldsprenkel'. (Nomenclatural standard: colour print provided by the registrant (WSY0070156). Published in Hachmann cat. 2004-2008 (2004), p.152, as 'Goldsprenkel' (an unacceptable name, already in use, Art. 27.1)).

(a) **'Hachmann's Juliette'**

cv. Evergreen azalea: 'Squirrel' (s) × 'Chippewa'. H: Holger Hachmann (1990), G: Holger Hachmann (1994), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls single, 2-3/truss, broadly funnel-shaped with a moderate to long tube, 30-35 × 38-44mm, with 5 slightly wavy-edged lobes, buds moderate red (47A), opening inside and out vivid red (46B), shaded deep yellowish pink (43C) and tinged with orange-red; dorsal lobe with sparse, vivid red (46B) markings; stamens 5, filaments strong purplish pink (55B), anthers deep red (53A); style strong/moderate purplish pink (62A/B), stigma dark red (187B); ovary greenish, silvery hairy. Lvs ovate, slightly convex, c.20-30 × 12-17mm, margins flat, dark shiny green above, cuneate at base, obtuse and slightly apiculate at apex, with silvery hairs; lasting well in winter. Shrub c.0.4 × c.0.8m in 10 years; habit flat-rounded. Early to late June (i.e. late). Syn: 'Juliette'. (Nomenclatural standard: colour print provided by the registrant (WSY0070163). Published in Hachmann cat. 2004-2008 (2004), p.176, as 'Juliette' (an unacceptable name, already in use, Art. 27.1)).

(r) **'Hachmann's Peggy'**

cv. Elepidote rhododendron: 'Paprika Spiced' (s) × 'Dolcemente'. H: H. Hachmann (1993), G: H. Hachmann (1998), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 7-9/rounded, upright, rather loose truss, broadly funnel-shaped, 50-55 × 65-85mm, with (5-)7 wavy-edged lobes, pale greenish yellow (13D) throughout, the dorsal lobes with conspicuous markings of dark red (187B/C) on a brilliant yellow (9C) ground; stamens 12-13, filaments yellowish white (158D), anthers moderate brown (165A); style brilliant yellow-green (150C) at base and tinged rosy near the apex, stigma moderate reddish brown/brownish orange (166B/C); ovary strong yellow-green (144B). Calyx hose-in-hose, 5-7-lobed, 10-18mm long, light yellow (12C). Lvs elliptic to ovate, 80-110 × 30-45mm, margins flat to slightly wavy, matt light green above, rounded at base, apiculate at apex, hairless. Shrub c.0.9 × c.1.1m in 10 years, with a dense, compact habit. Early to late May. Syn: 'Peggy'. (Nomenclatural standard: colour print provided by the registrant (WSY0070146). Published in Hachmann cat. 2004-2008 (2004), p.67, as 'Peggy' (an unacceptable name, already in use, Art.27.1)).

(r) **'Hachmann's Picobello'**

cv. Elepidote rhododendron: 'Hachmann's Kabarett' (s) × 'Schneespiegel'. H: Holger Hachmann (1991),

G: Holger Hachmann (1996), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 10-11/compact truss, broadly open funnel-shaped, 50-55 × 75-85mm, with 5(-6), wavy to very wavy-edged lobes, opening inside and out pure white (155D), dorsal lobe with a large, conspicuous, dark red (187A) blotch from the throat on to the lower half of the dorsal lobe and breaking up into markings of dark red to deep purplish red (187A-D); stamens 10, filaments white, anthers moderate orange-yellow/pale orange-yellow (165C/D); style brilliant yellow-green/pale yellow-green (149C/D), stigma greyish reddish orange (174D). Calyx 6-10mm long, light yellow-green (154D). Lvs ovate, 100-120 × 50mm, margins wavy, dark glossy green above, rounded at base, apiculate at apex, hairy. Shrub c.1.1 × c.1.4m in 10 years, with a broad, compact, rounded habit. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070135). Published in Hachmann cat. 2004-2008 (2004), p.50).

(r) **'Hachmann's Penguin'**

cv. Elepidote rhododendron: 'Hachmann's Kabarett' (s) × ('Scintillation' × 'Germania'). H: H. Hachmann (1994), G: H. Hachmann (2000), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 11-13/dense, ball truss, broadly open funnel-shaped, 50-55 × 75-90mm, with 5 wavy to very wavy-edged lobes, inside white, edged very pale purple (76C), outside similarly coloured but midveins light purple (84B), the colour spreading out onto the lobes; dorsal lobe with a large, conspicuous blotch of dark red (187A); stamens 10, filaments white, anthers moderate yellow (161A/B); style light yellow-green (150D), stigma dark red (187C); ovary greenish, silvery hairy. Calyx 5-8mm long, greenish/reddish. Lvs elliptic, c.120 × 45mm, margins flat or slightly wavy, matt green above, rounded at base, acute at apex, hairless. Shrub c.1.2 × c.1.5m in 10 years, with a broadly upright habit. Late May to early June. Syn: 'Penguin'. (Nomenclatural standard: colour print provided by the registrant (WSY0070147). Published in Hachmann cat. 2004-2008 (2004), p.68. as 'Penguin' (an unacceptable name, too similar to one already in use, Art.19.25)).

(a) **'Hachmann's Thekla'**

cv. Evergreen azalea: 'Elsie Lee' (s) × 'Purpurkissen'. H: H. Hachmann (1988), G: H. Hachmann (1992), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls double, 1-2(-3)/truss, with two whorls one within another, broadly funnel-shaped, 25-30 × 50-55mm, with 11-17 wavy-edged lobes, deep purplish pink (72D) inside and out; dorsal lobes with strong purplish red (72A) markings; stamens all transformed to make the inner corolla whorl; style unusually long, deep purplish pink (54B), stigma dark red (187B). Calyx 2-3mm long, strong yellow-green (145A). Lvs ovate, slightly concave, 25-35 × 15-23mm, margins flat, mid green and slightly glossy above, cuneate at base, apiculate at apex, with a few silvery hairs. Shrub c.0.5 × c.0.7m in 10 years,

with a compact, rounded habit. Late May to early June. Syn: 'Thekla'. (Nomenclatural standard: colour print provided by the registrant (WSY0070167). Published in Hachmann cat. 2004-2008 (2004), p.176, as 'Thekla' (an unacceptable name, already in use, Art.27.1)).

(r) **'Haithabu'**

cv. Elepidote rhododendron: 'Walküre' (s) × 'Germania'. H: Holger Hachmann (1991), G: Holger Hachmann (1997), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 16-18/ rounded truss, wide-open funnel-shaped, 50-55 × 90-110mm, with 5 narrowly wavy-edged lobes, strong purplish red (64B/C) in bud, opening inside deep/strong purplish pink (67C/D) tinged strong/vivid purplish red (67A/B), outside similarly coloured but slightly darker (vivid purplish red, 66B) on veins; dorsal lobe with deep purplish red (187D) spots; stamens 10, filaments deep/strong purplish pink (67C/D), anthers greyish reddish orange (174B); styles moderate purplish pink (68C), stigma dark red (187B); ovary light greyish olive (195A), whitish hairy. Calyx 2mm long, greenish yellow. Lvs ovate, 140-180 × 65-85mm, margins flat, dark green and matt to slightly glossy above, rounded to slightly oblique at base, acuminate at apex, with inconspicuous brownish hairs below. Shrub c.1.4 × c.1.6m in 10 years; habit broadly rounded. Mid May to early June. Fls long-lasting, not fading in the sun. (Nomenclatural standard: colour print provided by the registrant (WSY0070141). Published in Hachmann cat. 2004-2008 (2004), p.50).

(r) **'Harold Jordan'**

cv. Elepidote rhododendron: selection from *rex* subsp. *fictolactenum*. S: H. Jordan (1999), G: Canterbury Rhododendron Society (1999), N: Canterbury Rhododendron Society (2004), REG: Canterbury Rhododendron Society (2005). Fls 15/rounded truss, campanulate, 40 × 25mm, with 8 very wavy-edged lobes, pinkish, cream and white, with a minimal crimson blotch. Calyx very short. Lvs oblanceolate, 170 × 55mm, margins decurved, very dark matt green above, rounded at base, obtuse at apex, with buff to rusty brown hairs. Shrub 2.0 × 1.75m in 7 years. October (New Zealand). (Selected for its superb foliage. Etymology: named for the patron of the Canterbury Rhododendron Society. Published in Canterbury Rhododendron Society Newsletter, 2004).

(r) **'Herbstzauber'**

cv. Elepidote rhododendron: 'Guyens's Fasching' (s) × 'Marrakesch'. H: H.J.A. Guyens (1994), G: H.J.A. Guyens (1998), N: H.J.A. Guyens (2002), REG: H.J.A. Guyens (2005). Fls 12-16/loose, open-topped truss, out to slightly down-facing, funnel-campanulate, 70 × 80mm, with 5-6 slightly wavy-edged lobes, buds salmon pink, opening inside creamy pink, with two large, orange-brown, dorsal markings consisting of flecks and spots (mostly on or below the dorsal lobe). Pedicels 40-65mm long,

red-brown. Lvs lanceolate, 90-115 × 30-35mm, margins not wavy, but slightly decurved, matt dark green above, acute at base and apex, hairless; petioles and new shoots red-brown. Compact shrub 1.2 × 1.4m in 10 years. Flowering in spring, in the first half of May and then for a long period in the autumn (early September to mid November). (Nomenclatural standard: colour print provided by the registrant (WSY0052450)).

(r) **'Highborn'**

cv. Lepidote rhododendron: *nutallii* (s) × 'Alf Bramley'. H: H. O'Rourke (1990), G: H. O'Rourke (2000), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 4-5/loose truss, tubular funnel-shaped, 110 × 120mm, with 5 wavy-edged lobes, buds pink and cream, opening inside white with a yellow centre, outside white. Calyx 25mm long, green. Lvs oblanceolate, 150 × 60mm, wavy-edged, cuneate at base, broadly acute at apex. Shrub 1.7 × 1.7m in 13 years. October (Australia).

(r) **'Holly Snaza'**

cv. Elepidote rhododendron: 'Award' (s) × 'Betty Anderson'. H: E.T. Anderson (1995), G: E. Metcalfe, N: B.E. Anderson (2004), REG: B.E. Anderson (2005). Fls 13/ball truss, broadly funnel-shaped, 70-75 × 100mm, with 6 very wavy-edged lobes, strong purplish red (67A) in bud, opening white inside and out, edged deep purplish pink (67C) and with a strong greenish yellow (151B) basal blotch. Calyx 2mm long, light yellow-green (145B). Lvs elliptic, 100-175 × 40-65mm, semi-glossy and moderate olive green (137A) above, hairless. Shrub 1.2 × 1.2m in 9 years. Mid May.

(a) **'Ines Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls hose-in-hose, 2-4/truss, 30 × 23mm, with 5+5 very wavy-edged lobes, buds red-purple (N57B), opening red-purple (N57C) throughout; three lobes with red-purple (N57A) markings; stamens 5, exerted from the corolla, anthers greyish reddish orange (176D), filaments moderate purplish pink (68C); pistil just longer than stamens, style moderate purplish pink (68C), stigma deep yellowish pink (44D). Lvs obovate, 13 × 6mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (147A) above and moderate yellow-green (146D) below when mature; petioles 2-4mm long. Shrub 1.5 × 3.5m in 60 years (regularly pruned). May. (Etymology: named for the nominant's aunt).

(r) **'Innocent Air'**

cv. Lepidote rhododendron: (*johnstoneanum* × Chrysoamanic Group) (s) × 'Pink Silk'. H: H. O'Rourke (1994), G: H. O'Rourke (2003), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 8-9/rounded truss, funnel-shaped, 50 × 55mm, with 5 wavy-edged lobes, pale yellow and pink in bud, opening inside cream, outside pale yellow with pink on the lobes; yellow blotch. Calyx greenish yellow.

Lvs elliptic, 60 × 35mm, margins upcurved, matt green above, obtuse at base, acute at apex, scaly. Shrub 0.6 × 0.5m in 10 years. September to October (Australia).

(a) **'Intra'**

cv. Evergreen azalea: parentage unknown. R: Fratelli Savioli (1970), N: Fratelli Savioli (1985), I: Fratelli Savioli (1985), REG: Prof. E. Accati (2005). Fls single, 2-4/truss, 20 × 20mm, with 5 flat-edged lobes, strong red (53C) in bud and on opening, both inside and out, unmarked; stamens 5, exerted from the corolla, anthers greyish reddish brown (200B), filaments strong red (53C); pistil not or just exceeding stamens, style strong red (53D), stigma brownish orange (165B). Lvs lanceolate-obovate, 13 × 6mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (146D) below when mature; petioles 3mm long. Shrub 1.1 × 1.1m in 10 years. April. (Etymology: named for a village near the city of Verbania; the latter is where the Savioli nursery is situated).

(a) **'Jake Frey'**

cv. Evergreen azalea: unknown (a semi-double, evergreen/semi-evergreen plant referred to as CB-1) (s) × 'Mitey White'. H: Robert Stewart (1994), G: Robert Stewart (1997), N: Robert Stewart (2004), I: Robert Stewart (2004), REG: Robert Stewart (2005). Fls semi-double, saucer-shaped, 3/truss, 45 × 80mm, with 5 + 0.5 very wavy-edged lobes, white in bud, opening white, sanded and streaked with vivid purplish red (66A); some flowers open near strong purplish pink (67D) with vivid purplish red (66A) markings; blotch small, yellow-green. Calyx c.6mm long, brilliant yellow-green (149C). Lvs elliptic, 50 × 20mm, broadly acute at apex, cuneate at base, margins downcurved, semi-glossy and mid green above, hairless. Shrub 0.9 × 0.9m in 8 years. Midseason.

(a) **'Jhonatan Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls hose-in-hose, 2-4/truss, 30 × 21mm, with 5+5 flat-edged lobes, red-purple (N74B) in bud and throughout on opening; unmarked; stamens 5, exerted from the corolla, anthers light yellowish brown (199C), filaments red-purple (N74B); pistil exceeding stamens, style red-purple (N61C), stigma strong red (50A). Lvs obovate, 10 × 6mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (147A) above and moderate yellow-green (147C) below when mature; petioles 2mm long. Shrub 0.8 × 1.8m in 60 years (regularly pruned). April. (Etymology: named for the nominant's son).

(r) **'Jonathan Bryce'**

cv. Elepidote rhododendron: 'Blue Peter' (s) × 'White Dimples' H: T. Ahern (1994), G: J. Doppel (2000), N: J. Doppel (2004), REG: J. Doppel (2005). Fls 18/domed truss, open funnel-shaped, 50 × 55mm,

with 6 very wavy-edged lobes, light greyed lavender in bud, opening white inside and out, with a large, V-shaped dorsal flare composed of dark purple spots. Lvs obovate, 140 × 50mm, obtuse at apex, rounded at base, margins slightly downcurved, semi-glossy and dark green above, hairless. Shrub 1.1 × 1.1m in 10 years. Mid May.

(r) **'J.S. Garrett 3rd'**

cv. Elepidote rhododendron: 'Lockington Pride' (s) × 'Cup Day'. H: D.J. Dosser (1990), G: D.J. Dosser (2000), N: D.J. Dosser (2004), REG: D.J. Dosser (2005). Fls 13/domed truss, funnel-shaped, 75 × 70mm, with 5 wavy-edged lobes, inside white, lightly streaked strong purplish red (62B) and with numerous vivid purplish red (61C) spots on all the lobes (heaviest on 3 dorsal lobes); outside white, streaked strong purplish red (62B); slightly scented. Lvs oblong, 175 × 50mm, attenuate at base, cuspidate at apex, hairless. Shrub 2.0 × 1.0m in 14 years. November (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0070181)).

(a) **'Juliette'**

Syn. of 'Hachmann's Juliette'

(r) **'Juniflair'**

cv. Elepidote rhododendron: 'James Burchett' (s) × 'Tarantella'. H: H. Hachmann (1989), G: H. Hachmann (1995), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 11-14/dense, ball truss, broadly funnel-shaped, 55-60 × 75-90mm, with 5(-6) wavy-edged lobes, buds strong purplish red (60B/C), becoming tinged with deep/strong purplish red (60A/B), opening inside and out strong purplish red (63B), shaded strong purplish red (61B), outside similarly coloured but with a tinge of vivid purplish red (57C) on the veins; dorsal lobe with dark red (187A/B) markings on a light yellow/white ground; stamens 10, filaments white, anthers pale yellow (162D); styles light yellow-green (150D), stigma dark red (187B); ovary greenish, white hairy. Calyx 1-2mm long, reddish/greenish. Lvs elliptic to ovate, 120-150 × 50-60mm, margins flat to slightly wavy, matt mid green above, rounded at base, apiculate at apex, hairless. Shrub c.1.1 × c.1.3m in 10 years; habit broadly upright. Early to mid June. (Nomenclatural standard: colour print provided by the registrant (WSY0070136)). Published in Hachmann cat. 2004-2008 (2004), p.53.

(r) **'Kahlua'**

cv. Elepidote rhododendron: 'Sunspray' (s) × 'Lem's Cameo'. H: J. Olson, G: J. Olson (2005), N: J. Olson (2005), REG: J. Olson (2005). Fls 12/domed truss, open funnel-shaped, 75 × 100mm, with 6 wavy-edged lobes, strong purplish pink (55B) in bud, opening inside light to pale purplish pink (55C-D) at margins, shading to light yellow-green (2C) in the throat; outside strong to light purplish pink (55B-C), the slightly darker colour at lobe centres; no markings;

moderately scented. Calyx 50mm long, similar to the corolla. Lvs oblong, 150 × c.70mm, obtuse at apex, slightly cordate at base, with wavy margins, semi-glossy and dark yellowish green (139A) above, hairless. Shrub 1.5 × 1.5m in 6 years, with long-lasting flowers of heavy substance. Early May.

(r) **'Kangaro'**

cv. Elepidote rhododendron: 'Hachmann's Kabarett' (s) × 'Peter Alan'. H: Holger Hachmann (1994), G: Holger Hachmann (1999), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 15-17/rounded, fairly compact truss, broadly funnel-shaped, 45-50 × 75-80mm, with 5 wavy-edged lobes, opening inside vivid purplish red (78A), shaded vivid purple (80A), outside similarly coloured but with midveins shaded strong purplish red/strong reddish purple (72A/B); with a conspicuous blotch on 3 dorsal lobes of dark red (187A), shaded nearly dark greyish purple (202A) in the centre; stamens 10, filaments light to pale purplish pink (65B/C), anthers pale orange-yellow (159B); style near deep/strong purplish pink (73A/B) near apex, whitish towards base, stigma dark red (187A/B); ovary greenish, silvery hairy. Calyx 2-4mm long, greenish/reddish. Lvs narrowly elliptic to ovate, c.110 × 35-45mm, margins flat, slightly wavy, dark green and slightly glossy above, rounded at base, obtuse at apex, hairless. Shrub c.1.4 × c.1.7m in 10 years; habit compact, rounded. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070137)). Published in Hachmann cat. 2004-2008 (2004), p.54.

(r) **'Karen Swenson'**

cv. Elepidote rhododendron: (*calophytum* × Sarita Loder Group) (s) × *macabeanum*. H: Mrs E.M. Watson (1969), G: Mrs E.M. Watson (1978), N: Mrs E.M. Watson, REG: Mrs E.M. Watson (2005). Fls 14/ball truss, open campanulate, 80 × 80mm, with 5 wavy-edged lobes, light purplish pink (62C) in bud, opening light purplish pink (62C) and fading to white, with a dark purple (79A) basal flare (from 5-20mm long). Lvs elliptic, 300 × 105mm, acute at apex, cuneate at base, margins very slightly upcurved, glossy and dark yellowish green (139A) above, with a felted indumentum below (moderate orange-yellow (164C) when young, aging to pale yellow (164D)). Shrub of open habit, 4.2 × 3.6m in 35 years. Early March.

(r) **'Karminkissen'**

cv. Elepidote rhododendron: *yakushimanum* 'Koichiro Wada' (s) × 'Ruby Hart'. H: Holger Hachmann (1992), G: Holger Hachmann (1998), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 8/flattened ball truss, broadly funnel-shaped, 45-50 × 50-65mm, with 5 slightly wavy-edged lobes, opening inside and out strong red (53D), shaded strong red (53C); unmarked; stamens 10, filaments strong red (53D), anthers moderate reddish brown (174A); style strong red (53C), stigmas deep red (53A), ovary with dense silvery hairs. Calyx 35-40mm long, nearly as long as

the corolla, strong red (53D), shaded deep red (53A). Lvs narrowly elliptic, c.85 × 20-25mm, margins slightly decurved, dark glossy green above, rounded at base, mucronate at apex, with a dense, pale brown indumentum below. Shrub c.0.4 × c.0.7m in 10 years, with a semi-prostrate habit. Mid to late May. (Nomenclatural standard: colour print provided by the registrant (WSY0070150). Published in Hachmann cat. 2004-2008 (2004), p.97).

(r) **'Keen and Peachy'**

cv. Elepidote rhododendron: 'White Elation' (s) × 'Barbara Hardgrove'. H: D.B. Johnson (1998), G: D.B. Johnson (2005), N: D.B. Johnson (2005), REG: D.B. Johnson (2005). Fls 14/domed truss, broadly funnel-shaped, 65 × 100mm, with 7 flat-edged lobes, moderate yellowish pink (38B) in bud, opening inside pale yellow (11C) at base, changing to a 20mm wide, pale pink (49D) margin; outside pale yellow (8D) at base, shading paler to almost white, tinged pale pink (49D) at edge. Lvs elliptic, 125 × 55mm, obtuse at apex, oblique at base, margins wavy, glossy and moderate yellow-green (137C) above, hairless. Shrub 1.5 × 1.5m in 10 years. Mid May.

(r) **'King's Lady'**

cv. Elepidote rhododendron: Jalisco Group (s) × 'Mrs P.D. Williams'. H: H. O'Rourke (1991), G: H. O'Rourke (2000), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 10/rounded truss, broadly funnel-shaped, 50 × 100mm, with 5 lobes, cream in bud, opening cream with mauve/pink lobes both inside and out; large flare of reddish brown streaks on dorsal lobe. Calyx 20mm long, cream. Lvs oblong, 130 × 45mm, margins wavy, hairless. Shrub 1.25 × 1.2m in 12 years. October to November (Australia).

(r) **'Kinnor'**

cv. Elepidote rhododendron: 'Fantastica' (s) × *dicbroanthum* subsp. *apodectum*. H: Holger Hachmann (1993), G: Holger Hachmann (1998), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 7-10/ rather loose truss, funnel-campanulate, 35-40 × 55mm, with 5 wavy-edged lobes, inside light orange-yellow (23C), edged strong red (45D), outside strong red (45D), shaded vivid red (45B/C) on and beside veins; unmarked; stamens 10, filaments white (155B) - 3-5 of them with 1-2 flattened, greenish/white, hairy lobes (2-3mm long) at their base, anthers moderate/light reddish brown (177A/B); style light yellow-green (154D), stigma deep greenish yellow (153A); 5 conspicuous deep red nectaries in the throat. Calyx 1-2mm long, greenish red. Lvs ovate, 40-80 × 20-36mm, margins slightly decurved, matt dark green above, rounded at base, obtuse at apex, hairy (hairs silvery when young, aging to silvery/brownish). Shrub c.0.65 × c.0.9m in 10 years, with a rounded, very compact habit. Early to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070158). Published in Hachmann cat. 2004-2008 (2004), p.122).

(v) **'Kiri'**

cv. Lepidote rhododendron: 'Solar Flare' (s) × unknown. H: D. Brown (1999), G: D. Brown (2005), N: D. & P.J. Brown (2005), REG: D. & P.J. Brown (2005). Fls 14-16/full, rounded truss, tubular funnel-shaped, 75 × 70mm, with 7 flat-edged lobes, orange-red in bud, opening inside to 'Granny Smith' apple green in the tube shading to spanish orange (10/1) and with azalea pink (618/2) lobes, outside similarly coloured; unmarked; scented. Calyx 3mm long, reddish brown. Lvs elliptic, 110 × 55mm, margins flat, matt green above, cuneate at base, acute at apex, with a scaly indumentum (scales reddish brown, aging to brown); young foliage reddish. Upright shrub, up to 2.0 × 1.0m. Repeat-flowering.

(r) **'Kokette'**

cv. Elepidote rhododendron: 'Ovation' (s) × 'Schlaraffia'. H: H. Hachmann (1988), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 10-12/somewhat loose truss, campanulate, 40-50 × 50-60mm, with 5 strongly wavy-edged lobes, inside pure white except for strong red (53D) edge, outside nearly all strong red (53D) but paler near the base; unmarked; stamens 10, filaments light/pale purplish pink (55C/D), anthers moderate brown (165A); style light/pale purplish pink (55C/D), stigma moderate red (183C). Calyx 1-2mm long, light yellow-green (144D). Lvs elliptic to ovate, 80-110 × 45-50mm, margins slightly wavy, dark and slightly glossy green above, rounded at base, apiculate at apex, with a thin, fawn-coloured indumentum below. Shrub c.0.7 × c.1.1m in 10 years, with a broadly rounded, dense, compact habit. Early to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070151). Published in Hachmann cat. 2004-2008 (2004), p.98).

(r) **'Kyle Snaza'**

cv. Elepidote rhododendron: 'Van Nes Sensation' (s) × 'Wilburn'. H: E.T. Anderson (1975), G: B. Mendlin, N: B.E. Anderson (2004), REG: B.E. Anderson (2005). Fls 13-15/ball truss, broadly funnel-shaped, 65 × 100mm, with 6 wavy-edged lobes, light purple (80C) in bud, opening inside light purple (80C) at margins, changing abruptly to very light purple (76B) with a vivid purple (81B) stripe on each midrib and a large, prominent, speckled, dark red (59A) dorsal blotch; outside light purple (80C). Calyx 2-3mm long, vivid purplish red (57D). Lvs elliptic, 140-200 × 50mm, broadly acute at apex, cuneate to rounded at base, margins upcurved, semi-glossy and moderate olive green (137A) above, hairless. Shrub 1.2 × 0.9m in 6-8 years (2.4-3.0m tall in 29 years). Early May.

(r) **'Lachlan Arblaster'**

cv. Elepidote rhododendron: (Erebus Group × *yakushimanum*) (s) × 'Etta Burrows'. H: G. Arblaster (c. 1984-1985), G: G. Arblaster (1990), N: Mrs J. Arblaster (2005), REG: Mrs J. Arblaster (2005). Fls 9/hemispherical truss, broadly funnel-shaped, 65 × 70mm, with 5 more-or-less flat-edged lobes, inside

strong purple (77B) in a broad marginal band, sharply contrasting with the pure white centre, outside strong purple (77B) fading to light purple (75B) at the base; small dorsal fan of vivid red (45A) spotting in the dorsal throat and at base of dorsal lobe (fading to moderate orange (167A), overlaid vivid red (45A)). Stigma purplish; style yellowish white. Calyx up to 30mm long, strong yellow-green (144B). Lvs oblanceolate, up to 180 × 50mm, margins flat to slightly wavy edged, matt dark green (135A) above, leathery, cuneate at base, broadly acute at apex, minutely glandular on petiole and base of midvein. Shrub 2.7 × 2.15m in 15 years. Mid May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0103661)). Etymology: named for the second son of the registrant).

(a) **'La Coletta'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 1-2/truss, 49 × 38mm, with 5 wavy to very wavy-edged lobes, vivid reddish orange (42A) in bud and throughout on opening; three lobes with strong red (46A) markings shaded with vivid red (46B); stamens 4-5, not exerted from corolla, anthers black, filaments strong red (50A) shaded strong pink (50C); pistil distinctly exceeding stamens, style vivid red (45A) shaded moderate red (47A), stigma strong red (46A). Lvs lanceolate, 57 × 20mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (147B) below when mature; petioles 4mm long. Shrub 1.4 × 1.5m in 50 years. April to May. (Etymology: named for a place near Cannero Riviera (the latter being where the Carmine nursery is located)).

(a) **'Lady Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single (to semi-double), 2-3/truss, 75 × 45mm, with 5(-6) flat-edged or slightly wavy-edged lobes, deep purplish pink (67C) in bud and on opening, both inside and out, with deep purplish red (61A) markings on three lobes; stamens 8-11, exerted from the corolla, anthers moderate yellow (161B), filaments light purple (75A); pistil exceeding the corolla but shorter than the stamens, style moderate purplish pink (68C), stigma strong yellowish pink (32D). Lvs oblanceolate, 34 × 13mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (147A) above and moderate yellow-green (146C) below when mature; petioles 7-8mm long. Shrub 1.5 × 3.0m in 60 years (regularly pruned). May. (Etymology: the epithet marks both the femininity of this azalea and the family who produced it).

(a) **'La Grotta'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls

hose-in-hose, 1-2/truss, 31 × 30mm, with 5+5 flat to rarely wavy-edged lobes, buds strong purplish pink (55B), opening pale purplish pink (55D) shaded strong purplish pink (55B) both inside and out, unmarked; stamens 5, not exceeding the corolla, anthers moderate orange (167A), filaments strong purplish pink (55B); pistil equalling the corolla, style strong purplish red (54A), stigma moderate orange (167A). Lvs elliptic, 24 × 12mm, margins flat, glossy above, acute at base and apex, hairy, strong yellow-green (144A) above and moderate yellow-green (146C) below when mature; petioles 4mm long. Shrub 1.7 × 2.0m in 50 years. April. (Etymology: named for a place near Cannero Riviera (the latter being where the Carmine nursery is situated)).

(a) **'Laura Zegna'**

cv. Evergreen azalea: parentage unknown. N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single, 61 × 38mm, with 5 flat-edged lobes, buds purple (N78B), opening inside purple (N78C), outside purple (N78C); three lobes with moderate purplish red (59C) markings shaded deep purplish red (59B); stamens 9-10, exerted from the corolla, anthers violet-blue (N92A), filaments light purple (78C); pistil just exceeding the stamens, style deep pink (52B), stigma strong yellowish pink (35C). Lvs lanceolate, 39 × 18mm, margins flat, acute at base and apex, glossy above, hairy, moderate olive green (137A) above and moderate yellow-green (137C) below when mature; petioles 4mm long. Shrub 1.7 × 1.7m in 100 years (regularly pruned). May. (Etymology: named for Mrs Laura Zegna, director of Oasi Zegna, a location in Piemonte with an important collection of rhododendrons. Probably raised (c.1890) by Fratelli Rovelli and Fratelli Contini, gardeners at the Villa del Pascià in the nineteenth century - this being the villa where the then King of Egypt stayed during his Italian holidays on Lake Maggiore).

(a) **'Laveno'**

cv. Evergreen azalea: parentage unknown. R: Savioli Fratelli (1970), N: Savioli Fratelli (1985), I: Savioli Fratelli (1985), REG: Prof. E. Accati (2005). Fls single, 1-10/hemispherical truss, 51 × 33mm, with 5 wavy-edged lobes, buds strong reddish purple (78B), opening inside and out white, shaded light purple (75A) and strong reddish purple (78B); three lobes with strong red (46A) markings; stamens 5-7, exceeding corolla, anthers moderate olive brown (199A), filaments white; pistil just exceeding stamens, style light yellow-green (150D), stigma brilliant yellow-green (149A). Lvs lanceolate-obovate, 42 × 19mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (146A) above and moderate yellow-green (146D) below when mature; petioles 5mm long. Shrub 1.0 × 1.0m in 10 years. April. (Etymology: named for a village near the city of Verbania, Italy. The latter is where the Savioli nursery is situated).

(a) **'Lavinia Borromeo Elkan'**

cv. Evergreen azalea: parentage unknown. N: Prof.

E. Accati (2005), REG: Prof. E. Accati (2005). Fls single, 1-2/truss, 70 × 42mm, with 5 wavy to very wavy-edged lobes, buds strong red (50A), opening inside and out deep pink (50B); three lobes with deep red (53A) markings; stamens 8-10, exserted from the corolla, anthers purple (N77A), filaments deep pink (52B); pistil distinctly exceeding the stamens, style deep pink (52B), stigma strong yellowish pink (35C). Lvs lanceolate, 36 × 13mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (137A) above and moderate yellow-green (137C) below when mature; petioles 4-5mm long. Shrub 1.7 × 4.0m in c.100years (but regularly pruned). May. (Etymology: named for a descendent of the Borromeo family, owners of the islands on Lake Maggiore, Italy. Probably raised (c.1890) by Fratelli Rovelli and Fratelli Contini, gardeners at the Villa del Pascià in the nineteenth century. This is the villa where the then King of Egypt stayed during his Italian holidays on Lake Maggiore).

(a) **'Lea Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 36 × 26mm, with 5 flat-edged lobes, light reddish purple (74C) in bud and on opening, both inside and out, three lobes with deep purplish red (61A) markings; stamens 5, exserted from the corolla, anthers greyed orange (N167A), filaments very pale purple (75D); pistil exceeding the stamens, style moderate purplish pink (68C) shaded white (155D), stigma light orange-yellow (25D). Lvs oblanceolate, 18 × 8mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (137A) above and moderate yellow-green (146D) below when mature; petioles 3-4mm long. Shrub 1.0 × 3.0m in 60 years (regularly pruned). April. (Etymology: named for the nominant's aunt).

(r) **'Lemon Embers'**

cv. Elepidote rhododendron: 'Percy Wiseman' (s) × 'Recital'. H: J. Barlup (1996), G: J. Barlup (2000), N: J. Barlup (2005), REG: J. Barlup (2005). Fls 14/ball truss, open funnel-shaped, 40 × 65mm, with 5 wavy-edged lobes, strong red (53B) in bud, opening inside brilliant yellow (13C), blending at edges to moderate purplish pink (62B) and to strong purplish pink (63C) at centre of lobe tips; two moderate red (185B) dorsal flares consisting of striated rays terminating in small spots, extending upwards c.25mm from base; outside brilliant yellow (13C) with moderate purplish pink (62B) edges and strong purplish pink (63C) midribs. Lvs elliptic, 90 × 45mm, obtuse at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 9 years. Late May.

(a) **'Li'l Fella'**

cv. Evergreen azalea: ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'. H: W.L. Clagett (1996), G: W.L. Clagett (1999), N: W.L. Clagett (2005), I: White's Nursery (2006), REG: W.L. Clagett (2005). Fls hose-in-hose,

with 5 petaloids, 3/flat truss, broadly funnel-shaped, 25 × 35mm, with 5+5+5 wavy-edged lobes, inside and out deep yellowish pink (47C), paling to white at base, with insignificant spotting on the dorsal lobe. Lvs elliptic, 25 × 15mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (147A) above, hairless. Slow-growing shrub, 0.5 × 0.5m in 9 years. Mid May.

(a) **'Li'l Sweetie Pie'**

cv. Evergreen azalea: ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'. H: W.L. Clagett (1996), G: W.L. Clagett (1999), N: W.L. Clagett (2005), I: White's Nursery (2006), REG: W.L. Clagett (2005). Fls hose-in-hose, with some petaloids, 2-3/flat truss, broadly funnel-shaped, 35 × 55mm, with 5+5+c.2 wavy-edged lobes, moderate purple (82C) in bud, opening with a moderate purple (82C) picotee edge, paling to nearly white at centre; outside similar except green at base; sparse, but large, basal, dorsal spots of reddish purple to ochre or chartreuse. Calyx c.2mm long, pale green. Lvs elliptic, 45 × 20mm, broadly acute at apex, cuneate at base, margins flat, moderate yellow-green (146B) above, with very fine, white hairs above and below only when young. Shrub 0.75 × 0.75m in 9 years. Mid May.

(a) **'Li'l Tae'**

cv. Evergreen azalea: ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'. H: W.L. Clagett (1996), G: W.L. Clagett (2000), N: W.L. Clagett (2005), I: White's Nursery (2006), REG: W.L. Clagett (2005). Fls hose-in-hose, with some petaloids, 2-3/flat truss, broadly funnel-shaped, 35 × 50mm, with 14-16 very wavy-edged lobes, deep purplish pink (68A) in bud, opening inside deep purplish pink (68A) at margins, shading to moderate and light purplish pink (68C-D) in the centre, with bright yellow-green at the base; outside deep purplish pink (68A); bright reddish purple markings on the three dorsal lobes. Lvs elliptic, 45 × 20mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate yellow-green (146B) above, with very fine white hairs on lower surface when young and when mature. Dense shrub, 0.75 × 0.6m in 9 years. Mid May.

(a) **'Lina Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-7/hemispherical truss, 30 × 27mm, with 5 wavy-edged lobes, deep yellowish pink (43C) in bud and on opening, both inside and out; three lobes with vivid reddish orange (34A) markings; stamens 5-6, exserted from the corolla, anthers brownish orange (166C) shaded moderate orange (166D), filaments vivid reddish orange (43B); pistil distinctly longer than the stamens, style strong yellowish pink (43D), stigma vivid reddish orange (43B). Lvs elliptic, 28 × 15mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate yellow-green (147B) above and moderate yellow-green (146B) below when mature; petioles 5-7mm long. Shrub 1.1 × 2.0m in 60 years

(regularly pruned). April. (Etymology: named for the mother of the nominant).

(r) **'Little Hope'**

cv. Elepidote rhododendron: 'Bellringer' (s) × unnamed yellow hybrid. H: D.B. Johnson (1995), G: D.B. Johnson (2000), N: D.B. Johnson (2005), REG: D.B. Johnson (2005). Fls 13/ domed truss, 50 × 65mm, with 5 wavy-edged lobes, deep pink (50B) in bud, opening light greenish yellow (8C) throughout; small, deep pink (50B) blotch at base; slightly scented. Lvs elliptic, broadly acute at apex, rounded at base, dull and moderate olive green (137B) above, hairless. Shrub 0.9 × 0.9m in 7 years. Mid May.

(r) **'Live Embers'**

cv. Elepidote rhododendron: 'The Honourable Jean Marie de Montague' (s) × *elliottii*. H: H. O'Rourke, G: H. O'Rourke (1980), N: H. O'Rourke (1990), REG: H. O'Rourke (2005). Fls 20/rounded truss, campanulate, 50 × 65mm, with 5 wavy-edged lobes, red in bud, opening vivid red (46B) inside and out. Calyx 5mm long, pink. Lvs lanceolate, 110 × 40mm, margins decurved, matt green above, cuneate at base, acute at apex. October (Australia).

(r) **'Lockington Maggie'**

cv. Elepidote rhododendron: 'Marcus of Lockington' (s) × 'Australian Sunset'. H: D.J. Dosser (1994), G: D.J. Dosser (2003), N: D.J. Dosser (2005), REG: D.J. Dosser (2005). Fls 12/ball truss, funnel-shaped, 50 × 75mm, with 6 wavy-edged lobes, deep pink (50B) in bud, opening deep pink (50B) inside and out, with a strong red (46A) blotch at the base of the corolla; slightly scented. Calyx 25mm long, strong purplish red (58B). Lvs elliptic, 140 × 40mm, margins decurved, matt green above, cuneate at base, apiculate at apex, hairless. Shrub 1.0 × 1.0m in 10 years. November to December (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0060825)).

(r) **'Lockington's Glow'**

cv. Elepidote rhododendron: 'Marcus of Lockington' (s) × 'Australian Sunset'. H: D.J. Dosser (1994), G: D.J. Dosser (2003), N: D.J. Dosser (2005), REG: D.J. Dosser (2005). Fls 11/conical truss, funnel-shaped, 40 × 115mm, with 7 wavy-edged lobes, deep purplish pink (68A) in bud, opening inside moderate purplish pink (68C), flushed pale yellow (11C), the tips of the lobes deep purplish pink (68A), with a conspicuous, solid blotch of strong purplish red (71B) all over the base of the corolla and extended shortly as rays and breaking up into spots; slightly scented. Calyx 75mm long, moderate purplish pink (68C). Lvs elliptic, 180 × 50mm, margins flat, matt green above, attenuate at base, apiculate at apex, hairless. Shrub 1.0 × 1.0m in 10 years. November to December (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0060826)).

(r) **'Lockington's Sunset'**

cv. Elepidote rhododendron: 'Marcus of Lockington'

(s) × 'Australian Sunset'. H: D.J. Dosser (1994), G: D.J. Dosser (2003), N: D.J. Dosser (2004), REG: D.J. Dosser (2005). Fls 14/ball truss, broadly funnel-shaped, 40 × 115mm, with 6 wavy-edged lobes, strong purplish red (58B) in bud, opening inside pale greenish yellow (10D) shading to pink at edges, outside streaked moderate purplish red (58A); with a blotch around the base of the throat, up to 12mm long, of moderate purplish red (59C), extended slightly up each lobe. Calyx 40mm long, moderate purplish red (58A). Lvs elliptic, 150 × 50mm, glossy green, attenuate at base, acute at apex, hairless. Shrub 1.0 × 1.0m in 10 years. November (Australia). (Nomenclatural standard: colour print provided by the registrant (WSY0070187)).

(r) **'Lonnie Foley'**

cv. Elepidote rhododendron: 'Scintillation' (s) × 'Lem's Cameo'. H: D.A. Griswold (1974), G: Griswold Nursery (1984), N: Griswold Nursery, I: Griswold Nursery, REG: L. Malland (2005). Fls 14/lax truss, open funnel-campanulate, 50 × 25mm, with 5 wavy-edged lobes, strong yellowish pink (38A) in bud, opening very pale purple (69C), edged very pale purple (69A), the throat pale greenish yellow (2D) with short, spikey, deep yellowish pink (43C) markings in the centre; stigma green; anthers dark. Lvs elliptic, 100 × 40mm, acute at apex, cuneate at base, margins wavy, glossy and strong yellow-green (144A) above, hairless. Shrub with an upright and spreading habit, 1.5 × 1.5m in 30 years. May.

(r) **'Macarena'**

cv. Elepidote rhododendron: ('Hotei' × ('Goldrausch' × *wardii* hybrid)) (s) × Fred Wynniatt Group. H: H. Hachmann (1990), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 9-11/rather lax truss, broadly funnel-shaped, 55-60 × 90-105mm, with (5-)-7(-8) moderately wavy-edged lobes, inside and out brilliant yellow (11A), including the throat, the edges strong/moderate yellowish pink (38A/B); the three dorsal lobes with an area of intensified vivid/brilliant yellow (13A/B) and with markings of strong red (46A) deep in the throat; stamens 14, filaments pale greenish yellow (2D), anthers light reddish brown (177B); styles light orange-yellow (22C), stigma moderate orange-yellow (164B), ovary moderate yellow-green (147B). Calyx 10-30mm long, coloured as the corolla. Lvs ovate, 100 × 52mm, margins flat/slightly recurved, matt light green above, cordate at base, apiculate at apex, hairless. Shrub c.1.5 × c.1.7m in 10 years; habit broadly rounded. Early to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070138)). Published in Hachmann cat. 2004-2008 (2004), p.58.

(r) **'Maddalena'**

cv. Elepidote rhododendron: ('Kokardia' × ('Hachmann's Diadem' × 'Holger')) (s) × 'Sarasate'. H: H. Hachmann (1993), G: H. Hachmann (1999), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 10-18/fairly compact

truss, broadly funnel-shaped, 50-65 × 75-80mm, with 5 slightly wavy-edged lobes, buds strong purplish red (60D), opening inside strong purplish pink (67D), the centre of the lobes and the throat more-or-less whitish, lobe margins shaded deep purplish pink (67C), outside similarly coloured, but with midveins shaded vivid purplish red (67B); dorsal lobe with deep red (187A/B) markings; stamens 10, filaments white, anthers dark pink (182D); styles whitish/green, shaded strong purplish red/deep purplish pink (58C/D) near the apex, stigma dark red (59A); ovary reddish green. Calyx 1-3mm long, light purplish pink/very pale purple (73C/D). Lvs elliptic, 90-110 × 40-45mm, margins slightly waved and decurved, dark glossy green above, cuneate at base, apiculate at apex, with some hairs. Shrub c.1.4 × c.1.7m in 10 years; habit broadly rounded. Late May to mid June. (Nomenclatural standard: colour illustration provided by the registrant (WSY0070139). Published in Hachmann cat. 2004-2008 (2004), p.58).

(r) **'Maggie'**

cv. *Lepidote* rhododendron: 'Saint Merryn' (s) × 'Azurika'. H: Holger Hachmann (1992), G: Holger Hachmann (1998), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 2-5(-6)/individual truss (these gathered in a rounded ball inflorescence of 14-24 fls), rotate (to broadly funnel-shaped), c.20 × 23-30mm, with 5 wavy-edged lobes, opening light violet (89D) inside and out, with inconspicuous markings; stamens 6-10, filaments strong/brilliant violet (86C/D), anthers dark red (187B); style vivid reddish purple (74A/B), stigma dark red (187A). Calyx 3-5mm long, strong yellow-green (144B/C), silvery hairy. Lvs elliptic, 20 × 8mm, margins flat, glossy dark green above, cuneate at base, acute (to apiculate) at apex, with dark green scales below when mature. Shrub c.0.35 × c.0.6m in 10 years, with a dense, flat-rounded habit. Late April to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070159). Published in Hachmann cat. 2004-2008 (2004), p.132.).

(a) **'Maischnée'**

cv. Evergreen azalea: 'Elsie Lee' (s) × 'Purpurkissen'. H: H. Hachmann (1988), G: H. Hachmann (1992), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls single, 2-3/truss, open funnel-shaped, 35-40 × 65-75mm, with 5(-6) distinctly wavy-edged lobes, opening white (155D) throughout; dorsal lobe with soft markings of brilliant/light yellow-green (150C/D); stamens 8-9, some tending to be slightly petaloid, filaments white, anthers brownish; style light yellow-green (150D), stigma yellowish green. Calyx 3-4mm long, strong yellow-green (144B), silvery hairy. Lvs ovate, c.25-30(-40) × 12-22mm, margins flat, matt mid green above, cuneate at base, apiculate at apex, with only a few brownish hairs. Shrub c.0.4 × c.0.8m in 10 years, with a low-growing, broadly compact habit. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070164). Published in Hachmann cat. 2004-2008 (2004), p.169).

(r) **'Makayla'**

cv. *Elepidote* rhododendron: 'Blimey Limey' (s) × 'O'Harold'. H: M. Stockman (1993), G: M. & H. Stockman (2001), N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 4-5/domed truss, open funnel-shaped, 65 × 100mm, with 5 very wavy-edged lobes, strong purplish red (61B) in bud, opening inside and out deep purplish pink (66C), the inside with a solid dark red (59A) centre and speckled with the same colour on the dorsal lobe to within c.15mm of the edge, the speckling spreading onto adjacent lobes; outside deep purplish pink (66C) with a strong purplish red (61B) base and a narrow stripe up the centre of each lobe. Calyx lobes irregular, 35-50mm long, coloured as the corolla. Lvs elliptic, 120 × 50mm, margins flat, broadly acute at apex, rounded at base, margins downcurved, dark yellowish green (139A) above, moderate yellow-green (147C) below, hairless. Shrub 1.0 × 1.0m in 11 years. Early May.

(r) **'Marcus Agius'**

cv. *Elepidote* rhododendron: 'Crest' (s) × Loderi Group. H: E. de Rothschild, G: E. de Rothschild, N: E. de Rothschild (2004), REG: E. de Rothschild (2005). Fls 10-12/loose truss, funnel-shaped, 70 × 110-120mm, with 7 wavy-edged lobes, buds strong yellowish pink (31C), opening inside and out light greenish yellow (4B) in the centre, fading to pale yellow-green (4D) on lobes, sometimes flushed strong yellowish pink (37B); with a small, strong red (51A) blotch at the base of the throat; stamens in 2 groups of 7, filaments cream, anthers brown; style cream, stigma strong yellow-green (145A), distinctly exceeding the stamens. Calyx 8mm long, strong yellow-green (145A), edged deep yellowish pink (47C). Lvs elliptic, 140 × 60mm, margins flat, glossy above, obtuse at base and apex, hairless. Shrub 3.5 × 2.5m in c.20 years, with a loose habit. Mid to late May.

(r) **'Marengo'**

cv. *Elepidote* rhododendron: 'Ovation' (Wüstemeyer) (s) × 'Erato' (Hachmann). H: H. Hachmann (1987), G: H. Hachmann (1993), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 14-16/rather loose truss, broadly funnel-campanulate, 40-45 × 50-60mm, with 5 wavy-edged lobes, deep/strong red (53A/B) both inside and out; dorsal spotting of dark red (187A); stamens 10, filaments pale purplish pink (56D), anthers greyish brown (166A); style deep yellowish pink (39B), stigma dark red (183B); ovary greenish, silvery hairy. Calyx with 2 lobes 2mm long, the others apparently lacking, deep red (53A). Lvs ovate to elliptic, 90-140 × 35-70mm, margins flat, matt deep green above, rounded to slightly oblique at base, acute at apex, with fawn-coloured hairs below. Shrub c.1.1 × c.1.6m in 10 years; habit dense, broadly compact to rounded. Mid to late May. (Nomenclatural standard: colour print provided by the registrant (WSY0070160). Published in Hachmann cat. 2004-2008 (2004), p.129).

(r) 'Marianka'

cv. Elepidote rhododendron: ('Kokardia' × ('Hachmann's Diadem' × 'Holger')) (s) × 'Sarasate'. H: H. Hachmann (1993), G: H. Hachmann (1999), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 14-16/dense, ball truss, broadly open funnel-shaped, 50-55 × 65-75mm, with 5 wavy-edged lobes, buds vivid purplish red (61C), opening inside strong purplish pink (67D), centre of lobes and throat more-or-less white, margins shaded vivid purplish red (61C); outside similarly coloured but veins shaded strong/vivid purplish red (67A/B); with a conspicuous blotch and markings on the three dorsal lobes of strong red (46A); stamens 10, filaments white, anthers greyish red (182B); style light yellow-green (150D), stigma dark red (183B); ovary light olive grey (197B), white hairy. Calyx 2-5mm long, greenish purple. Lvs ovate, c.120 × 45-50mm, margins flat, slightly wavy, dark glossy green above, rounded at base, apiculate at apex, hairless. Shrub c.1.1 × c.1.3m in 10 years; habit broadly rounded. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070165). Published in Hachmann cat. 2004-2008 (2004), p.61).

(a) 'Marinja'

cv. Evergreen azalea: 'Elsie Lee' (s) × 'Purpurkissen'. H: H. Hachmann (1988), G: H. Hachmann (1992), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls double, 1-2(-3)/truss, with 3 whorls one on top of the other, 25-30 × 45-50mm, with a total of 17-21 slightly wavy-edged lobes, strong/moderate purplish pink (68B/C) inside and out, dorsal lobes with strong purplish red (60B) markings; stamens mostly petaloid, filaments pale purplish pink (65C), anthers moderate purplish red (70A); style strong/moderate purplish pink (68B/C), stigma deep purplish red (71A); ovary greenish, silvery hairy. Calyx always present, 3mm long, strong yellow-green (144C). Lvs narrowly elliptic to obovate, 20-25 × 12mm, margins flat, dark green and slightly glossy above, cuneate at base, acuminate at apex, with brownish hairs. Shrub c.0.45 × c.0.6m in 10 years; habit very compact, dense, rounded. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070165). Published in Hachmann cat. 2004-2008 (2004), p.170).

(r) 'Marylou'

cv. Elepidote rhododendron: 'Paprika Spiced' (s) × 'Kalamaika'. H: H. Hachmann (1992), G: H. Hachmann (1995), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 7-10/compact truss, broadly funnel-shaped, 35-45 × 55-65mm, with 5 wavy-edged lobes, inside and out pale orange-yellow (19D), edges shaded light yellowish pink to pale yellowish pink (36A-D), centre of lobes and throat pale yellow (18C/D); dorsal lobe with strong red (46A) markings; stamens 10, filaments pale yellow-green (4D), anthers moderate reddish brown/dark reddish brown (175A/B); style light yellow (14D), stigma moderate red (183C). Calyx 2-2.5mm long, yellowish/red. Lvs elliptic, 70-80 × 30-

35mm, margins slightly decurved, dark matt green above, cuneate at base, apiculate at apex, hairless. Shrub c.0.6 × c.1.0m in 10 years; habit broadly compact, low-growing. Early to late May. (Nomenclatural standard: colour print provided by the registrant (WSY0070142). Published in Hachmann cat. 2004-2008 (2004), p.61).

(r) 'Maspaloma'

cv. Elepidote rhododendron: ('Kokardia' × ('Hachmann's Diadem' × 'Holger')) (s) × 'Sarasate'. H: H. Hachmann (1993), G: H. Hachmann (1999), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 12-15/fairly compact truss, broadly open funnel-shaped, 45-50 × 70-75mm, with (5-)6-7 wavy-edged lobes, buds vivid purplish red (61C), opening inside with centre of lobes and throat deep purplish pink (66C), midveins slightly paler, margins vivid purplish red (61C), outside similarly coloured with the midveins tinged vivid purplish red (66B); dorsal lobes with a large, conspicuous blotch and markings of dark red (187A/B); stamens 13-18, some slightly petaloid, filaments glassy white, anthers dark pink (182D); style pale yellow-green (149D), stigma dark red (187B); ovary greenish, white hairy. Calyx 3-5mm long, reddish/green. Lvs ovate, sometimes twisted, 100-110 × 45-50mm, margins flat/slightly recurved, dark glossy green above, with yellowish green petioles and midveins, cuneate at base, apiculate at apex, with inconspicuous hairs. Shrub c.1.2 × c.1.4m in 10 years; habit broadly compact. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070143). Published in Hachmann cat. 2004-2008 (2004), p.61).

(a) 'May Young'

cv. Evergreen azalea: *nakaharae* (s) × 'Rosebud' (Gable). H: G. Griswold (1974), G: G. Griswold (1983), N: Griswold Nursery, I: Griswold Nursery, REG: Griswold Nursery (2005). Fls 3/lax truss, broadly funnel-shaped, c.30-40 × c.40-45mm, with 5 wavy-edged lobes, and 1-5 petaloids, strong purplish red (63A). Calyx 13mm long, vivid purplish red (66B). Lvs elliptic, 20 × 6mm, acute at apex, cuneate at base, margins flat, shiny and moderate olive green (147A) above, hairy above. Shrub 0.6 × 0.9m in 10 years. May.

(r) 'Mead'

cv. Elepidote rhododendron: 'Sunspray' (s) × 'Lem's Cameo'. H: J. Olson, G: J. Olson (2005), N: J. Olson (2005), REG: J. Olson (2005). Fls 12/domed truss, broadly funnel-shaped, 75 × 115mm, with 7 very wavy-edged lobes, light yellowish pink (36A) in bud, opening inside light yellowish pink (36B), shading to pale yellow (18C) in centre; outside light yellowish pink (36A) with pale yellowish pink (27C) at margins; unmarked; slightly scented; style long, with a large stigma. Calyx 40mm long, pale yellow (12D), edged red. Lvs oblong, 175 × 75mm, obtuse at apex, cordate at base, margins flat, semi-glossy and dark green (136A) above, hairless. Shrub 1.5 × 1.5m in 6 years, with flowers of very heavy substance. Mid May.

(a) 'Meina'

cv. Evergreen azalea: parentage unknown. R: Savioli Fratelli (1970), N: Savioli Fratelli (1985), I: Savioli Fratelli (1985), REG: Prof. E. Accati (2005). Fls single, 2-10/hemispherical truss, 42 × 28mm, with 5 very wavy-edged lobes, buds strong red (53B), opening strong red (53D) throughout; three lobes with deep red (53A) markings; stamens 5-7, exerted from the corolla, anthers greyish reddish brown (200B), filaments deep pink (53C); pistil exceeding stamens, style strong red (53C), stigma strong red (53B). Lvs elliptic, 24 × 12mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (147A) above and moderate yellow-green (147C) below when mature; petioles 5mm long. Shrub 1.0 × 1.0m in 10 years. April. (Etymology: named for a village near the city of Verbania; the latter is where the Savioli nursery is situated).

(r) 'Melody Bell'

cv. Elepidote rhododendron: parentage unknown. G: D. Dean (c.1988), S: D.S. Bell (2003), N: D.S. Bell (2003), REG: D.S. Bell (2005). Fls 10-12/domed truss, hose-in-hose, funnel-campanulate, 75 × 65mm, with 5 wavy-edged lobes, dark red (187C) in bud, opening inside with a dark red (187C) border, c.15mm wide, changing rather abruptly to white, with sparse, yellow, dorsal spotting; between the red margin and white star-shaped centre there is a narrow transition zone of feathery red markings; outside dark red (187C). Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins flat, dull green above, hairless, Shrub 1.8 × 1.8m in 20 years. Mid May.

(a) 'Mergozzo'

cv. Evergreen azalea: parentage unknown. R: Fratelli Savioli (1970), N: Fratelli Savioli (1985), I: Fratelli Savioli (1985), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 38 × 24mm, with 5 flat or slightly wavy-edged lobes, deep pink (52C) in bud and on opening, both inside and out; vivid red (52A) markings on three lobes; stamens 5, exerted from the corolla, anthers dark greyish yellow (199D), filaments deep pink (52B); pistil distinctly longer than the stamens, style deep pink (52B), stigma vivid red (46B). Lvs obovate, 28 × 13mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (146A) above and strong yellow-green (144B) below when mature; petioles 5mm long. Shrub 1.0 × 1.0m in 10 years. April. (Etymology: named for a village near the city of Verbania; the latter is where the Savioli nursery is situated).

(a) 'Miss Curlicue'

cv. (Evergreen azalea: ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'. H: W.L. Clagett (1996), G: W.L. Clagett (2000), N: W.L. Clagett (2005), I: White's Nursery (2006), REG: W.L. Clagett (2005). Fls hose-in-hose, with 5 petaloids, 3/flat truss, broadly funnel-shaped, 35 × 45mm, with 5+5 wavy-edged lobes, deep pink (50B) in bud, opening inside with an irregular edging of strong red (50A), shading to deep pink (50B) and then white, the centre bright yellow-green; outside

strong red (50A) fading to yellow at base; yellow ochre blotch on dorsal lobe, speckled with a few purple dots. Calyx 3mm long. Lvs elliptic, 25 × 13mm, broadly acute at base, margins flat, semi-glossy above, with very fine, white hairs below. Dense shrub, 0.6 × 0.6m in 9 years. Mid to late April.

(r) 'Miss Frances'

cv. Elepidote rhododendron: ('Yaku Sunrise' × 'Burgundy') (s) × 'Noyo Chief'. H: F. Fujioka (1988), G: F. Fujioka (1993), N: The children of Frances Maxwell (2005), I: Meerkerk Rhododendron Gardens (2005), REG: J. Neel (2005). Fls 20/conical truss, funnel-campanulate, 50 × 50mm, with 5 wavy-edged lobes, buds vivid red (45C) at base, blending to strong red (46A) at apex, opening inside vivid red (45C) with vivid red (46B) margins; outside vivid red (45B) with vivid red (45A) margins; unmarked. Lvs elliptic, 140 × 50mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and dark yellowish green (139A) above, with a plastered indumentum below (moderate yellow (161B) when young and maturing to moderate orange-yellow (165C)). Shrub 1.8 × 1.5m in 15 years, with red stems. Fls almost identical to 'Taurus'. Early April.

(r) 'Mogambo'

cv. Elepidote rhododendron: 'Rasputin' (s) × 'Monsieur Marcel Ménard'. H: Holger Hachmann (1989), G: Holger Hachmann (1995), N: Holger Hachmann (2003), I: Holger Hachmann (2005), REG: G. Stück (2005). Fls 10-12/firm truss, broadly open funnel-shaped, 55-60 × 65-80mm, with 5 wavy-edged lobes, buds deep purplish red (71A) shaded deep reddish purple (77A), opening inside strong purple (77B), edges tinged deep reddish purple (77A), lobe centres and veins shaded strong purplish red/strong reddish purple (72A/B); outside similarly coloured; dorsal lobe with a large, conspicuous blotch and markings of dark red (187A/B), the blotch area almost warty; stamens 10, filaments strong purplish pink (67D), anthers pale yellow (162D); style strong purplish red (58C), stigma dark red (187B); ovary light olive grey (197A). Calyx c.11mm long, reddish/greenish. Lvs elliptic, slightly convex, 110-180 × 35-50mm, margins flat, dark and slightly glossy green above, rounded at base, acute at apex, with inconspicuous, sparse, brownish hairs below. Shrub c.1.6 × c.1.9m in 10 years, with a broadly rounded habit. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0070144). Published in Hachmann cat. 2004-2008 (2004), p.64.).

(r) 'Morna's Piccolo'

cv. Elepidote rhododendron: 'Murray's Pink' (s) × 'Roy Stockman'. H: M. Stockman (1993), G: M. & H. Stockman (2002), N: H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 6-10/lax truss, campanulate, 40 × 50mm, with 6 flat-edged lobes, deep red (53A) in bud, opening strong reddish purple (70B) throughout, with a small, deep red (53A) spot at the base and no

other markings. Calyx c.1.5mm long, dark greenish yellow (152D). Lvs orbicular, 75 × 50mm, obtuse at apex, cordate at base, margins flat, dull and moderate olive green (147A) above, moderate yellow-green (148A) below, hairless. Shrub 0.6 × 0.75m in 11 years. Early April.

(r) **'Morning Dew'**

cv. Elepidote rhododendron: 'Mrs J.C. Williams' (s) × 'Plum High'. H: J. Barlup (1993), G: J. Barlup (1998), N: J. Barlup (2005), REG: J. Barlup (2005). Fls 16/ball truss, broadly funnel-shaped, 55 × 75mm, with 5 wavy-edged lobes, light purple (77C) in bud, opening throughout very pale purple (76D), the inside with slightly deeper tones of light purple (77D) in the throat and a flare (25 × 20mm) of discrete, moderate red (184B) dorsal spots; outside with light purple (77D) midribs; entire corolla fades to yellowish white (155D); the prominent filaments and style are purplish pink. Lvs elliptic, 90 × 45mm, broadly acute at apex, cuneate at base, margins flat, dull and moderate olive green (147A) above, hairless. Dense shrub, 0.9 × 1.2m in 12 years. Late May.

(r) **'Nancy Ledyard'**

cv. Lepidote rhododendron: 'Else Frye' (s) × 'Else Frye'. H: F. Renick (1995), G: F. Renick (2000), N: F. Renick (2005), REG: E.W. Moyles (2005). Fls 3/lax truss, open to broadly funnel-shaped, 65 × 75mm, with 5 wavy-edged lobes, white overall, with pink to white markings - ranging from random streaks to partial edging on lobes both inside and out; prominent orange/yellow flare in the dorsal throat. Calyx up to 3mm long, strong purplish red, hairy. Lvs elliptic, 75 × 25mm, semi-glossy green above, with dense, tiny, pale green scales below. Shrub 1.8 × 1.2m in 10 years. Mid April.

(a) **'Neve Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 47 × 37mm, with 5 wavy-edged lobes, white in bud and throughout on opening, unmarked; stamens 5, exerted from the corolla, anthers strong orange-yellow (22A), filaments white; pistil exceeding stamens, style brilliant greenish yellow (1B), shaded white, stigma brilliant greenish yellow (1B). Lvs obovate, 30 × 13mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (146C) below when mature; petioles 7-9mm long. Shrub 1.0 × 1.5 m in 60 years (regularly pruned). May. (Etymology: colour of the flowers resembles snow ("neve" in Italian)).

(r) **'Neville's Surprise'**

cv. Elepidote rhododendron: parentage unknown. S: N. Cleland (1996), G: N. Cleland (2000), N: N. Cleland (2004), REG: N. Cleland (2005). Fls 18-19/ ball truss, broadly funnel-shaped, 80mm wide, with 5 very wavy-edged lobes, buds off white shot with pink, opening white with a broad shading of strong

reddish purple (72C) around the corolla margin, outside deep purplish pink (72D); dorsal lobe heavily blotched vivid orange-yellow (23A). Calyx 3mm long, brownish green. Lvs elliptic, 120 × 45mm, margins flat, cuneate at base, acute at apex, hairless. Shrub 1.2 × 1.5m in 6 years. October to November (New Zealand). (Etymology: derived from the registrant's christian name and the chance origin of the plant).

(r) **'Nova Sunrise'**

cv. Elepidote rhododendron: ((*fortunei* × *smirnowii*) × (*yakushimanum* × 'Goldsworth Yellow')) (s) × (*awreum* × 'Prelude'). H: D.L. Craig (1988), G: D.L. Craig (1994), N: D.L. Craig & S. Gunn, REG: D.L. Craig & S. Gunn (2005). Fls 12-14/domed truss, open funnel-shaped, 40 × 60mm, with 5 slightly wavy-edged lobes, buds light yellow suffused strong yellowish pink (37B), opening inside light greenish yellow (5D), with light yellow-green (145B) flecks in the upper throat and scattered moderate red (180C) spots in the lower throat; outside light greenish yellow (5D), very lightly suffused with strong yellowish pink (37B). Calyx 3mm long, light greenish yellow (5D), suffused strong yellowish pink (37B). Lvs elliptic, 70 × 30mm, broadly acute at apex, cuneate at base, margins downcurved, semi-glossy and moderate yellow-green (147B) above, hairless. Shrub with a dense, compact habit, 0.6 × 0.9m in 11 years. Early May.

(r) **'November Moon'**

cv. Elepidote rhododendron: 'Unique' (s) × 'Jack's White'. H: D.J. Dosser (1994), G: D.J. Dosser (2002), N: D.J. Dosser (2004), REG: D.J. Dosser (2005). Fls 15/ball truss, funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, buds light greenish yellow (3D), opening pale yellow-green (4D) inside and out, without markings. Lvs broadly elliptic, 125 × 40mm, matt green, attenuate at base, mucronate at apex, hairless. Shrub 0.75 × 1.0m in 10 years. October (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0070182)).

(r) **'Noyo Charm'**

cv. Lepidote rhododendron: (*johnstoneanum* × *veitchianum*) (s) × *moupinense*. H: R.W. Scott (1969), G: R.W. Scott (1974), N: K. Jones (2005), I: Forest Lane Nursery (2005), REG: K. Jones (2005). Fls 2-5/lax truss, open funnel-shaped, 65 × 90mm, with 5 very wavy-edged lobes, inside white, randomly tinged light purplish pink (55C) and with prominent strong purplish red (54A) spotting at the base of the dorsal lobes; outside white tinged strong pink (48D), mainly at midribs. Calyx c.5mm long, moderate yellow-green (147D). Lvs obovate, 55 × 25mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (147A) above, with moderate yellow-green (146A) hairs and scales on both surfaces of leaves and on calyx. New growth dark green to reddish bronze; the bark of mature plants turns mahogany and then peels. Shrub 1.5 × 1.8m in 15 years. March to April.

(r) 'Orange Flirt'

cv. *Elepidote rhododendron*: 'Ken Janeck' (s) × unnamed hybrid of *dichroanthum* subsp. *scyphocalyx*. H: Holger Hachmann (1992), G: Holger Hachmann (1996), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 9-11/rather lax truss, funnel-campanulate, 50 × 50-65mm, with 5 wavy-edged lobes, inside pale orange-yellow (24D), edger moderate pink (35D), outside moderate yellowish pink (33D), tinged strong yellowish pink (38A), veins slightly darker; dorsal lobe with vivid reddish orange (33A/B) markings; stamens 4-5, c.10mm long, poorly developed, held deep in the throat, filaments greenish white (157D), anthers moderate orange (167A/B); style light yellow-green (154D), stigma brownish orange (165B). Calyx 12-15mm long, held horizontally, pale orange-yellow (24D) inside, light yellowish pink (27A) outside. Lvs elliptic, 90-100 × 35-45mm, margins flat, mid green and fairly glossy above, edges, venation and petioles yellow-green, cuneate at base, acute at apex, hairless. Shrub c.0.5 × c.0.7m in 10 years; habit dense; slow-growing. Mid to late May. (Nomenclatural standard: colour print provided by the registrant (WSY0070145). Published in Hachmann cat. 2004-2008 (2004), p.66).

(a) 'Paolo Pejrone'

cv. Evergreen azalea: parentage unknown. N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single, 1(-2)/truss, 53 × 35mm, with 5 flat or rarely wavy-edged lobes, buds vivid red (52A), opening vivid red (52A) inside and out; three lobes with markings of deep red (53A) shaded with strong red (53B); stamens 8-10, some exserted from the corolla, anthers purple (N79B), filaments deep pink (52B); pistil exceeding the stamens, style deep pink (52B), stigma orange-red (N34D). Lvs lanceolate, 37 × 14mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (146B) below when mature; petioles 6mm long. Shrub 1.7 × 2.0m in 100 years (regularly pruned). May. (Probably raised (c.1890) by Fratelli Rovelli and Fratelli Contini, gardeners at the Villa del Pascià in the 19th century; this villa is where the then King of Egypt stayed during his Italian holidays on Lake Maggiore. Etymology: named for one of the leading Italian landscape architects).

(r) 'Peggy'

Syn. of 'Hachmann's Peggy'.

(a) 'Pescherino'

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 2-3/truss, 67 × 45mm, with 5 wavy to very wavy-edged lobes, buds moderate purplish pink (62B), opening light purplish pink (62C) shaded strong purplish red (58B); three lobes with moderate purplish red (58A) markings; stamens 10, not exserted from the corolla, anthers dark red (59A), filaments white shaded vivid purplish red (61C); pistil

exceeding stamens, style strong red (53D) shading to strong purplish pink (55B) at the base, stigma dark red (59A) shaded deep yellowish pink (47C). Lvs lanceolate, 59 × 24mm, margins flat, rarely wavy, glossy above, acute at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (146B) below when mature; petioles 4mm long. Shrub 1.8 × 1.5m in 50 years. April to May. (Etymology: local name for a place near Cannero Riviera, the latter being where the Carmine nursery is situated).

(r) 'Petite Nancy Jean'

cv. *Lepidote rhododendron*: sport from the PJM Group. S: Corneliusen's Nursery (1998), G: Corneliusen's Nursery, N: Corneliusen's Nursery, I: Corneliusen's Nursery (2005), REG: Corneliusen's Nursery (2005). Fls 4/ball truss, open funnel-shaped, 40 × 25mm, with 5 wavy-edged lobes, vivid reddish purple (78A) in bud, opening inside light purple (77D); outside strong purple (77B), fading to light purple (77D); unmarked; style red. Lvs elliptic, 13-25 × 6-13mm, obtuse at apex, cuneate at base, margins slightly downcurved, glossy and moderate olive green (147A) above, dark greyish purple (202A) in winter, scaly below. Compact shrub, 40-45 × 50cm in 5 years. March.

(r) 'Philip Dee'

cv. *Elepidote rhododendron*: 'Casanova' (s) × 'Peter Alan'. H: J. Doppel (1995), G: J. Doppel (2000), N: J. Doppel (2004), REG: J. Doppel (2005). Fls 12/ball truss, broadly funnel-shaped, c.50 × 65mm, with 5 flat to slightly wavy-edged lobes, pinkish lavender in bud, opening inside light cream with a prominent, dark maroon, dorsal blotch; outside cream with some lavender tinges. Lvs obovate, 100 × 50mm, obtuse at apex, rounded at base, margins moderately downcurved, semi-glossy and mid green above, hairless. Shrub 0.9 × 1.2m in 10 years. Late May.

(r) 'Pinguin'

Syn. of 'Hachmann's Pinguin'.

(a) 'Premeno'

cv. Evergreen azalea: parentage unknown. R: Fratelli Savioli (1970), N: Fratelli Savioli (1985), I: Fratelli Savioli (1985), REG: Prof. E. Accati (2005). Fls single, 3-4/truss, 15 × 30mm, with 5 very wavy-edged lobes, buds strong reddish purple (78B), opening moderate purplish pink (78D) inside and out; three lobes with moderate red (47A) markings; stamens 7-9, exserted from the corolla, anthers deep reddish purple (77A), filaments strong reddish purple (78B); pistil just exceeding stamens, style pale yellow (11D), stigma brilliant greenish yellow (1B). Lvs lanceolate-obovate, 36 × 15mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate yellowish green (138A) above and moderate yellow-green (138C) below when mature; petioles 5mm long. Shrub 1.0 × 1.0m in 10 years. May. (Etymology: named for a village near the city of Verbania, the latter being where the Savioli nursery is situated).

(r) 'Princess Angeline'

cv. Elepidote rhododendron: 'Chevalier Félix de Sauvage' (s) × 'Purple Splendour'. H: Mrs E.M. Watson (1964), G: Mrs E.M. Watson (1975), N: Mrs E.M. Watson, REG: Mrs E.M. Watson (2005). Fls 23/ball truss, open funnel-shaped, 75mm wide, with 5 wavy-edged lobes, strong reddish purple (70B) throughout, with a prominent, darker, speckled, dorsal blotch. Calyx 65mm long, strong purplish red (59D). Lvs elliptic, 105 × 30mm, acute at apex, cuneate at base, with very slightly downcurved margins, dull and greyish yellow-green (191A) above, hairless. Shrub with a spreading habit, 0.5 × 0.75m in 30 years. Midseason.

(r) 'Pure Fire'

cv. Elepidote rhododendron: 'Anna' (s) × Antonio Group. H: H. O'Rourke (1988), G: H. O'Rourke (1997), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 20/ball truss, funnel-shaped, 60 × 80mm, with 5 wavy-edged lobes, strong red (46A) in bud and both inside and out on opening. Calyx 3mm long, red. Lvs oblanceolate, 150 × 80mm, margins decurved, matt green above, cuneate at base, acute at apex, with a scant, fawn indumentum when young. Shrub 1.0 × 1.2m in 14 years. September to October (Australia).

(r) 'Rachel Foster'

cv. Elepidote rhododendron: selection from *ungermii*. G: L. de Rothschild, N: E. de Rothschild (2005), REG: E. de Rothschild (2005). Fls 20-24/tall, domed, more-or-less compact truss, funnel-campanulate, 30 × 45mm, with 5 slightly wavy-edged lobes, buds pale purplish pink (62D) at base, gradually becoming strong purplish red (58B) along lobe edges, opening inside pale purplish pink (62D), flushing to deep purplish pink (61D) at the edges of the lobes and with the throat and centre of lobes remaining pale, outside pale purplish pink (62D) at base, flushing up to deep purplish pink (61D) unevenly over most of the tube and lobes; dorsal lobe lightly spotted with strong yellow-green (142A). Calyx 4mm long, light yellow-green (150D). Pedicels rose pink. Lvs oblanceolate, 110-150 × 40-55mm, margins flat, matt green above, narrowly cuneate at base, obtuse at apex, with hairs below (pale orange-yellow (159B) at first, aging to moderate orange-yellow (164C)). Shrub to at least 3.0 × 3.0m, with a rounded open habit (can be straggly). July to August. (Trusses smaller than usual in this species, on short stems; stamens and style shorter and darker than usual. Nomenclatural standard: colour print provided by the registrant (WSY0070475)).

(r) 'Radiant Beauty'

cv. Elepidote rhododendron: 'Coronation Day' (s) × 'Ruby Bowman'. H: H. O'Rourke (1990), G: H. O'Rourke (1999), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 18/loose truss, broadly funnel-shaped, 55 × 90mm, with 6 wavy-edged lobes, buds deep red (53A), opening inside and out strong red (53B). Calyx 1mm long, green. Lvs lanceolate, 140 × 45mm, margins decurved, matt green above, obtuse

at base and apex, hairless. Shrub 1.8 × 1.5m in 12 years. October (Australia).

(r) 'Red Lord'

cv. Elepidote rhododendron: *arboreum* (s) × 'Bernard Gill'. H: H. O'Rourke (1989), G: H. O'Rourke (2000), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 18-19/domed truss, funnel-shaped, 50 × 60mm, with 5 smooth-edged lobes, strong red (46A) in bud and strong red (46A) both inside and out on opening. Calyx 2mm long, yellowish green. Lvs lanceolate, 110 × 35mm, margins flat, matt green above, obtuse at base, acute at apex, with a grey, plastered indumentum below. Shrub 1.0 × 1.2m in 13 years. September to October (Australia).

(a) 'Red Panda'

cv. Evergreen azalea: 'Squirrel' (s) × 'Red Red'. H: K.N.E. Cox (1995), G: Glendoick Gardens (1998), N: K.N.E. Cox (2005), I: Glendoick Gardens (2005), REG: Glendoick Gardens (2005). Fls single, 1-2/truss, open funnel-campanulate, 30 × 37mm, with 5 slightly wavy-edged lobes, strong red (46A) in bud and on inside on opening, outside slightly darker; unmarked; stamens and style red. Calyx 4-5mm long, green. Lvs broadly elliptic, up to 20 × 15mm, margins upcurved, glossy green above, cuneate at base and apex, hairless. Shrub 0.3 × 0.5m in 5 years. May to June.

(r) 'Regality'

cv. Elepidote rhododendron: Calstocker Group (s) × 'Irene Stead'. H: H. O'Rourke (1989), G: H. O'Rourke (1999), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 14/flat-topped truss, broadly campanulate, 65 × 100mm, with 5 wavy-edged lobes, buds pink, opening inside and out pale pink, which fades to white. Calyx 7mm long, pink. Lvs ovate, 140 × 55mm, margins decurved, glossy green above, cuneate at base, broadly acute at apex, hairless. Shrub 1.0 × 1.2m in 14 years. October (Australia).

(r) 'Reileyridge Bright Day'

cv. Elepidote rhododendron: 'Janet Blair' (s) × 'Ed's Red'. H: H.E. Reiley (1983), G: H.E. Reiley, N: H.E. Reiley, REG: H.E. Reiley (2005). Fls 17/conical truss, saucer-shaped, c.70 × 90mm, with 5 wavy-edged lobes, deep purplish pink (55A) in bud, opening inside strong purplish pink (55B) with a strong orange-yellow (24B) dorsal flare lightly speckled with small, darker spots; outside strong purplish pink (55B). Lvs elliptic, 125 × 50mm, broadly acute at apex, cuneate at base, margins flat, dull and dark green above, hairless. Shrub 1.8 × 1.2m in 10 years. Midseason.

(r) 'Reileyridge Double Delight'

cv. Elepidote rhododendron: 'Evening Glow' (s) × 'Janet Blair'. H: H.E. Reiley (1987), G: H.E. Reiley (pre 1995), N: H.E. Reiley, REG: H.E. Reiley (2005). Fls 10/ball truss, broadly funnel-shaped, c.70 × 90mm, with 7 wavy-edged lobes, strong purplish pink (55B) in bud, opening inside light yellow (18A) with

pink spotting on the dorsal lobe; outside pink and yellow; moderately scented. Calyx c.10mm long, yellow, with pink margins. Lvs oblong, 125 × 55mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and dark green above, hairless. Dense shrub, 1.2 × 1.2m in 10 years. Midseason. (Etymology: "double" in the epithet refers to both the flowers and the foliage being a delight).

(r) **'Reileyridge Picotee'**

cv. Elepidote rhododendron: 'Janet Blair' (s) × 'Donna's Day'. H: H.E. Reiley (1983), G: H.E. Reiley (1990), N: H.E. Reiley, REG: H.E. Reiley (2005). Fls 13/conical truss, saucer-shaped, c.75 × 95mm, with 5 wavy-edged lobes, strong red (51A) in bud, opening inside strong red (51A) at margins and with a light pink (50A), star-shaped centre; outside strong red (51A); faint dorsal spotting of brilliant greenish yellow (6A); slightly scented. Calyx c.1.5mm long, light brown. Lvs oblong, 145 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and deep green above, hairless. Shrub 1.5-1.8 × 1.2m in 10 years. Mid May.

(r) **'Reileyridge Sweet Sue'**

cv. Elepidote rhododendron: *fortunei* (s) × 'Caractacus'. G: H.E. Reiley (1974), N: H.E. Reiley (2004), REG: H.E. Reiley (2005). Fls 9/domed truss, saucer-shaped, c.65 × 95mm, with 7 wavy-edged lobes, very light purple (76B) in bud, opening inside very pale purple (76D) with a solid, dark purple (79A) blotch in the centre and discrete dorsal spotting of the same colour; outside very light purple (76B); no stamens; strongly scented. Calyx 1mm long, moderate purplish red (186B). Lvs oblong, 115-125 × 50mm, obtuse at apex, rounded at base, margins flat, semi-glossy and dark yellowish green (139A) above, hairless. Dense shrub, 3.0 × 4.5m in 30 years. Early. (Grown from seed purchased from F.W. Schumacher in 1965).

(r) **'Reileyridge Target'**

cv. Elepidote rhododendron: (*fortunei* × 'Caractacus') (s) × 'Caractacus'. G: H.E. Reiley (1974), N: H.E. Reiley (2004), REG: H.E. Reiley (2005). Fls 8/ball truss, broadly funnel-shaped, c.65 × 100mm, with 7 wavy-edged lobes, deep purplish pink (55A) in bud, opening inside pale purplish pink (56A), with light purplish pink (55C) edges and a solid, deep red (53A) centre; outside strong purplish pink (55B), with strong purplish red (54A) midribs; no stamens; strongly scented. Calyx 2mm long, moderate yellow-green (139C). Lvs oblong, 125 × 50mm, broadly acute at apex, oblique at base, semi-glossy and moderate olive green (137A) above, hairless. Dense shrub, 2.1 × 3.6m in 30 years. Early. (Grown from seed purchased from F.W. Schumacher, 1965).

(r) **'Richard Cree'**

cv. Elepidote rhododendron: 'Etta Burrows' (s) × (*yakushimanum* hybrid × *barbatum*). H: Mrs E.M. Watson (1995), G: Mrs E.M. Watson (2003), N: Mrs E.M. Watson, REG: Mrs E.M. Watson (2005). Fls

29/ball truss, open funnel-shaped, 50 × 70mm, with 5 smooth-edged lobes, strong red (51A) throughout. Lvs elliptic, 150 × 75mm, acute at apex, cuneate at base, margins wavy and downcurved, dull and moderate olive green (137A) above; petioles visibly hairy. Upright shrub, 1.5 × 0.9m in 10 years. Late April.

(v) **'Richard Marques'**

cv. Lepidote rhododendron: (*zoelleri* × *leucogigas*) (s) × *laetum*. H: R.A. Mitchell (2000), G: R.A. Mitchell (2002), N: R.A. Mitchell, REG: R.A. Mitchell (2005). Fls 4/flat truss, broadly funnel-shaped, with 5 very wavy-edged, large, deeply cut lobes, 108 × 80mm, inside strong purplish pink (62A) at margins, blending into light purplish pink (62C); outside light purplish pink (62C); both inside and outside colours blend very subtly with random blotches of strong purplish red (63B). Lvs elliptic, 165 × 50mm, acuminate at apex, cuneate at base, margins upcurved, midribs dark red (187C) and c.5mm wide at base, tapering to a fine point at apex; lvs change colour with age, from moderate yellow-green (137C) to dark red (187C); no scales visible to the naked eye. Shrub 1.2 × 0.75m in 5 years. Late January.

(r) **'Rio de Janeiro'**

cv. Elepidote rhododendron: ('Souvenir of Anthony Waterer' × 'Direktor E. Hjelm') (s) × 'Marrakesch'. H: H.J.A. Guyens (1994), G: H.J.A. Guyens (1998), N: H.J.A. Guyens (1998), I: H.J.A. Guyens, REG: H.J.A. Guyens (2005). Fls 14-18/hemispherical truss, funnel-shaped, 70 × 80mm, with 5-7 slightly wavy-edged lobes, inside creamy salmon pink with darker salmon pink edges, outside salmon pink; a dark brown to orange blotch at the base of the corolla, the three dorsal lobes also with brown and olive green markings on a creamy yellow ground. Calyx 20-30mm long, creamy salmon pink inside and out, with brown-orange markings. Lvs lanceolate, 100-160 × 25-45mm, margins wavy and upcurved, broadly acute at base and apex, matt mid green above, hairless. Shrub 0.8 × 0.7m in 7 years. Early to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0052451)).

(a) **'Rob Bruce Stewart'**

cv. Evergreen azalea: 'Desiree' (Shammarello) × semi-double hybrid of unknown parentage (no.CB-1). H: Robert Stewart (1990), G: Robert Stewart (1993), N: Robert Stewart (2005), I: Robert Stewart (2005), REG: Robert Stewart (2005). Fls broadly funnel-shaped, 65 × 40mm, with 5 very wavy-edged lobes, buds white with markings close to light purple (75B), opening white striped and speckled with strong reddish purple (72C) and with some selfs near light purple (75B); blotch of light yellow-green (near 142C) and some some strong purplish pink (66B). Calyx 20mm long, mid green. Lvs elliptic, 65 × 25mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and moderate olive green (146A) above. Shrub 0.9 × 1.1m in 14 years. Midseason. (Etymology: named for the son of the hybridizer).

(a) 'Rocky River Red'

cv. Deciduous azalea: *cumberlandense* (s) × *arborescens*. H: J.E. Schild (1986), G: J.E. Schild (1993), N: J.E. Schild (1996), I: Schild Azalea Gardens and Nursery (2006), REG: J.E. Schild (2005). Fls 20-24/ball truss, tubular funnel-shaped, 45 × 50mm, with 5 flat-edged lobes, strong reddish orange (40C) in bud, opening inside vivid reddish orange (N30A) with a vivid reddish orange (N30B) blotch on the dorsal lobe; outside vivid reddish orange (40B) with many glandular hairs along the tube; filaments vivid reddish orange (41A); style vivid red (44A); stigma deep purplish red (71A); slightly scented. Calyx c.3mm long, strong yellow-green (145A). Lvs elliptic, 85 × 35mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (137A) above, moderate yellowish green (138A) below, hairs on leaf margins and on midveins below (hairs white at first, aging to yellowish white). Dense shrub, 0.9 × 0.9m in 10 years. June. (Published in *The Azalean* 27(2):32 (2005)).

(a) 'Ronc de Gira'

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 2-4/truss, 73 × 48mm, with 5 flat to rarely wavy-edged lobes, buds strong red (51A), opening deep pink (51B) throughout; three lobes with deep red (53A) markings, shaded with strong red (53B); stamens 5, long exerted from the corolla, anthers deep red (53A) shaded greyed purple (N186A), filaments moderate pink (49B) shaded deep pink (48A); pistil much shorter than the stamens but still exerted from the corolla, style strong red (39A) shaded deep yellowish pink (39B), stigma strong red (47B). Lvs elliptic, 32 × 20mm, margins flat, glossy above, obtuse at base, acute at apex, hairy, strong yellow-green (144A) above and moderate yellow-green (147C) below when mature; petioles 4mm long. Shrub 1.0 × 1.5m in 50 years. April to May. (Etymology: named for a place near Cannero Riviera, the latter being the location of the Carmine nursery).

(a) 'Ronc de Masok'

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 2-4/truss, 70 × 49mm, with 5 flat or rarely wavy-edged lobes, deep pink (50B) in bud and throughout on opening; three lobes with deep red (53A) markings, shaded strong red (53C); stamens 5, exerted from the corolla, anthers dark red (187A), shaded moderate purplish red (186A), filaments strong pink (52D), shaded vivid red (52A). Lvs lanceolate, 49 × 13mm, margins very wavy, glossy above, acute at apex and base, hairy, moderate olive green (146A) above and moderate yellow-green (146B) below when mature; petioles 4mm long. Shrub 1.5 × 1.0m in 50 years. April to May. (Etymology: named for a place near Cannero Riviera, the latter being the location of the Carmine nursery).

(t) 'Ronny'

cv. Lepidote rhododendron: 'Saint Merry' (s) × 'Azurika'. H: Holger Hachmann (1992), G: Holger Hachmann (1998), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls 3-5/individual truss, the trusses gathered at shoot tips into a dense ball inflorescence, broadly funnel-shaped, c.20 × 26-32mm, with 5 wavy-edged lobes, brilliant violet (90D) inside and out; unmarked; stamens 6-10, filaments strong/brilliant violet (86C/D), anthers dark red (187B); style vivid reddish purple (74A/B), stigma dark red (187A). Calyx 1-2mm long, light yellow-green (144D). Lvs elliptic, 18 × 5-6mm, margins flat, glossy above, cuneate at base, acute at apex, with scales above and below. Shrub c.25 × c.45cm in 10 years, with a dense, low/rounded habit. Late April to early May. (Nomenclatural standard: colour print provided by the registrant (WSY0070161). Published in Hachmann cat. 2004-2008 (2004), p.132).

(a) 'Rosa Confetto Cavadini'

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 3-9/truss, 40 × 29mm, with 5 flat, rarely wavy or very wavy-edged lobes, buds strong purplish pink (68B), opening strong purplish pink (62A) inside and out; three lobes with strong red (53C) markings; stamens 5, exerted from the corolla, anthers moderate reddish brown (176B), filaments deep purplish pink (64D); pistil just exceeding stamens, style deep purplish pink (58D), stigma orange-red (N34C). Lvs oblanceolate, 29 × 12mm, margins flat, glossy above, obtuse at base, acute at apex, hairy, moderate olive green (146A) above and moderate yellow-green (146D) below when mature; petioles 5-7mm long. Shrub 2.0 × 2.5m in 60 years (regularly pruned). May. (Etymology: flower colour resembles that of a sugared almond).

(a) 'Rosa Doppio Cavadini'

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls hose-in-hose, 2-5/hemispherical truss, 46 × 30mm, with 5+5 flat, rarely wavy-edged lobes, deep purplish pink (55A) in bud and throughout on opening; three lobes with strong red (53C) markings; stamens 6, exerted from the corolla, anthers moderate reddish brown (176B), filaments strong purplish pink (55B); pistil not or distinctly exceeding the stamens, style deep purplish pink (55A), stigma strong red (53D). Lvs elliptic, 34 × 17mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (146C) below when mature; petioles 3-4mm long. Shrub 1.6 × 4.0m in 60 years (regularly pruned). May. (Etymology: the name suggests the flower shape and colour).

(a) 'Rosa Intenso Cavadini'

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini

(1960), REG: Prof. E. Accati (2005). Fls single, 2-5/hemispherical truss, 31 × 26mm, with 5 flat-edged lobes, strong purplish red (61B) in bud and throughout on opening; unmarked; stamens 0-5, exerted from the corolla, anthers light reddish brown (177B), shaded moderate reddish brown (177A), filaments strong purplish red (61B); pistil not or just exceeding stamens, style strong purplish red (61B), stigma strong purplish red (71B). Lvs oblanceolate, 17 × 8mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (147A) above and moderate yellow-green (147C) below when mature; petioles 4-5mm long. Shrub 0.8 × 2.0m in 60 years (regularly pruned). April to May. (Etymology: the name suggests the colour of the flower).

(a) **'Rosa Pastello Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-8/hemispherical truss, 48 × 28mm, with 5 flat to very wavy-edged lobes, buds deep pink (52B), opening deep pink (52C) inside and out; three lobes with dark reddish orange (178B) markings; stamens 5, exerted from the corolla, anthers brownish orange (165B), filaments deep pink (52C); pistil exceeding stamens, style deep pink (52C), stigma vivid reddish orange (33B). Lvs lanceolate, 33 × 13mm, margins flat, glossy above, acute at base, obtuse at apex, hairy, moderate olive green (146A) above and strong yellow-green (144B) below when mature; petioles 7-8mm long. Shrub 1.7 × 2.0m in 60 years (regularly pruned). April. (Etymology: the name suggests the colour of the flower).

(a) **'Rosa Salmone Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls hose-in-hose, single or double, 2-5/truss, 32 × 29mm, with 5 (+5) flat or slightly wavy-edged lobes, buds vivid reddish orange (43B), opening deep yellowish pink (43C) inside and out; three lobes with strong red (53D) markings; stamens 0-5, exerted from the corolla, anthers light reddish brown (177B), filaments strong yellowish pink (43D); pistil exceeding stamens, style deep yellowish pink (43C), stigma vivid reddish orange (43B). Lvs elliptic, 22 × 13mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (147C) below when mature; petioles 2-3mm long. Shrub 1.6 × 1.5m in 60 years (regularly pruned). April to May. (Etymology: the name suggests the colour of the flower).

(a) **'Rosbevan Star'**

cv. Evergreen azalea: 'Diana Trask' (s) × 'Windmill'. H: D.J. Dosser (1990), G: D.J. Dosser (1995), N: D.J. Dosser (2004), REG: D.J. Dosser (2005). Fls semi-double, 3/truss, broadly funnel-shaped, 40 × 75mm, corolla with 5 wavy-edged lobes and with several shorter petaloids in the centre of the flower, strong

red (53D) inside and out, with a few, deeper, dorsal spots. Lvs elliptic, 40 × 15mm, cuneate at base, obtuse at apex. Shrub 0.6 × 0.6m in 14 years. September to October (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0070188)).

(r) **'Royal Fusilier'**

cv. Elepidote rhododendron: 'Noyo Chief' (s) × *elliottii*. H: H. O'Rourke (1982), G: H. O'Rourke (1998), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 25/flat-topped truss, funnel-campanulate, 45 × 50mm, with 5 wavy-edged lobes, vivid red (46B) in bud and both inside and out on opening, with a few dark spots on the dorsal lobes. Calyx 2mm long, red. Lvs elliptic, 130 × 45mm, margins decurved, matt green above, cuneate at base, acute at apex, with fawn hairs. Shrub 1.5 × 1.2m in 17 years. September to October (Australia).

(r) **'Royal Treasure'**

cv. Elepidote rhododendron: 'Kubla Khan' (s) × 'Ayers Rock'. H: H. O'Rourke (1985), G: H. O'Rourke (1996), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 10/loose truss, tubular funnel-shaped, 50 × 90mm, with 5 wavy-edged lobes, pink and yellow in bud, opening inside yellow in the tube, the lobes edged pink; outside yellow, edged pink; with a small red blotch in the centre of the flower. Calyx 15mm long, pink and yellow. Lvs oblong, 70 × 30mm, margins wavy, matt green, cuneate at base, acute at apex, hairless. Shrub 1.2 × 1.2m in 12 years. October to November (Australia).

(a) **'Sabioncella'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls hose-in-hose, 1-4/truss, 45 × 33mm, with 5+5 wavy to rarely very wavy-edged lobes, red-purple (N66A) in bud and throughout on opening; three lobes with deep red (60A) markings shaded strong purplish red (60B); stamens 5-7, exerted from the corolla, anthers dark red (59A), filaments red-purple (N57D), with red-purple (N57A) near the base; pistil held within the corolla, style strong red (53C), stigma red-purple (N79B). Lvs lanceolate, 46 × 21mm, margins flat, glossy above, obtuse at base, acute at apex, hairy, moderate olive green (147A) above and light yellow-green (145C) below when mature; petioles 4mm long. Shrub 0.9 × 0.6m in 50 years. April to May. (Etymology: named for a place near Cannero Riviera, the latter being where the Carmine nursery is situated).

(r) **'Sakura-kage'**

cv. Elepidote rhododendron: (R × *nikomontanum*) (s) × *degronianum* (with double white fls). H: H. Maehara (1993), G: H. Maehara (1998), S: H. Maehara (2002), N: H. Maehara (2004), REG: H. Maehara (2005). Fls double, with no normal stamens, 7-9/hemispherical truss, open funnel-shaped, 75 × 90mm, with 10-13 very wavy-edged lobes, pink in bud, opening white

throughout inside and out, unmarked. Calyx 2mm long, light brown. Lvs elliptic-oblong, 85 × 28mm, margins flat, glossy above, rounded at base, acute at apex, hairless. Very compact shrub, 27 × 45cm in 12 years. Late April. (Nomenclatural standard: colour slide provided by the registrant (WSY0070244)).

(a) **'Salmellia'**

cv. Evergreen azalea: parentage unknown. H: F. Furman (c.1975), G: F. Furman (c.1980), N: F. Furman, REG: F. Furman (2005). Fls 1-2/truss, saucer-shaped and irregularly double, c.40 × 55mm, with 5-10 wavy lobes, deep pink (50B) at edges, shading to deep pink (48C) at centre. Calyx 3mm long, green. Lvs oblanceolate, spring lvs 25 × 13mm, summer lvs c.13 × 6mm, mid green. Shrub 0.4 × 0.6m in 10 years. Midseason.

(r) **'Sammy'**

cv. Elepidote rhododendron: parentage unknown. H: H. O'Rourke (1986), G: R. & E. Tavner (1989), N: R. & E. Tavner (2005), REG: R. & E. Tavner (2005). Fls 18/flat-topped truss, campanulate, 70 × 62mm, with 5 flat-edged lobes, pale yellow-green (4D) in bud both inside and out when open, unmarked. Calyx 37mm long, light yellowish pink (36C). Lvs elliptic, 140 × 50mm, margins flat, matt green above, obtuse at base, acute at apex, hairless. Shrub 1.5 × 1.5m in 19 years. Late September (Tasmania).

(r) **'Samtkrone'**

cv. Elepidote rhododendron: 'Tarantella' (s) × Kilimanjaro Group. H: H. Hachmann (1990), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 19-21/fairly compact truss, broadly funnel-campanulate, 45-50 × 70-75(-80)mm, with 5 wavy-edged lobes, buds dark red (187A/B), opening moderate red (47A) inside and out, narrowly edged dark red/ deep purplish red (59A/B); dorsal lobe markings of dark red (59A); stamens 10, filaments strong purplish red/strong purplish pink (63B/C), anthers moderate reddish brown (166B); style light purplish pink (62C), stigma dark red (187A/B); ovary light green, silvery hairy. Calyx 1mm long, reddish. Lvs narrowly elliptic, 90-130 × 35-55mm, margins slightly wavy, matt dark green above, edges and petioles yellowish green, rounded to oblique at base, apiculate at apex, hairless. Shrub c.1.5 × c.1.8m in 10 years, with a neat, dense habit. Late May to early June. (Fls long-lasting, not fading in the sun; later flowering than most reds. Nomenclatural standard: colour print provided by the registrant (WSY0070148). Published in Hachmann cat. 2004-2008 (2004), p.74).

(a) **'San Bernardo'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 1-2/truss, 60 × 40mm, with 5 wavy to rarely very wavy-edged lobes, deep purplish pink (66C) in bud and throughout on opening; three lobes with deep red (60A) markings; stamens 4-5, exerted from

the corolla, anthers dark red (187A), shaded very pale purple (92D), filaments pale purplish pink (62D), shaded strong purplish red (63A); pistil distinctly exceeding the corolla, style deep pink (52C), shaded yellow-green (N144C), stigma deep pink (48A). Lvs elliptic, 35 × 16mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (147B) below when mature; petioles 4mm long. Shrub 1.3 × 2.1m in 50 years. April to May. (Etymology: San Bernardo and San Rocco are the patron saints of Cannero Riviera, where this cultivar originated).

(r) **'Sanderling'**

cv. Lepidote rhododendron: *dauricum* 'Hokkaido' (s) × *leucaspis*. H: P.A. Cox (1970s), G: Det Norske Arboret (c.1990), N: P.M. Jørgensen (2005), I: Glendoick Gardens (c.1990), REG: P.M. Jørgensen (2005). Fls 5/lax, open truss, flat saucer-shaped, 20 × 15mm, with 5 flat-edged lobes, white throughout, unmarked; style slightly curved, stigma pink; anthers black. Calyx up to 5mm long, green. Lvs narrowly elliptic, 35-50 × 25mm, margins flat, glossy above, tapered at base, mucronate at apex, with brownish black scales; scented when crushed. Shrub 1.0 × 0.5m in 10 years. (Nomenclatural standard: colour slide provided by the registrant (WSY0060829). Originally introduced under parentage only).

(a) **'Sandra's Harlequin'**

cv. Evergreen azalea: unknown (s) × 'Blushing Angel'. H: S.F. McDonald (1984), G: S.F. McDonald (1988), N: S.F. McDonald, REG: S.F. McDonald (2005). Fls 2-3/truss, open funnel-shaped, 43 × 70mm, with 5 wavy-edged lobes, white, with stripes, sectors and flecks of deep yellowish pink (39B) in bud, opening white, with stripes, sectors and flecks of deep yellowish pink (39B) inside and out; no blotch. Calyx 10mm long, strong yellow-green (144C). Lvs elliptic, 50 × 20mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (147A) above, hairy above and below, hairs tan-coloured. Shrub 1.5 × 1.2m in 20 years. Mid April to early May. (Seed parent was an unknown, evergreen azalea found as a rogue at de Wilde Greenhouses).

(a) **'San Rocco'**

cv. Evergreen azalea: parentage unknown. R: M. Carmine (1940), N: Mrs O. Carmine (2004), I: M. Carmine (1960), REG: Prof. E. Accati (2005). Fls single, 2/truss, 70 × 46mm, with 5 wavy-edged lobes, deep pink (52B) in bud, and throughout on opening; three lobes with strong red (53B) markings, shaded with deep red (53A); stamens 5-6, not exerted, anthers strong red (51A), shaded red-purple (N79B), filaments strong purplish red (55B) shaded with white; pistil distinctly exceeding the corolla, style light yellow-green (2C), shaded deep pink (50B), stigma dark greenish yellow (152C). Lvs lanceolate, 55 × 21mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (146A) above and moderate yellow-green (146B) below when mature;

petioles 4mm long. Shrub 1.8 × 2.0m in 50 years. April to May. (Etymology: San Rocco and San Bernardo are the patron saints of Cannero Riviera, where this cultivar originated).

(r) **'Santorina'**

cv. *Elepidote rhododendron*: ('Hotei' × ('Goldrausch' × *wardii* hybrid)) (s) × Fred Wynniatt Group. H: H. Hachmann (1990), G: H. Hachmann (1996), N: H. Hachmann (2003), I: H. Hachmann (2004), REG: G. Stück (2005). Fls 8-10/fairly compact truss, broadly campanulate, 45-50 × 75-85mm, with (5-)7 wavy-edged lobes, inside pale greenish yellow (13D), the centre and throat brilliant yellow (13C), the three dorsal lobes brilliant yellow (13B), lobes edged light purplish pink (62C); outside light purplish pink (62C) at edges, shading basally to a mixture of yellow and purple and to brilliant yellow (13C) at base; inconspicuous markings and a small basal spot of deep red (185A) in the dorsal throat; stamens 14, filaments pale greenish yellow (13D), anthers moderate reddish brown (166B); style brilliant yellow-green (150C), stigma brilliant yellow-green (150B); ovary moderate yellow-green (138C), with silvery hairs. Calyx hose-in-hose, 10-15mm long, mixed yellow and purple. Lvs ovate, 75-85 × 35-45mm, margins slightly wavy, matt light green above, rounded or occasionally oblique at base, minutely apiculate at apex, hairless. Broadly compact, upright shrub c.1.2 × c.1.4m in 10 years. Early to mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070149). Published in Hachmann cat. 2004-2008 (2004), p.74).

(r) **'Saveta'**

cv. *Elepidote rhododendron*: 'Girard Joshua' (s) × *yakushimanum*, Exbury form. H: J.D. Hooper (c.1992), G: J.D. Hooper (c.1997), N: J.D. Hooper (2005), REG: J.D. Hooper (2005). Fls 14-16/domed truss, funnel-campanulate, 50 × 60mm, with 5 wavy-edged lobes, strong purplish red (61B) in bud, opening inside light purple (76A), narrowly edged moderate purplish red (64A), with sparse, strong orange-yellow (163B) dorsal spotting; outside strong purplish red (64C) with slightly darker midribs; colour fading slightly with age. Calyx c.3mm long, cream. Lvs elliptic, 140 × 45mm, broadly acute at apex, rounded at base, margins slightly downcurved, semi-glossy and dark olive green above, young lvs covered with white tomentum, hairs thinly covering underside and petioles of mature lvs (hairs almost white when young, aging to orange tan). Shrub 0.9 × 1.5m in c.13 years. Midseason.

(r) **'Scotian Bells'**

cv. *Lepidote rhododendron*: *mucronulatum* var. *taquetii* (s) × *camphylogynum* Charopaum Group. H: J.K. Weagle (1987), G: J.K. Weagle (1990), N: J.K. Weagle, I: Glendoick Gardens, REG: J.K. Weagle (2005). Fls in lax trusses, pendent, campanulate, (2-)3/bud and with 2-3 buds/terminal shoot, 26 × 30mm, with slightly wavy-edged lobes, deep purplish red (71A) in bud, opening inside light purple (81C), outside vivid

purple (80A); unmarked; stigma 23mm long, curved sharply downwards. Calyx 4mm long. Lvs elliptic to orbicular, 27 × 17mm, broadly acute at apex, cuneate at base, glossy and between moderate olive green (147A) and dark yellowish green (139A) above, with reddish scales, many spaced 2-3 times their width apart; petioles 5.5mm long. Shrub 0.3 × 0.5m in 12 years, very compact even in shade. May be deciduous in very cold, snowless winters without impairment. Mid to late May.

(a) **'Scotian Breeze'**

cv. Evergreen azalea: *keiusianum* (RSF no.81/11) (s) × 'Lady Louise'. H: J.K. Weagle (c.1985-1986), G: J.K. Weagle, N: J.K. Weagle, I: Rhodoland Nursery, REG: J.K. Weagle (2005). Fls 2/bud, open funnel-shaped, with a flat face, 30 × 35mm, with 5 slightly wavy-edged lobes, light purplish pink (55C) in bud, opening inside and out with a 5mm wide margin of strong purplish pink (68B) that fades abruptly to a white centre; occasional fls are solid strong purplish pink (68B); bicolors have faint (68B) spotting; solid colours have darker spotting. Calyx c.1mm long, covered with silky white hairs. Summer lvs elliptic/obovate, 20 × 15mm, broadly acute/obtusate at apex, cuneate at base, moderate yellow-green (146B-C); winter lvs obovate/elliptic, 15 × 6mm, broadly acute/obtusate at apex, cuneate at base, moderate olive green (146A) above; summer lvs very slightly hairy above, winter lvs shiny, with flat, very short, reddish hairs. Very compact shrub, 0.8 × 1.2m in c.20 years. Late June.

(a) **'Senior Prom'**

cv. Deciduous azalea: *arborescens* (s) × *cumberlandense*. H: J.E. Schild (1986), G: J.E. Schild (1994), N: J.E. Schild (2003), I: Schild Azalea Gardens and Nursery (2006), REG: J.E. Schild (2005). Fls 6-8/domed truss, tubular funnel-shaped, 55 × 55mm, with 5 flat-edged lobes, buds strong purplish red (63A) with yellowish white ribs, dorsal lobe opening white with strong purplish red (63B) margins and a vivid yellow (15A) blotch in the centre; other lobes strong purplish red (63B) inside; outside strong purplish pink (62A), randomly flushed with vivid purplish pink (N66B); style moderate purplish pink (62B); filaments light purplish pink (62C); stigma greyish reddish purple (N77C); moderately scented. Calyx 5mm long, brilliant yellow-green (142B). Lvs obovate, 65 × 35mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate yellow-green (138C) above, moderate yellow-green (138B) below, margins hairy and with sparse hairs on the blade above, many hairs on midvein below (hairs white, aging to green-white). Shrub 1.2 × 1.2m in 10 years. Early June. (Published in *The Azalean* 27(2):31 (2005)).

(r) **'Sheila Gay'**

cv. *Elepidote rhododendron*: (*maximum* × 'Mary Garrison') (s) × ('Caroline' (Gable) × ('King of Shrubs' × 'Crest')). H: M.E. Byrkit (1993), G: J.D. Hooper (2000), N: J.D. Hooper (2005), REG: J.D. Hooper (2005). Fls 10-15/flat to slightly lax truss,

broadly funnel-shaped, 60 × 95mm, with 5(-6) wavy-edged lobes, strong purplish red (60D) in bud, opening inside very light purple (75C), the throat flushed pale yellow (11D) with a deep to moderate red (185B) blotch at the base; outside light purplish pink (62C), the midribs slightly darker; dorsal lobe has a narrow, dark reddish orange (175C) flare with strong orange-yellow (163B) speckles. Calyx c.3mm long, apple green. Pedicels wine red. Lvs elliptic, 165 × 45mm, broadly acute at apex, cuneate at base, margins flat, dull and dark olive green above, hairless. Shrub 1.2 × 1.7m in c.12 years, with decumbent lower branches. Late (early July).

(v) **'Sherla'**

cv. Lepidote rhododendron: ((*leucogigas* × *laetum*) × *javanicum*) (s) × *konori*. H: R.A. Mitchell (1999), G: R.A. Mitchell (2005), N: R.A. Mitchell (2005), REG: R.A. Mitchell (2005). Fls 12/truss, the individual flowers disposed horizontally, tubular funnel-shaped, 65 × 75mm, with 7 wavy-edged lobes, inside light purplish pink (65B) at edges, blending to pale purplish pink (65C) and then to light yellow-green (2C) at centre; outside pale purplish pink (65C), blending to pale greenish yellow (2D) at base; unmarked. Lvs obovate, 125 × 65mm, broadly acute at apex, cuneate at base, margins flat, dull and moderate olive green (137B) above, with no scales visible to the naked eye. Shrub 0.9 × 0.6m in 4.5 years. February. (The *konori* used is one known locally as Jakes's form).

(r) **'Shin-shun'**

cv. Elepidote rhododendron: *brachycarpum* (s) × *moulmainense* (as *leopodum*). H: K. Kasai (1995), S: K. Kasai (2003), G: K. Kasai (2003), N: K. Kasai (2004), REG: K. Kasai (2005). Fls 16-18/hemispherical truss, open funnel-shaped, 45 × 65mm, with 5-6 wavy-edged lobes, deep red (53A) in bud, opening inside strong purplish red (60D), outside strong purplish red (61B), with moderate purplish red (59C) spots on the dorsal lobe. Calyx 2mm long, moderate purplish red (59C). Lvs oblong-elliptic, 110 × 70mm, margins distinctly recurved at first, later they may become upcurved, matt green above, rounded at base, obtuse at apex, hairless above. Compact shrub 20 × 45cm in 10 years. Early June.

(a) **'Shiroi-shōjo'**

cv. Evergreen azalea: (white-flowered *kinsianum* × (white-flowered *kaempferi* f. *latisepalum* × *yedoense* var. *poukhanense*)) (s) × (*schlippenbachii* × (white-flowered *kaempferi* f. *latisepalum* × white-flowered *yedoense* var. *poukhanense*)). H: T. Saitō (1998), G: T. Saitō (2004), S: T. Saitō (2004), N: T. Saitō (2004), REG: T. Saitō (2005). Fls hose-in-hose, 2-3/open truss, funnel-shaped, 32 × 20mm, with 5+5 wavy-edged lobes, white in bud and throughout on opening, unmarked; stamens 5, filaments white; style white. No normal calyx present. Lvs elliptic, 37 × 15mm, margins flat, matt green above, obtuse at apex, cuneate at base, with white hairs. Shrub 25 × 43cm in 7 years. Mid May. (Nomenclatural standard: colour slide provided by the registrant (WSY0070239)).

(r) **'Silent Angel'**

cv. Elepidote rhododendron: *wardii* (s) × 'Purple Splendour'. H: Ben Staab (1995), G: Billie Staab (2005), N: J.L. Huisingh, REG: J.L. Huisingh (2005). Fls 8/flat truss, broadly funnel-shaped, 50 × 65mm, with 5 wavy-edged lobes, deep purplish pink (54B) in bud, opening inside pale purplish pink (55D) with a pale yellow (11C) throat, the yellow colour extending upwards along the dorsal lobe; outside strong purplish pink (55B); no spotting. Calyx 2mm long, brownish orange (166C). Lvs elliptic, 90 × 40mm, obtuse at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137B) above, hairless. Shrub 1.2 × 1.2m in 10 years. Mid May.

(r) **'Siltcoos Sunrise'**

cv. Elepidote rhododendron: 'Paprika Spiced' (s) × 'Jezebel'. H: E. Cockeram (1995), G: E. Cockeram (2000), N: E. Cockeram (2000), REG: E. Cockeram (2005). Fls 10-13/domed truss, open funnel-shaped, 45 × 50mm, with 5 wavy-edged lobes, strong yellowish pink (37A) in bud, opening inside brilliant yellow (8A) in the throat, fading to brilliant yellow (11A); outside brilliant yellow (11A); irregularly shaped, vivid reddish orange (33B) markings inside all corollas - varying from single or multiple flares to entire centres. Calyx 30mm long, brilliant yellow (11A), with vivid reddish orange (33B) spots. Lvs lanceolate, 100 × 30mm, broadly acute at apex, cuneate at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.6m in 9 years. Early April.

(a) **'Silvio della Valle di Casanova'**

cv. Evergreen azalea: parentage unknown. R: Gardeners of the Villa San Remigio (pre 1920), N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single to semi-double, 1-3/truss, 80 × 48mm, with 5 flat-edged lobes, red-purple (N57B) in bud and throughout on opening, three lobes with deep purplish red (59B) markings; stamens 10, not exerted from the corolla, anthers purple (N77A), filaments strong red (53B); pistil distinctly exceeding stamens and corolla, style pale yellow-green (157A), stigma strong red (53B). Lvs lanceolate, 48 × 20mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (147C) below when mature; petioles 8mm long. Shrub 3.0 × 4.0m in c.100 years. May. (Etymology: Silvio della Valle di Casanova, together with his wife Sofia Browne, built the garden of Villa San Remigio in the 19th century).

(a) **'Sky Hill Pose'**

cv. Evergreen azalea: probably 'Edna Bee' (s) × 'Hahn's Red'. H: F. Furman (c.1975), G: F. Furman (c.1980), N: F. Furman, REG: F. Furman (2005). Fls 1-2/truss, saucer-shaped, single, semi-double or fully double, c.50 × 75mm, with 5-10 wavy-edged lobes, vivid purplish red (57C) at edges, shading to strong purplish red (58C) at centre, with a moderate purplish red (58A) dorsal flare. Calyx 5mm long, green. Lvs oblanceolate, spring leaves 40 × 20mm, summer

leaves c.30 × 13mm, mid green, darkening in winter. Shrub 0.5 × 0.8m in 10 years. Midseason.

(r) **'So Chic'**

cv. Elepidote rhododendron: 'College Pink' (s) × Dido Group. H: H. O'Rourke (1985), G: H. O'Rourke (1987), N: H. O'Rourke (2005), REG: H. O'Rourke (2005). Fls 16/rounded truss, tubular funnel-shaped, 65 × 85mm, with 5 wavy-edged lobes, light purplish pink (55C) in bud and throughout on opening. Calyx green. Lvs ovate, 130 × 50mm, margins wavy, matt green above, cuneate at base, acute at apex, hairless. Shrub 1.2 × 1.5m in 18 years. October (Australia).

(a) **'Sofia Browne'**

cv. Evergreen azalea: parentage unknown. R: Gardeners of the Villa San Remigio (pre 1920), N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single, 1-3/truss, 58 × 33mm, with 5 flat-edged lobes, vivid red (52A) in bud, opening deep pink (52B) inside and out; strong red (53C) markings on one lobe (rarely on three); stamens 7-9, exerted from the corolla, anthers deep red (53A), filaments deep pink (52B); pistil not or sometimes exceeding stamens, style deep pink (52B), stigma brownish orange (165B). Lvs lanceolate 38 × 14mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (146C) below when mature; petioles 5mm long. Shrub 2.0 × 5.0m in c.100 years. May. (Etymology: Sofia Browne, with her husband Silvio della Valle di Casanova, built the garden at Villa San Remigio in the 19th century).

(r) **'Something Royal'**

cv. Elepidote rhododendron: 'Sir Frederick Moore' (s) × un-named seedling from 'Australian Sunset'. H: H. O'Rourke (1988), G: H. O'Rourke (1998), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 14/loose truss, tubular-campanulate, 70 × 100mm, with 6 wavy-edged lobes, strong purplish pink (54B) in bud, opening inside salmon pink, outside deep purplish pink (54B). Calyx 2mm long, pink. Lvs lanceolate, 120 × 45mm, margins decurved, obtuse at base, acute at apex. Shrub 1.5 × 1.5m in 16 years. October to November (Australia).

(r) **'Southern Alps'**

cv. Elepidote rhododendron: parentage unknown, but evidently a hybrid derived from sect. *Fortunea*. G: W.R. & M.N. Philipson, N: W.R. & M.N. Philipson, REG: M.N. Philipson (2005). Fls 7-9/globular truss, open campanulate, 70 × c.110mm, with 7 slightly wavy-edged lobes, pale pink in bud, opening white inside and out with a very small, soft pink or pale yellow blotch at the base of the dorsal throat; ovary and style glandular. Pedicels glandular. Calyx irregularly lobed, to c.3mm long. Lvs oblanceolate to oblong, (110-)170-190(-200) × (35-)40(-45)mm, margins upcurved, broadly acute at apex, hairless. Shrub with a fastigiate habit with many branches, 3.5m tall in 20 years. September to October (New

Zealand). (Found by the Philipsons as a very small seedling growing on a tree stump in the former garden of Edgar Stead at Ilam, Christchurch, New Zealand in c.1960. Nomenclatural standard: colour print provided by the registrant (WSY0070519). Etymology: named after the snow-covered mountain range in the South Island of New Zealand).

(r) **'Spot of Gold'**

cv. Elepidote rhododendron: 'Molly Smith' (s) × 'Mrs J.G. Millais'. H: C. Nuys (1990), G: C. Nuys (1993), N: C. Nuys (2005), REG: C. Nuys (2005). Fls 16/ball truss, broadly funnel-shaped, c.70 × 90mm, with 5 flat-edged lobes, pink in bud, opening white, tinged pale pink throughout, with a large, vivid yellow (17B), speckled, dorsal blotch; as the flowers age they become pure white and the blotch becomes darker. Calyx c.10mm long, yellowish green. Lvs elliptic, 125 × 40mm, acute at apex, cuneate at base, margins flat, moderate olive green (147A) above, hairless. Shrub 1.8 × 1.8m in 15 years. Mid May.

(r) **'Star Spangled'**

cv. Elepidote rhododendron: 'Lem's Goal' (s) × 'Skipper Hal'. H: M. Stockman (1994), G: M. & H. Stockman (2001), N: H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 6-7/lax truss, funnel-campanulate, 50 × 75mm, with 5 very wavy-edged lobes, strong purplish red (61B) in bud, opening inside pale purplish pink (62D), tinged deep purplish pink (66C) randomly and along margins, with a large, deep red (53A), freckled blotch on the dorsal lobe and on half of each of the adjacent lobes; outside strong purplish red (61B). Lvs elliptic, 105 × 40mm, acute at apex, cuneate at base, margins wavy, dull and moderate olive green (137B) above, moderate yellow-green (146D) below, hairless. Shrub 0.75 × 0.9m in 10 years. Mid May.

(a) **'Stefania Cavadini'**

cv. Evergreen azalea: parentage unknown. R: L. Cavadini (1930), N: G. Cavadini (2004), I: L. Cavadini (1960), REG: Prof. E. Accati (2005). Fls single, 2-7/hemispherical truss, 48 × 24mm, with 5 flat-edged lobes, strong purplish red (58C) in bud and on opening, both inside and out; three lobes with deep purplish red (61A) markings; stamens 5-6, exerted from the corolla, anthers light olive grey (197A), filaments deep purplish pink (61D); pistil exceeding stamens, style deep purplish pink (58D), stigma strong red (41B). Lvs lanceolate, 27 × 12mm, margins flat, glossy above, obtuse at base and apex, hairy, moderate olive green (137B) above and moderate yellow-green (146C) below when mature; petioles 4-5mm long. Shrub 1.8 × 3.0m in 60 years (regularly pruned). April. (Etymology: named for the nominant's daughter-in-law).

(r) **'Stockman's Black Eye'**

cv. Elepidote rhododendron: 'Van Nes Sensation' (s) × 'Kubla Khan'. H: M. Stockman (1992), G: M. & H. Stockman (2000), N: H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman

(2005). Fls 8/conical truss, broadly funnel-shaped, 65 × 90mm, with (5-6-)7 wavy-edged lobes, strong purplish pink (62A) in bud, opening inside moderate purplish pink (62B), shading to light purplish pink (62C), with a prominent basal blotch darker than deep red (53A), surrounded by a jagged, strong purplish red (63B) blotch; outside strong purplish pink (62A), with some colour showing through from the blotch. Calyx lobes c.25-30mm long, reflexed, coloured as the corolla. Lvs elliptic, 120 × 45mm, broadly acute at apex, rounded at base, margins wavy, dull and moderate olive green (147A) above, moderate yellow-green (147C) below, hairless. Shrub 0.75 × 0.9m in 12 years. Mid May.

(r) **'Stockman's Firecracker'**

cv. Elepidote rhododendron: 'Blimey Limey' (s) × 'Gallacci's Gold'. H: M. Stockman (1992), G: M. & H. Stockman (2000), N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 9-11/flattened ball truss, funnel-campanulate, 55 × 70mm, with 6 wavy-edged lobes, vivid red (45D) in bud, opening inside deep pink (48A) with a slight yellowish tinge, shading to solid deep pink (51B) at edges, and with two narrow lines of strong red (45D) spotting on dorsal lobe; outside deep pink (50B). Calyx with dorsal lobes c.15-20mm long, ventral lobes c.10-15mm long, dorsal deep pink (48B), ventral moderate yellowish pink (31D), all lobes somewhat reflexed. Lvs elliptic, 165 × 45mm, acute at apex, cuneate at base, margins flat and downcurved, glossy and moderate yellow-green (137C) above, moderate yellow-green (143D) below, hairless. Shrub with a sprawling habit, 0.75 × 1.25m in 12 years. Mid May.

(r) **'Stockman's Flirt'**

cv. Elepidote rhododendron: 'Blimey Limey' (s) × 'Christie Ess'. H: M. Stockman (1993), G: M. & H. Stockman (2002), N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 7-8/loose ball truss, open funnel-shaped, 45 × 65mm, with 6 very wavy-edged lobes, strong red (51A) in bud, opening inside light yellow (17D), edged strong red (51A); outside light yellow (17D) at base, becoming light yellowish pink (29C), with c.5mm wide stripes of strong red (51A) up centre of each lobe and on edges; unmarked. Calyx lobes c.15-20mm long, held tight to the corolla, light yellow (17D). Lvs elliptic, 150 × 55mm, broadly acute at apex, rounded at base, margins wavy and downcurved, dull and moderate olive green (137A) above, greyish yellow-green (148C) below, hairless. Shrub 0.9 × 1.0m in 11 years. Early May.

(r) **'Stony Brook'**

cv. Elepidote rhododendron: 'Voluptuous' (s) × 'Point Defiance'. H: W. Brack (1988), G: W. Brack (1996), N: W. Brack (2004), I: Roslyn Nursery (2004), REG: W. Brack (2005). Fls 16/domed truss, open funnel-shaped, 65 × 80mm, with 5-6 wavy-edged lobes, vivid red (46C) in bud, opening pale yellowish pink (27D) inside, with prominent strong purplish

pink (55B) markings on the sides of the lobes and a brilliant yellow-green (154C) flare on three dorsal lobes; outside pale yellowish pink (27D), with light purplish pink (55C) markings. Calyx c.3mm long, light yellow-green (154D). Lvs elliptic, 120 × 55mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137A) above, hairless. Shrub 1.2 × 1.5m in 16 years. Midseason.

(r) **'Striking'**

cv. Elepidote rhododendron: Calstocker Group (s) × 'Auguste van Geert'. H: H. O'Rourke (1989), G: H. O'Rourke (2000), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 14/flat-topped truss, broadly funnel-shaped, 75 × 110mm, with 5 wavy-edged lobes, white in bud and throughout on opening, with a small pink eye. Calyx 2mm long, green. Lvs obovate, 150 × 60mm, margins upcurved, glossy green above, cuneate at base, obtuse at apex, hairless. Shrub 1.2 × 1.6m in 14 years. October (Australia).

(r) **'Sucre d'Orge'**

cv. Elepidote rhododendron: 'September Song' (s) × ('Fred Wynniatt' (i.e. the FCC clone) × 'Ring of Fire'). H: M. Colombel (1998), G: M. Colombel (2003), N: M. Colombel (2005), REG: M. Colombel (2005). Fls 7-10/domed truss, broadly funnel/saucer-shaped, 50 × 80mm, with 7 flat-edged lobes, deep red (53A) in bud, opening inside light purplish pink (55C), outside strong pink (54C), with very few spots on the three dorsal lobes; stigma green. Lvs oblong, 120-140 × 50-55mm, margins flat, matt green above, oblique at base, broadly acute at apex, hairless. Shrub 1.2 × 1.2m in 7 years. Early to mid May. (Nomenclatural standard: colour slide provided by registrant (WSY0070278). Etymology: epithet suggested by the colour of the flowers).

(a) **'Summer Heat'**

cv. Deciduous azalea: (*cumberlandense* × *prunifolium*) (s) × *cumberlandense*. H: J.E. Schild (1984), G: J.E. Schild (1989), N: J.E. Schild (2003), I: Schild Azalea Gardens & Nursery (2006), REG: J.E. Schild (2005). Fls 6-10/domed truss, tubular funnel-shaped, 55 × 65mm, with 5 flat-edged lobes, vivid red (45A) in bud, opening inside vivid red (45B), tinged strong reddish orange (N34B), with an inconspicuous strong red (46A) dorsal blotch; outside vivid reddish orange (N30A), with sparse white hairs along the tube; filaments and style vivid red (46B); stigma moderate yellow (138B); anthers light yellow (11B). Calyx 5mm long, light green. Lvs obovate, 65 × 55mm, broadly acute at apex, cuneate at base, margins upcurved, dull and strong yellow-green (143A) above, moderate yellow-green (147B) below, with hairs along margin and on midvein below (hairs white at first, aging green-white). Shrub 1.2 × 0.9m in 10 years. Mid to late June. (Published in *The Azalean* 27(2):33 (2005)).

(r) **'Sun Bonnet'**

cv. Elepidote rhododendron: ('Paprika Spiced' × ('C.I.S.' × 'Crest' selfed)) (s) × 'Marie Starks'. H: J.D. Winberg (1988), G: J.D. Winberg (1992), N: J.D.

Winberg (2005), REG: J.D. Winberg (2005). Fls 13-14/domed truss, open funnel-shaped, 45 × 50mm, with 6 wavy-edged lobes, strong red (53B) in bud, opening inside light yellow (10B), with a strong red (47B) flare in the throat; outside light yellow (10C). Lvs elliptic, 125 × 65mm, obtuse at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137A) above, hairless. Shrub 2.0 × 2.4m in 17 years. Mid April.

(a) **'Sunday Morning'**

cv. Deciduous azalea: *arborescens* (s) × *cumberlandense*. H: J.E. Schild (1986), G: J.E. Schild (1994), N: J.E. Schild (2003), I: Schild Azalea Gardens & Nursery (2006), REG: J.E. Schild (2005). Fls 6-8/flat truss, tubular funnel-shaped, 70 × 55mm, with 5 flat-edged lobes, buds pale yellow (11D) on tube shading to white, with a deep pink (47D) tip, opening inside bright white with two prominent, brilliant yellow (7A) blotches on dorsal lobe; outside white, the midribs deep pink (47D), with many yellowish white hairs and glandular setae on the tube; style strong red (41B); stigma moderate red (N34A); filaments strong yellowish pink (38A); anthers light orange (26C); strongly scented. Calyx c.3mm long, yellow-green. Lvs obovate, 65 × 25mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (137B) above, moderate yellow-green (138B) below, hairs on margin and midveins below (hairs white, aging to yellow-green). Dense shrub, 0.9 × 0.75m in 10 years. Early June. (Published in *The Azalean* 27(2):32 (2005)).

(r) **'Sun Shower'**

cv. Elepidote rhododendron: 'Hotei' (s) × 'Satin Gold'. H: J. Barlup (1984), G: J. Barlup (1989), N: J. Barlup (1994), REG: J. Barlup (2005). Fls 19/ball truss, broadly funnel-shaped, 50 × 75mm, with 7 wavy-edged lobes, moderate reddish orange (173B) in bud, opening throughout pale greenish yellow (10D), with touches of moderate yellowish pink (38B) on some edges; with a moderate reddish orange (42D) dorsal flare of brush-like streaks of very small spots, extending upwards c.20mm from the base. Calyx c.15mm long, pale greenish yellow (10D) with moderate reddish orange (42D) dorsal spots. Lvs elliptic, 90 × 35mm, broadly acute at apex, rounded at base, margins upcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.45 × 0.75m in 5 years. Late April.

(r) **'Sweet Peace'**

cv. Elepidote rhododendron: 'The Master' (s) × *wardii*. H: H. O'Rourke (1989), G: H. O'Rourke (1999), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 12-13/rounded truss, broadly funnel-shaped, 45 × 90mm, with 5 wavy-edged lobes, creamy pink in bud, opening inside pale yellow in throat with cream lobes, outside pale yellow with cream lobes; very small flare of red spots which turn green. Calyx 4mm long, red. Lvs oblong, 145 × 50mm, margins decurved, glossy green above, cuneate at base, acute at apex. Shrub 1.4 × 1.2m in 13 years. October to November (Australia).

(r) **'Sweet Persuasion'**

cv. Lepidote rhododendron: *johnstoneanum* (s) × Chrysomanicum Group. H: H. O'Rourke (1985), G: H. O'Rourke (1994), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 5/loose, flat-topped truss, broadly funnel-shaped, 50 × 70mm, with 5 wavy-edged lobes, bronze yellow in bud, opening inside cream with a yellow centre, outside cream. Calyx green. Lvs elliptic, 50 × 30mm, margins flat, matt green above, cuneate at base, broadly acute at apex, scaly. Shrub 1.4 × 1.5m in 9 years. September to October (Australia).

(r) **'Ted's Big Red'**

cv. Elepidote rhododendron: 'Smokey' (s) × 'Goldsworth Orange'. H: E.T. Anderson (1998), G: D.W. Holcomb (2004), N: Mrs E.T. Anderson (2004), REG: Mrs E.T. Anderson (2005). Fls 14-17/domed truss, broadly funnel-shaped, 65 × 65mm, with 5 wavy-edged lobes, deep red (185A) in bud, opening deep red (53A) throughout; dorsal lobe with an inconspicuous, greyish red (184A) blotch. Calyx c.13mm long, deep red (185A). Lvs elliptic, 100-140 × 40-65mm, acute at apex, rounded at base, margins slightly downcurved, dull and moderate olive green (147A) above, new growth strong yellow-green (144A), hairless. Shrub 0.9 × 0.75m in 20 years. Late May.

(r) **'Te Harinui'**

cv. Lepidote rhododendron: selection from *maddenii* subsp. *maddenii*. S: Canterbury Rhododendron Society (1985), G: Canterbury Rhododendron Society (2002), N: Canterbury Rhododendron Society (2002), REG: Canterbury Rhododendron Society (2005). Fls 4-6/lax truss, tubular funnel-shaped, with 5 wavy-edged lobes, cream and white in bud, and both inside and out on opening, unmarked; spicily scented. Calyx 5mm long, cream. Lvs oblanceolate, 120-180 × 40-60mm, margins flat, dark glossy green above, narrowed at base, pointed at apex, scaly. Tree-like shrub, 2.0 × 1.0m in 15 years. Late. (Etymology: first flowered on Advent Sunday, 2002 - Te Harinui is the title of a New Zealand Christmas hymn. Published in New Zealand Rhododendron Association Newsletter, March 2004, p.10).

(r) **'Terai-homare'**

cv. Elepidote rhododendron: *yakushimanum* (s) × 'A1'. H: K. Terai (1975), G: K. Terai (1982), N: K. Terai (2005), REG: K. Terai (2005). Fls 15/compact, ball truss, open funnel-shaped, 35 × 60-70mm, with 5 wavy-edged lobes, buds strong purplish red (67A), opening inside light purplish pink (68D) and paling towards the centre, outside vivid purplish red (67B); with a small, dark blotch at the base of the corolla. Lvs elliptic, 80 × 30mm, margins decurved, glossy green above, with a silvery white indumentum below (hairs remaining this colour). Compact shrub 0.4 × 0.4m in 4 years. Mid May. (Nomenclatural standard: colour print provided by registrant (WSY0070569)).

(r) 'Terra'

cv. Elepidote rhododendron: 'Recital' (s) × 'Nelda Peach'. H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2005), REG: J. Barlup (2005). Fls 15/ball truss, broadly funnel-shaped, 55 × 90mm, with 6 wavy-edged lobes, moderate red (47A) in bud, opening inside pale yellow (11C-D) blending to shades of pink (near pale purplish pink, 56A) at margins; outside pale yellow (11C) turning to strong pink (54C); solid dark red (59A) basal blotch, c.15mm deep, transforming into a spotted dorsal flare of the same colour extending upwards another 25mm. Calyx c.15mm long, a blend of pale yellow (11D) and strong pink (54C), with two dark red (59A) dorsal flares. Lvs elliptic, 100 × 45mm, broadly acute at apex, rounded at base, with wavy margins, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 9 years. Late May.

(r) 'The Duchess'

cv. Elepidote rhododendron: 'Trude Webster' (s) × 'Grand Slam'. H: H. O'Rourke (1986), G: H. O'Rourke (1997), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 22/domed truss, broadly funnel-shaped, 60 × 105mm, with 5 wavy-edged lobes, strong purplish pink (55B) in bud and both inside and out on opening; red flare on dorsal lobe comprising red dots and streaks. Calyx 5mm long, green. Lvs oblong, 180 × 65mm, margins decurved, matt green above, cuneate at base, acute at apex, hairless. Shrub 1.3 × 1.0m in 16 years. November (Australia).

(a) 'Thekla'

Syn. of 'Hachmann's Thekla'.

(a) 'Tim-Tim'

cv. Evergreen azalea: ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'. H: W.L. Clagett (1996), G: W.L. Clagett (2000), N: W.L. Clagett (2005), I: White's Nursery (2006), REG: W.L. Clagett (2005). Fls 3/flat truss, broadly funnel-shaped, c.20 × 35mm, with a variable number of wavy-edged lobes (5-8 plus 1-4 petaloids), deep yellowish pink (43C) in bud, opening deep yellowish pink (43C) throughout, with large, dark salmon spotting on the dorsal lobe; ventral lobes "butterfly" outwards. Calyx 5mm long. Lvs elliptic, 30 × 10mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate yellow-green (146B) above, with very fine, white hairs above and below when young. Shrub 0.6 × 0.6m in 9 years. Mid May.

(r) 'Tisofee'

cv. Elepidote rhododendron: 'Blimey Limey' (s) × ('Cotton Candy' × 'Van Nes Sensation'). H: M. Stockman (1993), G: M. & H. Stockman (2000), N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 5-6/lax truss, funnel-campanulate, 65 × 90mm, with 6 wavy-edged lobes, strong purplish pink (55B) in bud, opening inside light purplish pink (55C), shading to pale yellow-green (4D) in centre, with two speckled,

strong red (50A) lines (c.25 × 3mm) in the centre of two dorsal lobes; outside light purplish pink (55C) with a pale yellow-green (4D) base and stripes of strong purplish pink (55B) up the centre of each lobe. Calyx lobes irregular, 20-40mm long, pale yellow-green (4D), edged light purplish pink (55C), with strong red (50A) speckling on the dorsal lobes. Lvs lanceolate, 120 × 40mm, acute at apex, oblique at base, margins flat and downcurved, semi-glossy and moderate yellow-green (147B) above, greyish yellow-green (148D) below, hairless. Shrub 1.2 × 1.5m in 11 years. Mid May.

(r) 'Tom Garbutt'

cv. Elepidote rhododendron: 'Charlotte de Rothschild' (s) × 'Alice Hennessey'. H: T.A. Garbutt (1993-1994), G: Mrs A. Sim (2004), N: Mrs A. Sim (2005), REG: T.A. Garbutt (2005). Fls 15/lax truss, funnel-shaped, 90 × 150mm, with 6 wavy-edged lobes, deep bright pink in bud, opening bright pink and paling with age; unmarked; stigma green, style white. Calyx 2mm long, deep pink. Lvs oblong, unusually long, 240 × 80mm, margins flat, matt green above, acute at apex, cuneate to truncate at base. Shrub 1.5 × 1.5m in 10 years. Midseason. (Nomenclatural standard: colour slide provided by the registrant (WSY0070520)).

(r) 'Tony Law'

cv. Elepidote rhododendron: selection from a selfed form of *brachycarpum* known as "Gable's Yellow" or "Gable's Cream" which may itself be a hybrid. H: R.M. Steele (c.1970), G: R.M. Steele, N: R.M. Steele, I: Bayport Plant Farm, REG: J.K. Weagle (2005). Fls 26/domed truss, broadly funnel-shaped, c.45 × 60mm, with 5 wavy-edged lobes, very pale purple (69B) in bud, opening inside pale purplish pink (65D) and fading to greenish white (155C); outside greenish white (155C) with very pale purple (69B) midribs; deep greenish yellow (153A) spotting on the upper two-thirds both inside and out. Calyx 3mm long. Lvs oblong to oblong-elliptic, convex, centre of leaf depressed longitudinally from petiole for about four-fifths of the leaf, sides curved downwards, obtuse to broadly acute at apex, rounded to very slightly oblique at base, glossy and near moderate olive green (147A) above, with a very thin coating on underside of lamina, with little on the midrib, this indumentum brownish orange (165B). Shrub 4.5 × 3.3m in 35 years. Early to mid July. (Has larger flowers and improved foliage, size and colour compared to parent. Very vigorous).

(r) 'Tootie Flutey'

cv. Elepidote rhododendron: 'Van Nes Sensation' (s) × 'Kubla Khan'. H: M. Stockman (1992), G: M. & H. Stockman (1999), N: M. & H. Stockman (2004), I: Hammond Nursery (2006), REG: M. & H. Stockman (2005). Fls 4-5/ball truss, saucer-shaped, 65 × 100mm, with 7 very wavy-edged lobes, light purplish pink (55C) in bud, opening inside pale purplish pink (56A) at margins, shading paler to pale purplish pink (56D) towards the centre, with a vivid red (45A) solid base and conspicuous (45A) dorsal spotting on a pale

yellow (11D) ground; outside margins pale purplish pink (56B), shading to pale purplish pink (56A) at base and pale purplish pink (56D) along lobe centres. Calyx with 2 dorsal lobes c.15mm long, the others c.3mm long, light yellow-green (145D). Lvs elliptic, 140 × 50mm, broadly acute at apex, rounded at base, margins flat and downcurved, dull and moderate yellowish green (138A) above, moderate yellow-green (147C) below, hairless. Shrub 0.6 × 0.75m in 12 days. Early May.

(a) **'Tootsie'**

cv. Evergreen azalea: 'Squirrel' (s) × 'Chippewa'. H: Holger Hachmann (1990), G: Holger Hachmann (1994), N: Holger Hachmann (2003), I: Holger Hachmann (2004), REG: G. Stück (2005). Fls single, 2-3/truss, broadly funnel-shaped, 30-35 × 40-45mm, with 5 slightly wavy-edged lobes, buds strong purplish red (54A), opening inside deep purplish pink (61D), shaded vivid purplish red (61C), outside similar but a bit darker on the veins; dorsal lobe with strong purplish red (54A) markings; stamens 5, filaments deep purplish pink (61D), anthers dark red (187B); style deep purplish pink (61D), stigma deep purplish red (61A); ovary greenish, silvery hairy. Calyx 4-5mm long, brilliant/pale yellow-green (149C/D), silvery hairy. Lvs elliptic to ovate, 20-28 × 10-14mm, margins flat, mid green and slightly glossy above, cuneate at base, apiculate at apex, with white hairs above and brownish hairs below; lasting well in winter. Shrub c.0.35 × c.0.6m in 10 years, with a dense, low-growing habit. Early to late June (i.e. late). (Nomenclatural standard: colour illustration provided by the registrant (WSY0070166). Published in Hachmann cat. 2004-2008 (2004), p.177).

(r) **'Touch of Magic'**

cv. Elepidote rhododendron: *aberconwayi* (s) × 'Coronation Day'. H: H. O'Rourke (1994), G: H. O'Rourke (2002), N: H. O'Rourke (2004), REG: H. O'Rourke (2005). Fls 11-12/rounded truss, broadly funnel-shaped, 60 × 120mm, with 7 lobes, white in bud, opening white inside with a small red flare at the centre of the flower, outside white. Calyx 2mm long, red. Lvs oblanceolate, 80 × 35mm, margins flat, glossy above, cuneate at base, acute at apex, hairless. Shrub 1.2 × 1.0m in 9 years. October to November (Australia).

(r) **'Tout Feu Tout Flamme'**

cv. Elepidote rhododendron: ('Kernéostic' × 'Jalisco Elect') (s) × ('Fred Wynniatt' (i.e. the FCC clone) × 'Ring of Fire'). H: M. Colombel (1998), G: M. Colombel (2003), N: M. Colombel (2005), REG: M. Colombel (2005). Fls 9/domed truss, broadly funnel-shaped, 50 × 70mm, with 7 flat-edged lobes, strong red (47B) in bud, opening brilliant yellow (11A), outside strong red (51A); a few red spots on the dorsal lobes and darker marking around the base of the throat; stigma green. Lvs oblong, 100 × 50-55mm, margins flat, matt green above, rounded at base, obtuse at apex, hairless. Shrub 0.7 × 1.2m in 7 years. Mid to late May. (Nomenclatural standard: colour

slide provided by the registrant (WSY0070277). Etymology: epithet means "all flower and flame", suggested by the flower colour).

(r) **'Toyama 21'**

cv. Elepidote rhododendron: *yakushimanum* (s) × 'A1'. H: K. Terai (1975), G: K. Terai (1982), N: K. Terai (2005), REG: K. Terai (2005). Fls 20/ball truss, open funnel-shaped, 40 × 70mm, with 5 very wavy-edged lobes, buds strong purplish red (67A), opening inside light purplish pink (68D), paling towards base and with a tiny basal blotch, outside strong purplish pink (67B). Lvs narrowly elliptic, 100 × 35mm, glossy green above, obtuse at base, acute at apex, margins slightly recurved, with a silvery white indumentum below (not changing colour with age). Shrub 0.6 × 0.5m in 4 years, with broadly spreading branches. Mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070567). Etymology: Toyama is the prefecture in Japan where the plant originated).

(r) **'Traffic Light'**

cv. Elepidote rhododendron: *elliottii* (s) × 'Gill's Crimson'. H: H. O'Rourke (1987), G: H. O'Rourke (1999), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 20/rounded truss, campanulate, 50 × 60mm, with 5 very wavy-edged lobes, vivid red (45A) in bud and both inside and out on opening. Calyx 3mm long, red. Lvs oblong, 170 × 60mm, margins decurved, matt green above, obtuse at base, acute at apex, hairless. Shrub 1.5 × 1.2m in 15 years. October (Australia).

(v) **'Tropic Alpine Survivor'**

cv. Lepidote rhododendron: (*zoelleri* × *lochiae*) (s) × *commonae*. H: J.W. Gerdemann (1985), G: J.W. Gerdemann, N: J.W. Gerdemann, I: The Bovees Nursery, REG: J.W. Gerdemann (2005). Fls 4-7/flat truss, elongated tubular funnel-shaped, with 5, flat-faced, reflexed lobes, 50 × 45mm, vivid reddish orange (44C) inside and out, the inside blending to vivid yellow (17C) in the basal part of the tube and the outside with some strong yellowish pink (33C) on the tube. Calyx inconspicuous. Lvs elliptic, 65 × 40mm, broadly acute to obtuse at the apex, cuneate at base, margins flat, semi-glossy and moderate olive green (137A) above, scaly. Shrub 0.9 × 0.5m in 18 years: has survived being frozen back 50% or more (once to the ground) but always recovers. The only survivor of 300 plants from this cross. Flowering at any time of the year.

(v) **'Veryl Ann'**

cv. Lepidote rhododendron: aff. *laetum* (s) × (*macgregoriae* × *aurigeranum*). H: R.A. Mitchell (1994), G: R.A. Mitchell (1998), N: R.A. Mitchell (2005), REG: R.A. Mitchell (2005). Fls 7/truss, held horizontally, tubular funnel-shaped, 55 × 90mm, with 5 wavy-edged lobes, opening inside and out light orange (29B) at margins, blending to vivid yellow (17C). Lvs elliptic, 100 × 50mm, acute at apex, cuneate at base, margins flat (but lvs V-shaped in cross

section), semi-glossy and moderate green (135B) above; no scales visible to the naked eye. Shrub 1.2 × 0.9m in 5 years. December to January.

(a) **'Villa San Remigio'**

cv. Evergreen azalea: parentage unknown. R: Gardeners of the Villa San Remigio (pre 1920), N: Prof. E. Accati (2004), REG: Prof. E. Accati (2005). Fls single, 1-3/truss, 76 × 56mm, with 5 very wavy-edged lobes, buds red-purple (N74A), opening red-purple (N74B) throughout, three lobes with markings of strong red (53C), shaded deep red (53A); stamens 10, not exerted from the corolla, anthers red-purple (N77C), filaments red-purple (N74B); pistil distinctly exceeding the stamens and the corolla, style strong purplish red (58B), stigma strong red (53C). Lvs lanceolate, 50 × 21mm, margins flat, glossy above, acute at base and apex, hairy, moderate olive green (147A) above and moderate yellow-green (147C) below when mature; petioles 11mm long. Shrub 5.0 × 7.0m in c.100 years. May. (Etymology: named after its place of origin).

(a) **'Violajana'**

cv. Evergreen azalea: 'Friedhelm Scherrer' (s) × 'Mevrouw Jozef Heursel'. R: G. Harmsen (pre 1985), N: G. Harmsen, REG: G. Harmsen (2005). Fls hose-in-hose, double, few/truss, erect, dish-shaped, whorls 5-6-lobed, lobes not wavy-edged, inside vivid reddish purple (78A) at margin, centre strong reddish purple (78B), outside similarly coloured; throat darker, with small, deep purplish red (71A) spots. Calyx petaloid. Lvs elliptic, acute at apex, cuneate at base, bluish green and slightly glossy above, convex. Very early. (Previously listed in the Checklist but not registered.)

(r) **'Violet Mist'**

cv. Elepidote rhododendron: *yakushimanum* (s) × ('Frank Galsworthy' × 'Purple Splendour'). H: J. Barlup (1995), G: J. Barlup (2000), N: J. Barlup (2005), REG: J. Barlup (2005). Fls 16/ball truss, broadly funnel-shaped, 45 × 70mm, with 5 very wavy-edged lobes, strong purplish red (72A) in bud, the dorsal lobe opening yellowish white (158D) at the base and extending upwards c.45mm where it blends into light purple (77D), the predominant colour of the interior; outside light purple (77D) blending to strong purple (77B) edges and with random 77B streaks; a prominent, spotted, moderate red (183C) dorsal flare, c.25mm wide, extends upwards c.30mm from the base. Lvs elliptic, 105 × 40mm, broadly acute at apex, cuneate at base, margins upcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 10 years. Mid May.

(r) **'Wally Harper'**

cv. Elepidote rhododendron: 'Minnie' (s) × 'Barbara Cook'. H: A.T. & S.A. Anderson (1990), G: A.T. & S.A. Anderson (1998), N: Mrs P. Walton (2005), REG: Mrs P. Walton (2005). Fls 10-14/domed truss, open funnel-shaped, 50 × 75mm, with 5-7 wavy-edged lobes, strong purplish pink (63C) in bud, opening pale

yellowish pink (159D) throughout, with twin, speckled, dark red (187C) dorsal flares and rays of strong purplish pink (63C) freckles extending from base to sinuses. Lvs oblong, 125 × 40mm, broadly acute at apex, rounded at base, margins wavy, dull and moderate yellow-green (147B) above, hairless. Shrub 0.9 × 1.1m in 15 years. Late May to early June.

(r) **'White Elation'**

cv. Elepidote rhododendron: 'Scintillation' (s) × 'Carolyn Grace'. H: D.B. Johnson (1985), G: D.B. Johnson (1991), N: D.B. Johnson (2005), REG: D.B. Johnson (2005). Fls 8/domed truss, saucer-shaped, 50 × 75mm, with 7 wavy-edged lobes (plus petaloids), strong purplish pink (55B) in bud, opening white (155D) throughout; unmarked; no normal stamens present. Lvs oblong, 140 × 65mm, obtuse at apex, cordate at base, margins downcurved, glossy and moderate olive green (137A) above, hairless. Shrub 2.4 × 2.4m in 20 years. Early May.

(r) **'White Opal'**

cv. Elepidote rhododendron: 'Peach Charm' (s) × 'Pirouette'. H: J. Barlup (1992), G: J. Barlup (1996), N: J. Barlup (2005), REG: J. Barlup (2005). Fls 23/ball truss, broadly funnel-shaped, 55 × 75mm, with 5 wavy-edged lobes, strong purplish pink (63B) in bud, opening throughout light purplish pink (65B) and then changing to white (155D); two small, dorsal flares of strong yellowish pink (37A) spots extending upwards c.15mm from the base. Lvs elliptic, 125 × 40mm, broadly acute at apex, cuneate at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.1 × 1.4m in 12 years. Late May.

(r) **'Winter Spice'**

cv. Elepidote rhododendron: 'Hachmann's Charmant' (s) × 'Dreamland' (Waterer). H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2005), REG: J. Barlup (2005). Fls 21/conical truss, broadly funnel-shaped, 50 × 70mm, with 5 very wavy-edged lobes, strong purplish red (63B) in bud, opening inside pale yellow-green (155A), blending to pale purplish pink (65C) on the dorsal lobe near the margin, with a large dorsal flare of deep pink spots extending upwards c.25mm from the base on the two dorsal lobes; outside yellowish grey (156A) shading to pale purplish pink (65C) on the dorsal lobes. Calyx c.5mm long, pale yellow-green (155A). Lvs elliptic, 105 × 40mm, broadly acute at apex, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 9 years. Late May.

(r) **'Wonderlight'**

cv. Elepidote rhododendron: 'Van Nes Sensation' (s) × 'Queen Souriya'. H: H. O'Rourke (1988), G: H. O'Rourke (1997), N: H. O'Rourke (2003), REG: H. O'Rourke (2005). Fls 9-10/loose truss, broadly funnel-shaped, 65 × 90mm, with 7 wavy-edged lobes, mauve in bud, opening inside pale mauve, with a small red ray, the lobes fading with age; outside pale mauve, fading with age. Calyx 2mm long, green. Lvs oblong,

110 × 30mm, margins flat, matt green above, cuneate at base, acute at apex, hairless. Shrub 1.6 × 2.0m in 14 years. October to November (Australia).

(r) **'Yellow South'**

cv. Elepidote rhododendron: unknown (s) × 'Golden Delicious'. H: D.B. Johnson (1995), G: D.B. Johnson (2000), N: D.B. Johnson (2005), REG: D.B. Johnson (2005). Fls 12/domed truss, open funnel-shaped, 50 × 90mm, with 5 wavy-edged lobes, strong yellowish pink (43D) in bud, opening throughout light yellow (12C), edged pale yellow (20D), unmarked. Lvs oblong, 125 × 45mm, broadly acute at apex, rounded at base, with wavy margins, dull and moderate olive green (137B) above, hairless. Shrub 1.1 × 1.8m in 8 years, growing well in the southern heat in the USA. Mid May.

(r) **'Yume-tzuzuri'**

cv. Elepidote rhododendron: *yakushimanum* (s) × 'A1'. H: K. Terai (1975), G: K. Terai (1982), N: K. Terai (2005), REG: K. Terai (2005). Fls 15/ball truss, open funnel-shaped, 35 × 60-70mm, with 5 very wavy-edged lobes, strong purplish red (67A) in bud, opening inside light purplish pink (68D), paling towards the base and with a small basal blotch, outside

vivid purplish red (57B). Lvs narrowly elliptic, 80 × 30mm, glossy green above, margins recurved, obtuse at base, acute at apex, with a silvery white indumentum below (hairs not changing colour with age). Shrub 0.5 × 0.4m in 4 years. Mid May. (Nomenclatural standard: colour print provided by the registrant (WSY0070566)).

(a) **'Zoverallo'**

cv. Evergreen azalea: parentage unknown. R: Fratelli Savioli (1970), N: Fratelli Savioli (1985), I: Fratelli Savioli (1985), REG: Prof. E. Accati (2005). Fls single, 2-5/truss, 44 × 30mm, with 5 wavy-edged lobes, buds deep pink (48A), opening white shaded strong pink (48C) inside and out, three lobes with deep yellowish pink (46D) markings; stamens 5, exserted from the corolla, anthers moderate orange (167B), filaments white; pistil distinctly exceeding the stamens, style white, stigma brilliant greenish yellow (1B). Lvs ovate, 24 × 13mm, margins flat, glossy above, obtuse at base and apex, hairy, strong yellow-green (144A) above and moderate yellow-green (146C) below when mature; petioles 7mm long. Shrub 1.0 × 1.0m in 10 years. April. (Etymology: named for a village near the city of Verbania, the latter being where the Savioli nursery is situated).

NAMES AND ADDRESSES

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

* = deceased

- | | |
|--|--|
| Accati, Prof.E.
Università degli Studi di Torina, Grugliasco, Italy | Brown, D. & P.J.
Browns Nursery, Tauranga, New Zealand |
| Ahern, T.
Bethlehem, Pennsylvania, USA | Brown, P.
Whispering Pines Nursery, Lakeville,
Massachusetts, USA |
| Anderson, A.T. & S.A.
Franklin Lakes, New Jersey, USA | Canterbury Rhododendron Society Inc.
Christchurch, New Zealand |
| Anderson, B.E.
Longview, Washington, USA | Carmine, M.
Vivaio La Margotta, Cannero Riviera, Italy |
| Anderson, Mr & Mrs E.T.
Longview, Washington, USA | Carmine, Mrs O.
Vivaio La Margotta, Cannero Riviera, Italy |
| Anderson, J.D.
Anderson Garden, Enumclaw, Washington, USA | Carroll, C.
West Pymble, New South Wales, Australia |
| Anderson Garden
Enumclaw, Washington, USA | Carroll's Country Gardens Nursery
Mount Tomah, New South Wales, Australia |
| Arblaster, *G. & Mrs J.
Eyemouth, Berwickshire, Scotland | Cavadini, G.
Cavadini Azienda Vivaistica, Intra, Italy |
| Arborglen Pty Ltd
Glenorie, New South Wales, Australia | Clagett, W.L.
Derwood, Maryland, USA |
| Arneson, I. & R.
Arneson Nursery, Canby, Oregon, USA | Cleland, N.
Stratford, New Zealand |
| Barber, D.
Phoenix, Arizona, USA | Cockeram, E.
Florence, Oregon, USA |
| Barlup, J.
Bellevue, Washington, USA | Colombel, M.
Menez-Rohou, Fouesnant, France |
| Bayport Plant Farm
Lunenburg County, Nova Scotia, Canada | Coners, G.
Westerstede, Germany |
| Beaudry, N. | Corneliussen's Nursery
Gervais, Oregon, USA |
| Bethesda, Maryland, USA | Cox, P.A. & K.N.E.
Glendoick Gardens, Glencarse, Perthshire,
Scotland |
| Bell, D.S.
Hendersonville, North Carolina, USA | Craig, D.L.
Kentville, Nova Scotia, Canada |
| Bovees Nursery, The
Portland, Oregon, USA | Doppel, J.
Lenhartsville, Pennsylvania, USA |
| Brack, W.
St James, New York, USA | Dosser, D.J.
Lower Wilmot, Tasmania, Australia |
| Briggs Nursery
Olympia, Washington, USA | |

- Forest Lane Nursery
Fort Bragg, California, USA
- Fujioka, F.
Freeland, Washington, USA
- Furman, F.
Bridgewater, New Jersey, USA
- Garbutt, T.A.
Oamaru, North Otago, New Zealand
- Gerdemann, J.W.
Yachats, Oregon, USA
- Glassman, A.
Allentown, Pennsylvania, USA
- Glendoick Gardens
Glencarse, Perthshire, Scotland
- Griswold, D.A.
Griswold Nursery, Kirkland, Washington, USA
- Griswold Nursery
Kirkland, Washington, USA
- Gunn, S.
Centerville, Nova Scotia, Canada
- Guyens, H.J.A.
Kamp-Lintfort, Germany
- *Hachmann, H.
(Hans Hachmann), Barmstedt, Germany
- Hachmann, Holger
Barmstedt, Germany
- Hammond Nursery
Arlington, Washington, USA
- Holcomb, D.W.
Longview, Washington, USA
- Hooper, J.D.
Uniontown, Ohio, USA
- Huisingh, J.L.
Puyallup, Washington, USA
- Imrie, S.T.
Bonney Lake, Washington, USA
- Jago, E.
Stratford, New Zealand
- Johnson, D.B.
Mountain Rest, South Carolina, USA
- Jones, K.
Fort Bragg, California, USA
- Jordan, H.
Ashburton, New Zealand
- Jørgensen, P.M.
Hjellestad, Norway
- Kasai, K.
Nakatsugaru-district, Aomori-prefecture, Japan
- Klimavicz, J.F.
Vienna, Virginia, USA
- Lake Tapps Rhododendron Nursery
Bonney Lake, Washington, USA
- Lehmann, C.A.
Denmark
- Loder Plants
Lower Beeding, West Sussex, England
- Maehara, H.
Higashi-Osaka-city, Osaka-prefecture, Japan
- Malland, L.
Griswold Nursery, Kirkland, Washington, USA
- McDonald, S.F.
Le-Mac Nurseries, Hampton, Virginia, USA
- McNew, C.
Kelso, Washington, USA
- Meerkerk Gardens
Greenbank, Washington, USA
- Mendlin, B.
Longview, Washington, USA
- Metcalf, E.
Beaverton, Oregon, USA
- Millais, D.G. & *E.G.
Millais Nurseries, Farnham, Surrey, England
- Millais Nurseries
Farnham, Surrey, England
- Miller, G.T.
New Oxford, Pennsylvania, USA
- Mitchell, R.A.
Volcano, Hawaii, USA
- Moyles, E.W.
Oakland, California, USA
- Newcomb, L.L. & E.F.
Snohomish, Washington, USA
- Nouws, G.
Zundert, The Netherlands

- Nys, C.
Longview, Washington, USA
- Olson, J.
Fall Creek, Oregon, USA
- O'Rourke, H.
Burnie, Tasmania, Australia
- Parks, J.B.
Dover, New Hampshire, USA
- Philipson, M.N. & *W.R.
Christchurch, New Zealand
- Pilkington, Mrs E.
Monument Beach, Massachusetts, USA
- Reiley, H.E.
Woodsboro, Maryland, USA
- Renick, F.
Camarillo, California, USA
- Rhodoland Nursery
Niagara-on-the-Lake, Ontario, Canada
- Robinson, M.L.A.
Forest Row, East Sussex, England
- Roslyn Nursery
Dix Hills, New York, USA
- Rothschild, E. de & *L. de
Exbury Gardens, Hampshire, England
- Sahara, G.
Keaau, Hawaii, USA
- Saitō, T.
Morioka-city, Iwate-prefecture, Japan
- *Salvatore, B.
Zoverallo, Italy
- Savioli Fratelli
Verbania, Italy
- Schild, J.E.
Hixson, Tennessee, USA
- Schild Azalea Gardens & Nursery
Hixson, Tennessee, USA
- Schmalscheidt, W.
Oldenburg, Germany
- *Scott, R.W.
Kensington, California, USA
- Sim, Mrs A.
Oamaru, New Zealand
- Staab, B. & B.
Puyallup, Washington, USA
- Starling, B.N.
Drewsteignton, Devon, England
- Steele, R.M.
Bayport Plant Farm, Lunenburg County,
Nova Scotia, Canada
- Stein, T.
Espergaerde, Denmark
- Stewart, Robert
Springfield, Virginia, USA
- Stockman, M. & H.
Port Angeles, Washington, USA
- Stück, G.
Kronshagen/Kiel, Germany
- Tavner, R. & E.
Ulverstone, Tasmania, Australia
- Terai, K.
Tonami-city, Toyama-prefecture, Japan
- Villa San Remigio
Pallanza, Italy
- Watson, Mrs E.M.
Kirkland, Washington, USA
- Weagle, J.K.
Halifax, Nova Scotia, Canada
- Weeks, C.
Westgate Gardens Nursery, Eureka, California
- Westgate Gardens Nursery
Eureka, California, USA
- Whispering Pines Nursery
Lakeville, Massachusetts, USA
- White's Nursery
Germantown, Maryland, USA
- Wieder, R.
Kent, Washington, USA
- *Wilkes, W.B.
Seattle, Washington, USA
- Winberg, J.D.
Fall City, Washington, USA
- Zachmann, S.
Port Angeles, Washington, USA

CORRECTIONS TO THE INTERNATIONAL RHODODENDRON REGISTER AND CHECKLIST (2004)

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to the First Supplement published in 2006. A further probably extended list should be ready for inclusion with the Third Supplement. The Registrar is grateful to those correspondents who have brought such errors to his notice and welcomes knowledge of further amendments that need to be made. Minor changes (e.g. to spelling within a description) are not noted here but such changes have been made to the database and will be available in any future electronic version. The database will also be taking into account additional information that comes to light relating to any individual entry.

In the following list the page number refers to the page in the *Register and Checklist* on which the entry concerned is to be found.

- | | |
|---|---|
| <p>'Admiraal de Ruyter' (p.21). Add the synonyms: 'Admiraal de Ruiter', 'Admiral Ruyter'</p> | <p>'Bryan Harris' (p.181). Change J. Harris to J.O. Harris</p> |
| <p>'Admiraal Ruyter' Add new entry: (Syn. of 'Admiraal de Ruyter')</p> | <p>'Buddy McMakin' (p.182). Change J. Harris to J.O. Harris</p> |
| <p>'Admiral de Ruiter' Add new entry: (Syn. of 'Admiraal de Ruyter')</p> | <p>'Butheanum' (p.187). Change 'Bluteanum' to 'Blutheanum'</p> |
| <p>'Adolphe d'Haene' (p.21). Add the synonym: 'Madame Adolphe d'Haene'</p> | <p>'California Sunset' (p.193). Change 'Haerens Saumoneum' to 'Haerens Saumoneus'</p> |
| <p>'All Glory' Add new entry: (Syn. of 'Nuccio's All Glory')</p> | <p>'Carolyn's Day' (p.215). Change J. Harris to J.O. Harris</p> |
| <p>'Ambrose Bristow' (First Supplement p.3) Add nomenclatural standard reference: WSY0046884</p> | <p>'Chidori' (p.242). Change category from (r) to (a)</p> |
| <p>'Amiral Ruyter' Add new entry: (Syn. of 'Admiraal de Ruyter')</p> | <p>'Cille Shaw' (p.252). Change J. Harris to J.O. Harris</p> |
| <p>'Ann Lee McPhail' (p.64). Change J. Harris to J.O. Harris</p> | <p>'Coronado Cascade' (p.206). Change J. Harris to J.O. Harris</p> |
| <p>'Anthony Roland' (p.65). Delete entry and replace with: An error for 'Antony Ronald'</p> | <p>'Dorothy Clark' (p.367). Change J. Harris to J.O. Harris</p> |
| <p>'Antony Roland' (p.67). Delete name and replace with 'Antony Ronald'</p> | <p>'Edith Henderson' (p.390). Change J. Harris to J.O. Harris</p> |
| <p>'Antony Roland' Add new entry: An error for 'Antony Ronald'</p> | <p>'Equisitum' (p.415). Change epithet to 'Exquisitum'</p> |
| <p>'Aya-no-kamuri' (p.100). Change reference at end of entry to 'Aya-kanmuri'</p> | <p>'Fourth of July' (p.464). Change deciduous azalea to evergreen azalea</p> |
| <p>'Bouquet de Roses' (p.167). Change the synonym 'Roseiflorum Plenum' to 'Rosiflorum Plenum'</p> | <p>'Francis E. Seidler' (p.476). Change J. Harris to J.O. Harris</p> |
| <p>'Bruce Hancock' (p.179). Change J. Harris to J.O. Harris</p> | <p>'Frosted Orange' (p.479). Change J. Harris to J.O. Harris</p> |
| | <p>'Genji-miyoshi' (p.498). Change 'Asagasumi' to 'Asagasumi'</p> |

- 'Gentain Blue Diamond' (p.498). Change spelling to 'Gentian Blue Diamond'
- 'Georgia Giant' (p.501). Change J. Harris to J.O. Harris
- 'Gesproet' (p.504). Change category from (r) to (a)
- 'Gloria Still' (p.519). Change J. Harris to J.O. Harris
- 'Harris Purple' (p.577). Change J. Harris to J.O. Harris
- 'Hatsuki' (p.580). Add: Kurume
- 'Helena Neumann' (p.586). Change category from (r) to (a)
- 'Helene Thelemann' (p.588). Change synonym 'Helen Thelman' to 'Helen Theleman'
- 'Hikkasen' (p.600). Change 'Hiyakasan' to 'Hiyakasen'
- 'Himura-shiro' (First Supplement p.18). Add: Nomenclatural standard: colour slide provided by the registrant (WSY0041911)
- 'Hitachi' (p.605). Change Elepidote rhododendron to Evergreen azalea
- 'Imperatrix Eugenie' Add new entry: (Syn. of 'Impératrice Eugénie')
- 'Jeni Slocock' (First Supplement, p.19). Change nomenclatural standard reference from WSY0052447 to WSY0046894
- 'Juliana Nova' (p.695). Add the synonym 'Julia Second'
- 'Kagetsu' (p.701). Change informal grouping from Kurume to Satsuki
- 'Kagetsu-muji' (p.701/702). Change informal grouping from Kurume to Satsuki
- 'Karl Glaser' (p.707). Add to entry: R: K. Glaser (pre 1964), N: K. Glaser
- 'Kerneostic' (p.717). Change spelling to 'Kernéostic'
- 'Kinsai' (p.725). Change to 'Kin-sai'
- 'Koasabi' (p.730). Change reference to 'Ko-osobi' in the note to 'Ko-asobi'
- 'Ko-mane' (p.733). Change reference to 'Sene' in the note to 'Sen'e'
- 'La Tempérance' (p.758). Change category from (r) to (a)
- 'Lauren' (p.760). Change category from (r) to (a)
- 'Ledifolium Noordtanium' (p.766). Entry should read: (Syn. of 'Noordtanium')
- 'Lee Thomas' (p.767). Change J. Harris to J.O. Harris
- Lehmann, K.A. (p.1518). Change to Lehmann, C.A.
- 'Lockington Velvet' (First Supplement, p.24). Add: Nomenclatural standard: colour print provided by the registrant (WSY0038787)
- 'Low Boy' (p.819). Change J. Harris to J.O. Harris
- Ludeike (p.1519). An error for Lüdicke, W.
- 'Madame Rhatert' (p.841). Change 'Helen Theleman' to 'Helene Thelemann'
- 'Mademoiselle Marie Planchon' (p.844). Add: Syn. 'Marie Planchon'
- 'Mai-ogi' (second entry, p.851). Change 'Maisogi' to 'Maioogi'
- 'Margaret Rowell' (p.861). Change J. Harris to J.O. Harris
- 'Mary Ann Egan' (p.876). Change J. Harris to J.O. Harris
- 'Meteor' (second entry, p.903). Change 'Roshomon' to 'Rashomon'
- 'Midnight Flare' (p.906). Change J. Harris to J.O. Harris
- 'Monsieur G.J. Bier' (p.928). Add: Syn. 'G.J. Bier'
- 'Monsieur Victor Savant' (p.929). Add: Syn. 'M. Victor Savant'
- 'Novita Antilescens' (p.991). Change 'Novita Antecellens' to 'Novitas Antecellens'
- 'Ogi-no-odorikarki' (p.1003). Change spelling to 'Ogi-no-odorikaraki'
- 'Petrick Alba' (p.1058). Change 'Schepen's Kerstperel' to 'Schepens Kerstperel'
- 'Président Comte Oswald de Kerchove' (p.1096). Add the synonym 'Président Oswald de Kerchove'
- 'Prinz Rudolph' (p.1111). Change 'Prince Rudolphe' to 'Prince Rodolphe'
- 'Rivermist' (p.1165). Change J. Harris to J.O. Harris
- (a) 'Roi des Panachées' Add new entry: (See note under 'Mrs Wright')
- 'Rosy Cheeks' (p.1189). Change 'Pamela Mallard' to 'Pamela Malland'

'Salmon King' (first entry, p.1213). Change 'Mrs L.C. Fisher' to 'Mrs L.C. Fischer'

(a) 'Saturnus' Add new entry: (Syn. of 'Saturne' (Ghent))

'Silver Jubilee' (p.1262). The parentage has now been corrected by the registrant to *yakushimanum* × 'Coronation Day'

'Venus' (p.1409). The correction listed under this epithet in the *First Supplement* (p.47) referred to the plant raised and named by K. Glaser



