



Royal
Horticultural
Society

Sharing the best in Gardening

The International Rhododendron Register & Checklist (2004)

Eighth Supplement



© 2014 The Royal Horticultural Society
80 Vincent Square, London SW1P 2PE, United Kingdom
www.rhs.org.uk

International Rhododendron Registrar:
Dr A C Leslie

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057427

Printed and bound in the UK by Page Bros, Norwich

The previous supplement (Seventh Supplement) was published on 30 May 2012

Cover: *Rhododendron* 'Mrs Charles E. Pearson', by Alfred John Wise
Image courtesy of RHS Lindley Library

The International Rhododendron Register and Checklist (2004)

Eighth Supplement

Names registered, January–December 2012	page 2
Names and addresses	page 22
Corrections	page 24

Introduction

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first seven Supplements included names registered from 1 January 2003 to 31 December 2011. This eighth Supplement includes all those registered in 2012.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7–10 of the Introduction. It may be helpful here to repeat that the following abbreviations are employed:

(a)	azalea	(az)	azaleodendron
cv.	cultivar	G	grown to first flower by
Gp	Group	H	hybridised by
I	introduced by	N	named by
(r)	rhododendron	R	raised by
REG	registered by	S	selected by
(s)	seed parent	(v)	vireya

Where possible, colour references are given to the *RHS Colour Chart* (1966, 1986, 1995 & 2001), with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website (www.rhs.org.uk/plants/plant-science/plant-registration-forms/rhodoform) or from one of the national or regional registrars as follows:

North America	Mr Michael Martin Mills, 632 Burnham Road, Philadelphia, Pennsylvania 19119-3510, USA
Australia	Mrs Lesley Eaton, PO Box 597, Snug, Tasmania 7054, Australia
New Zealand	Mr B.J. Coker, 42 Massey Crescent, St Albans, Christchurch 8014, New Zealand
Japan	Dr K. Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

Dr A.C. Leslie, International Rhododendron Registrar, 109 York Street, Cambridge CB1 2PZ, UK
(Email: alanleslie@rhs.org.uk)

Names registered

January–December 2012

(r) 'Aisai'

cv. Elepidote rhododendron: *brachycarpum* (as *f. nemotoanum*) (s) × *ponticum*

H: H. Suzuki (1997), G: H. Suzuki (2006), N: H. Suzuki (2008), REG: H. Suzuki (2012)

Fls double, 11–13 (average 12)/ball truss, open funnel-shaped, 50 × 70mm, with two whorls of 5(–6), wavy-edged lobes (the inner usually shorter and narrower and representing partially petaloid stamens), deep pinkish red in bud, opening purplish pink-red, inside shading paler towards the centre of the lobes and the centre of the flower; dorsal blotch of yellowish spots. Lvs narrowly elliptic, 140 × 35mm, margins flat, acute at apex, cuneate at base, glossy green above, hairless. Shrub 70 × 40cm, in 4 years after grafting. Mid May.

(Nomenclatural standard: colour print provided by the registrant (WSY))

(r) 'Amber Honey'

cv. Elepidote rhododendron: 'Apricot Fantasy' (s) × 'Sunstone'

H: J. Barlup (1999), G: J. Barlup (2004), N: J. Barlup (2011), REG: J. Barlup (2012)

Fls 18/domed truss, open funnel-shaped, 50 × 65mm, with 6 wavy-edged lobes, moderate red (47A) in bud, opening inside pale orange-yellow (10D), blending to light purplish pink (55C) and edged strong purplish pink (55B), with strong red (53B) nectar pouches and a (53B) spotted flare extending c.12mm from the base of the dorsal lobe; outside pale orange-yellow (19D), blending to light purplish pink (55B) in the midrib area; not scented. Calyx c.20mm long, pale orange-yellow (19D), with strong red (53B) flares extending c.7mm from the base. Lvs elliptic, c.90 × 40mm, margins flat, broadly acute at apex, rounded at base, dull and moderate olive green (147A) above, hairless. Shrub 0.5 × 0.6m in 5 years. May.

(Etymology: flower colour resembles that of some types of honey)

(a) 'Ance'

cv. Deciduous azalea: 'Aida' (de Smet/Vuylsteke) (s) × 'Uguns'

H: R. Kondratovičs (1978), G: R. Kondratovičs (1986), N: R. Kondratovičs (2001),

I: Rhododendron Nursery Babīte, University of Latvia (2001), REG: Rhododendron Nursery Babīte, University of Latvia (2012)

Fls 11–12/slightly domed truss, tubular funnel-shaped, 60mm wide, with 5 strongly wavy-edged lobes (and shorter petaloid stamens), buds strong red (39A) with brilliant yellow (20A), opening inside deep yellowish pink (39B) with light yellow (11B) and with a vivid yellow (12A) blotch on the dorsal lobes, outside deep yellowish pink (39B), with a light yellow (11B) central line down the lobes; style red, stigma green. Calyx 2mm long, green. Lvs elliptic, 115–120 × 40–45mm, pointed at base and apex, margins wavy, glossy green above, hairless. Shrub 2.0 × 1.8m in 10 years. Early June.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: Latvian female christian name)

(r) 'Anne Marie de Beau Vallon'

cv. Elepidote rhododendron: 'Kerneostic' (s) × 'September Song'

H: Société Bretonne du Rhododendron (1997), G: J.F. Saint-Jalm (2003), N: J.F. Saint-Jalm, REG: J.F. Saint-Jalm (2012)

Fls 10/domed truss, funnel-shaped, 55 × 60mm, with 5–6 wavy-edged lobes, buds moderate red (N34A), opening inside brilliant orange-yellow (23B) and strong red (47B), outside similarly coloured; dorsal spotting of brownish orange (N167A); stamens 12, filaments white; style yellow, stigma green; ovary green, with white glands. Calyx 2mm long, strong yellowish pink (37B). Lvs elliptic, 125 × 49mm, margins flat, obtuse at apex, cuneate at base, matt green above, hairless. Shrub 1.35 × 1.65m in 15 years. Early to mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named after the registrant's wife (Anne Marie) and the village in which they live (Beau Vallon))

(a) **'Arima'**

cv. Deciduous azalea: 'Washington State Centennial' (s) × 'Arouca'
H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999), N: K.-H. Hübbers (2006), REG: J. Westhoff (2012)
Fls up to 14/loose truss, tubular funnel-shaped, 50 × 60mm, with 5 flat to slightly wavy-edged lobes, buds white, blushed red, opening inside yellowish white (158D), outside yellowish white (158C) on the lobes, deep pink (50B) on the tube; a vivid orange-yellow (23A) blotch covers about three-quarters of the dorsal lobe and there are deep pink (50B) veins in the throat all around the centre of the flower; filaments and style suffused deep pink, especially towards the base; anthers brown; stigma green; slightly scented. Calyx 3mm long, light green. Lvs elliptic, 70 × 30mm, margins flat, acute at apex, cuneate at base, matt green above, margins ciliate and with tiny white hairs along the midrib when young. Shrub 0.8 × 0.7m in 10 years. Mid May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a city in Trinidad)

(r) **'Austin's Gold'**

cv. Elepidote rhododendron: 'Jawndice' (s) × 'Early Light'
H: J. Doppel (2000), G: J. Doppel (2004), N: J. Doppel (2005), REG: J. Doppel (2012)
Fls 10/conical truss, open funnel-shaped, 40 × 65mm, with 5 wavy-edged lobes, moderate red (181A) in bud, opening inside vivid red (44A), shading to brilliant greenish yellow (6A), with moderate orange (173C) dorsal spots; outside similarly coloured, but without the spotting. Lvs elliptic, 90 × 32mm, broadly acute at apex, cuneate to rounded at base, with flat margins, dull and mid green above, hairless. Shrub 0.8 × 0.9m in 12 years. Early May. (Etymology: Austin is the name of the registrant's grandson)

(r) **'Azuma-jukō'**

cv. Elepidote rhododendron: *degronianum* subsp. *heptamerum* (s) × *degronianum* subsp. *degronianum*

H: H. Suzuki (1995), G: H. Suzuki (2008), N: H. Suzuki (2008), REG: H. Suzuki (2012)
Fls 12–15 (average 13)/hemispherical truss, open funnel-shaped, 50 × 70mm, with 5 slightly wavy-edged lobes, deep red in bud, opening deep red throughout, with inconspicuous red dorsal spotting. Lvs narrowly elliptic to narrowly obovate, 100 × 30mm, margins flat to slightly recurved, broadly acute at apex, cuneate at base, glossy green above, hairs below white when young, aging to brown. Shrub 50 × 40cm, in 6 years after grafting. Early May. (Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **'Babītes Kompaktais'**

cv. Elepidote rhododendron: in vitro mutation of 'Nova Zembla' (s)
G: Rhododendron Nursery Babīte, University of Latvia (2002), N: Rhododendron Nursery Babīte, University of Latvia (2007), I: Rhododendron Nursery Babīte, University of Latvia (2007), REG: Rhododendron Nursery Babīte, University of Latvia (2012)
Fls 15/slightly domed truss, funnel-shaped, 40 × 50mm, with 5 flat-edged lobes, strong purplish red (60C) in bud, opening inside strong purplish red (60D), outside strong purplish red (63A), with dark red (187A) spots on the dorsal lobes; filaments red-purple, anthers white; style red-purple, stigma black. Calyx 2mm long, light green. Lvs oblanceolate, 140–170 × 55–60mm, acute at apex, obtuse at base, hairless. Shrub 0.6 × 1.0m in 15 years. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: epithet refers to the nursery of origin and its compact habit. Originated in 1997)

(a) **'Baiba'**

cv. Deciduous azalea: Exbury azalea seedling (no. Ex-5) (s) × 'Uguns'
H: R. Kondratovičs (1976), G: R. Kondratovičs (1982), N: R. Kondratovičs (2001), REG: Rhododendron Nursery Babīte, University of Latvia (2012)

Fls single, 10–11/ slightly domed truss, tubular funnel-shaped, with 5–6 wavy-edged lobes, strong red (41B) in bud, opening inside strong reddish orange (32B) and strong orange (25B), outside strong reddish orange (32B), with a vivid orange-yellow (23A) blotch on the dorsal lobes; filaments and anthers yellow; style orange-red, stigma light green. Calyx 2mm long, light green. Lvs elliptic, 100–105 × 40–45mm, pointed at base and apex, margins wavy, glossy green above, hairless. Shrub 3.0 × 2.5m in 20 years. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: a Latvian female christian name)

(r) **‘Beni-hibiki’**

cv. Elepidote rhododendron: *degronianum* subsp. *degronianum* (s) × *arboreum* × *degronianum* subsp. *heptamerum* var. *kyomaruense*
H: H. Suzuki (1992), G: H. Suzuki (2002), N: H. Suzuki (2006), REG: H. Suzuki (2012)
Fls 8–12 (average 10)/ball truss, open campanulate, 60 × 90mm, with 5–7 (average 5) very wavy-edged lobes, red in bud, opening red, with black spots all around the corolla, but most dense in the dorsal throat. Lvs oblong-elliptic, 90 × 40mm, broadly acute at apex, cuneate at base, margins flat, matt green above, with a discontinuous indumentum below which is not visible to the naked eye. Shrub 1.0 × 0.8m in 16 years. Mid May. (Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **‘Berkeley 89’**

cv. Elepidote rhododendron: *auriculatum*, open-pollinated
G: G. Gilberg (2009), N: G. Gilberg (2012), REG: G. Gilberg (2012)
Fls 8/lax truss, narrowly funnel-shaped, 80–90 × 75–80mm, with 7 wavy-edged lobes, moderate yellow-green (137C) in tight bud, opening inside white (whiter than 155D), with a flare of reddish purple markings extending for 10–15mm at the base of the dorsal side of the corolla, outside white (whiter than 155D); stamens 14–16, 38–52mm long, filaments pale yellow-green (4D), anthers near

pale orange-yellow (159C); style 70mm long, pale greenish yellow (1D), stigma brilliant greenish yellow (1B); strongly scented. Calyx 2mm long, brilliant yellow-green (150C). Lvs elliptic, 70–150 × 20–60mm, margins decurved, obtuse at apex, oblique at base, matt and moderate yellow-green (146B) above, hairless. Spreading shrub, 2.8 × 2.4m in 12 years. Late July to early August. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: grown from seed collected in 1989 by the registrant from cultivation in Berkeley, California, USA and a reminder of the significant earthquake experienced there on that day)

(r) **‘Blanzeflor’**

cv. Elepidote rhododendron: *R. × nikomontanum* (s) × *decorum*
G: K.E. Grov (2010), N: P.M. Jørgensen (2012), I: Gimle Planteskule, REG: P.M. Jørgensen (2012)
Fls double, 6–8/flat-domed truss, 35–40 × 30–40mm, with 5–6 wavy-edged, slightly reflexed lobes in outer corolla and with petaloid stamens, creamy white in bud, opening inside creamy white, with a more yellowish tone towards the centre, unmarked, outside creamy white; slightly scented. Calyx 1–2mm long, reddish green, slightly glandular. Lvs elliptic, 40–50 × 20–30mm, cuneate at base, rounded at apex, margins flat, hairless. Shrub 0.5 × 0.5m in 10 years. Late April. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). First described and illustrated in *Lapprosen 2*: 4–6 (2012). Etymology: Blanzeflor is a mythical princess, from a medieval Swedish song, who was fond of white flowers. Seed originally obtained from the Swedish Rhododendron Society: hybridiser unknown)

(r) **‘Bleuñ Skaer’**

cv. Elepidote rhododendron: ‘Double Date’ (s) × *griersonianum*
H: L. Le Duigou (2000), G: A. Le Duigou (2006), N: Scaer Town Council (2012), REG: L. & A. Le Duigou (2012)
Fls 5–7/lax truss, funnel-shaped, 70 × 85mm, with 5 wavy-edged lobes, buds vivid red (46B) and with

white hairs, opening inside deep yellowish pink (46D), with vivid red (46B) towards the base, outside deep yellowish pink (47C); numerous, small, discrete, vivid red (46B) spots on all lobes; stamens usually 10, filaments vivid red (46B); style vivid red (46B), stigma dark red (59A). Calyx 3mm long, strong red (46A), covered with white hairs. Lvs elliptic, 165 × 40mm, margins flat, acute at apex, cuneate at base, matt and moderate olive green (N137B) above, with an indumentum of hairs below (the hairs greenish white, 192D, at first, aging to light orange, 26C). Shrub 1.45 × 1.2m in 12 years. May to June.

(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY). Etymology: the name, chosen by the Town Council of Scaer (Brittany, France), means “flower of Scaer”. The spelling “Skaer” is deliberate and represents the Breton spelling for the town)

(r) **‘Breizh-Izel’**

cv. Elepidote rhododendron: ‘Nancy Evans’ (s) × ‘Fireman Jeff’

H: L. Le Duigou (2000), G: A. Le Duigou (2008), N: L. & A. Le Duigou (2012), REG: L. & A. Le Duigou (2012)

Fls 12–14/rounded truss, open funnel-shaped, 55 × 80mm, with 5–6 wavy-edged lobes, buds vivid red (46B), opening inside light orange (28D) and maturing to light orange (19B), outside light orange (26C), maturing to light orange-yellow (24C); vivid reddish orange (34A) spots on all lobes, but more numerous on dorsal lobes; stamens 11; style brilliant yellow-green (150C), stigma dark greenish yellow (152C). Calyx 35mm long, vivid red (45A), maturing to moderate orange (173C). Lvs elliptic, 75 × 40mm, margins flat, broadly acute at apex, rounded at base, matt green above, hairless. Compact shrub, 1.0m × 1.2m in 12 years. May.

(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY). Etymology: Breizh-Izel is the Breton name for Basse-Bretagne (Lower Brittany), that part of Brittany where the Breton language was traditionally spoken and where the culture associated with this language is most prolific)

(r) **‘Bulduri’**

cv. Elepidote rhododendron: ‘Lavanda’ (s) × ‘Marka’

H: R. Kondratovičs (1985), G: R. Kondratovičs (1991), N: R. Kondratovičs (2010), REG: Rhododendron Nursery Babīte, University of Latvia (2012)

Fls 16/slightly domed truss, open funnel-shaped, 80mm wide, with 5 very wavy-edged lobes, vivid reddish purple (N74B) in bud, opening very pale purple (76C) in the centre, light purple (75A) at the margins, outside vivid reddish purple (N74B), dorsal lobes with strong greenish yellow (151A) spots; filaments white, anthers grey; style white, stigma red. Calyx 2mm long, light green. Lvs oblanceolate, 160–170 × 65mm, pointed at apex, rounded at base, margins flat, glossy green above, with white hairs below when young, which age to brown. Shrub 3.0 × 3.0m in 25 years. Late May to early June.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the Bulduri Gardening School, Latvia)

(r) **‘Burnie Gold’**

cv. Lepidote rhododendron: *chrysodoron* (s) × ‘Chrysomanicum’

H: N. Sullivan (1991), G: Emu Valley Rhododendron Garden (2000), N: N. Sullivan (2000), I: Emu Valley Rhododendron Garden (2011), REG: Emu Valley Rhododendron Garden (2012)

Fls 7/loose truss, broadly campanulate, 35 × 68mm wide, with 5 wavy-edged lobes, buds vivid yellow-green (154A) and contrasting with the reddish-tipped, strongly white ciliate bud scales, opening light greenish yellow (1C) inside and out; dorsal spotting of brilliant greenish yellow (1A); filaments and style greenish yellow; anthers reddish brown. Calyx 8mm long, pale greenish yellow (9D). Lvs elliptic, 65 × 30mm, margins flat, acute at apex, cuneate at base, matt green above, with a scaly indumentum and the margins of the young leaves with long, white, marginal hairs. Shrub 0.8 × 0.8m in 10 years. Early August (Tasmania, Australia). (Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **'Catherine Josephine'**

cv. Elepidote rhododendron: un-named Gable hybrid (no. 41/64) (s) × *glischrum* (from RSF 66/656)
H: H.B. Kline (2004), G: H.B. Kline (2012),
N: H.B. Kline (2012), REG: H.B. Kline (2012)
Fls 12–15/ball truss, open to broadly funnel-shaped, 55 × 70mm, with 5–6 wavy-edged lobes, deep rose pink in bud (the lobe edges with a picotee effect close to light purple, 75A), opening inside slightly off white to very pale purple (close to 75D), with a very dark purple basal blotch; outside similarly coloured, shading lighter towards the base. Calyx 6mm long, light red-purple. Lvs elliptic/lanceolate, 115–150 × 30–50mm, broadly acute at apex, oblique at base, margins downcurved, semi-glossy and mid to dark green above, with distinct veining above and below, somewhat leathery, hairless. Stems and petioles bristly. Shrub 0.8 × 0.7m in 7 years. Late April to early May. (Etymology: named after the late wife of the registrant)

(r) **'Catherine White'**

cv. Elepidote rhododendron: parentage unknown
G: R.H. White (c. 2000), S: R.H. White (c. 1995),
N: R.H. White (2012), REG: J. Delano (2012)
Fls 11/conical truss, broadly funnel-shaped, 63 × 100mm, with 6 very wavy-edged lobes, buds moderate yellowish pink (38B), opening inside cream (pale orange-yellow, 159C), with a pale greenish yellow (2D) throat; outside cream (pale orange-yellow, 159C); dorsal lobes with moderate red (47A) spotting emanating from the throat; stamens cream (pale orange-yellow, 159C), style brilliant yellow-green (150B), stigma strong red (46A). Lvs elliptic, 130 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 2.4 × 2.7m in 17 years. Late (May on Cape Cod, Massachusetts, USA). (Etymology: named for the wife of the nominant)

(r) **'Champagne Lace'**

cv. Elepidote rhododendron: 'Glenna' (s) × 'Janet Blair'
H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2012), REG: J. Barlup (2012)

Fls 19/ball truss, broadly funnel-shaped, 50 × 90mm, with 5 very wavy-edged lobes, strong purplish red (54A) in bud, opening inside to light greenish yellow (4C) in the centre, blending to pale yellow-green (4D) toward the edge, dorsal lobes light greenish yellow (4B) with a moderate yellow (162B) flare; outside overall pale yellow-green (4D), with the base light greenish yellow (4C). Lvs elliptic, 125 × 45mm, broadly acute at the tip, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 7 years. Late. (Etymology: the colour is descriptive of the flower)

(r) **'Charming Pink'**

cv. Elepidote rhododendron: 'Pink Flourish' (s) × 'Cape White'
H: J. Doppel (2004), G: J. Doppel (2008), N: J. Doppel (2011), REG: J. Doppel (2012)
Fls 10/ball truss, funnel-shaped, 45 × 57mm, with 5 very wavy-edged lobes, deep pink (52B) in bud, opening inside moderate purplish pink (68C) at margins, blending into pale purplish pink (56C) towards the centre, with a light yellow (15D) dorsal blotch, outside moderate purplish pink (68C), with pale purplish pink (56C) midribs. Lvs elliptic, 100 × 45mm, broadly acute at apex, rounded at base, with downcurved margins, glossy and mid green above, hairless. Shrub 0.8 × 0.9m in 8 years. Mid May.

(r) **'Cherries and Merlot'**

cv. Elepidote rhododendron: 'Pretty Baby' × *pachysanthum* (s) × 'Whid Bee'
H: F. Fujioka (2001), G: F. Fujioka (2005), N: F. Fujioka (2012), REG: F. Fujioka (2012)
Fls 18/domed truss, funnel-campanulate, 50 × 50mm, with 5 wavy-edged lobes, moderate red (47A) in bud, opening inside vivid red (45A), with two elongated clusters of small, discrete black spots on the dorsal lobe; outside moderate red (47A) to vivid red (46B). Calyx c.1.5mm long, deep purplish red (71A). Lvs elliptic, 90 × 40mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and dark green (136A) above, deep purplish red (close to 71A) below, hairy below (the hairs moderate purplish red (59C) at first, aging to dark red (59A)). Shrub with a dense habit, 0.9 ×

0.8m in 10 years. Late April.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).
Etymology: named in reference to the colour of the flowers and the colour of the indumentum on the lower leaf surface. The 'Pretty Baby' involved in the parentage is an unregistered hybrid derived from *yakushimanum* × 'Cornubia')

(r) '**Cheryl's Choice**'

cv. Elepidote rhododendron: parentage unknown
H: J.C. Cowles (1960s), G: Heritage Museums & Gardens (1970), N: C. Lilly (2011), I: Briarwood Gardens (1988), REG: Heritage Museums & Gardens (2012)
Fls 7–8/lax truss, broadly funnel-shaped, 55 × 90mm, with 6–7 wavy-edged lobes, deep purplish pink (66C) in bud, opening inside very pale purple (75D), with a moderate yellow-green (146C) throat and light purplish pink (63D) margins; outside light purplish pink (63D); stamens large, petaloid, moderate purplish pink (74D). Lvs elliptic, 115 × 50mm, broadly acute at apex, oblique at base, dull and moderate olive green (137A–B) above, hairless. Shrub 1.2 × 1.2m in 10 years.
(Etymology: Cheryl Lilly is the nominant)

(r) '**Dalette Atkins**'

cv. Elepidote rhododendron: 'Janet Blair' (s) × 'Helen Vieira'
H: J.D. Brim (2002), G: J.D. Brim (2005), N: J.D. Brim (2012), REG: J.D. Brim (2012)
Fls 18/ball truss, open funnel-shaped, 57 × 75mm, with 6 wavy-edged lobes, inside pale purplish pink (56D), with strong red (53B) blotches, outside light purplish pink (55C). Lvs oblong, 200 × 64mm, obtuse at apex, cuneate at base, margins downcurved, semi-glossy and moderate olive green (137B) above, hairless. Shrub 1.2 × 1.5m in 10 years. Midseason (May in western North Carolina, USA).
(Etymology: named for the daughter of the hybridiser)

(r) '**Don's Blue Star**'

cv. Lepidote rhododendron: *augustinii*
subsp. *augustinii* (s) × *augustinii*
subsp. *chasmanthum*

G: A. Meerkerk, S: Meerkerk Rhododendron Gardens (2005), N: Meerkerk Rhododendron Gardens (2012), I: Meerkerk Rhododendron Gardens (2012), REG: Meerkerk Rhododendron Gardens (2012)

Fls 3/lax truss, broadly funnel-shaped, 40 × 65mm, with 5 wavy-edged lobes, strong purple (84A) in bud, opening inside with a 6mm wide band of light purple (85A) at the margins, shading to very pale purple (84D) in the centre; with moderate yellow (162B) spots on the dorsal lobe; outside light purple (84C) at the edge, shading to light purple (84D) in the centre. Lvs ovate, 75 × 20mm, acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, scaly. Shrub 2.8 × 1.9m in over 30 years. Early.
(Etymology: named for a friend of Meerkerk Rhododendron Gardens)

(r) '**Eric Szabo**'

cv. Elepidote rhododendron: parentage unknown
H: unknown, G: Emu Valley Rhododendron Garden (2007), N: L. Coombes (2011), I: Emu Valley Rhododendron Garden (2011), REG: Emu Valley Rhododendron Garden (2012)
Fls 13/ball truss, funnel-shaped, 30 × 45mm, with 5 wavy-edged lobes, buds moderate yellowish pink (37C), opening pale yellow (8D) inside and out, with sparse, scattered, purplish pink spots on the dorsal lobes and similar markings and a suffusion around the base of the throat; filaments and style white in upper parts, anthers brownish, stigma green. Lvs elliptic, 95 × 38mm, margins flat, acute at apex, cuneate at base, matt green above, hairless. Shrub 1.0 × 1.0m in 10 years. Late August (Tasmania, Australia).
(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named in memory of a young man who died in a tragic accident)

(r) '**Fryderyk**'

cv. Elepidote rhododendron: 'Belkanto' (s) × *aureum*
H: P. Muras (1996), G: P. Muras (2001), N: P. Muras (2010), REG: Mrs M. Muras (2012)

Fls 10/slightly domed truss, open funnel-campanulate, 35 × 55mm, with 5 wavy-edged lobes, deep pink (48B) in bud, opening pale yellow-green (4D) and tinged light purplish pink (62C), fading to pale yellow-green (4D), outside pale yellow-green (4D), with light purplish pink (62C) stripes on the midveins; conspicuous strong yellow-green (144B) spotting on the dorsal lobe and parts of the adjacent lobes; filaments light to pale purplish pink (62C–D); style deep pink (48B), stigma brown. Calyx 2mm long, reddish. Lvs elliptic, 40–80 × 25–40mm, obtuse at apex, rounded at base, margins flat, slightly glossy and moderate green above, hairless; petioles red on the sunny side. Dense, flat-rounded shrub, 0.5 × 0.8m in 10 years. Late April to early May (in southern Poland). (Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for Fryderyk Chopin, the Polish composer and pianist, the anniversary of whose birth was celebrated in 2010)

(r) **‘George Sand’**

cv. Elepidote rhododendron: ‘Belkanto’ × *aureum*
H: P. Muras (1996), G: P. Muras (2001), N: P. Muras (2011), REG: Mrs M. Muras (2012)
Fls 10–14/slightly domed truss, open funnel-campanulate, 35–40 × 55–60mm, with 5 wavy-edged lobes, strong yellowish pink (37A) in bud, opening light yellow (10C), blending to pale yellow (11D), both inside and out, with brilliant yellow-green (154B) spotting on the dorsal lobe and parts of the adjacent lobes; filaments light yellow (10C); style strong yellowish pink (37A), stigma reddish brown. Lvs elliptic, 40–80 × 25–40mm, obtuse at apex, rounded at base, margins flat, slightly glossy and moderate green above, hairless; petioles red on sunny side. Rounded shrub, with a dense habit, 0.5 × 0.5m in 10 years. Late April to early May (in southern Poland). (Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the French writer who was a close friend of the Polish composer and pianist Fryderyk Chopin)

(r) **‘Golden Rhapsody’**

cv. Elepidote rhododendron: ‘Golden Star’ (s) × ‘Fort Cream’ × *vernicosum*, Rock18139

H: W.L. Rhein, G: W.L. Rhein, N: W.L. Rhein, REG: Rhein Registration Committee (2012)
Fls 8–12/domed truss, broadly funnel-shaped, 50–55 × 90–95mm, with 7 wavy-edged lobes, pale yellow (11C) in bud, opening inside pale greenish yellow (13D), with a small, brilliant yellow (13C) speckled blotch on the dorsal lobe; outside pale yellow (11C); anthers prominent, dark red. Calyx 3mm long, strong yellow-green (143C). Lvs oblong, 115–150 × 65–75mm, broadly acute at apex, rounded at base, with slightly downcurved margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.3m in 10 years. Mid May.

(r) **‘Golly Gumdrops’**

cv. Elepidote rhododendron: ‘Ditto’s Mom’ (s) × ‘Francesca’
H: F.E Burns (1995), G: F.E. Burns (2009), N: F.E. Burns (2011), REG: F.E. Burns (2012)
Fls 15/domed truss, funnel-shaped, 50 × 70mm, with 5 very wavy-edged lobes, vivid red (52A) in bud, opening inside deep pink (52B) at margins and all except the dorsal lobe changing abruptly to strong pink (52D) and fading to light pink (50D) down to the centre; the dorsal lobe fades quickly to almost creamy white; on the outside the lobes are strong red (51A) at the apex, fading somewhat to the edges, with a broad strong red (51A) stripe running down to the base; internally five strong pink (52D) rays rise upwards around the nine pale pink stamens. Calyx 3mm long, moderate yellow-green (147C). Lvs elliptic, 115 × 45mm, margins flat, broadly acute at apex, oblique at base, dull and moderate olive green (147A) above; with light olive brown (199B) hairs, most numerous on the petioles and down the midvein. Shrub 1.5 × 0.9m in 15 years. Late May to early June. (Etymology: slang expression used by the registrant, who suggests that ‘the colourful trusses stand up on the plant like big bright pink gumdrops’)

(r) **‘Hachgold’**

cv. Elepidote rhododendron: ‘Bernadette’ (s) × ‘Goldzauber’
H: H. Hachmann (1994), G: H. Hachmann (1999), N: Holger Hachmann, I: Holger Hachmann (2009), REG: M. Oppenrieder (2012)

Fls 7–12/flat rounded, slightly loose truss, campanulate, 50–72mm wide, with 5 flat- to slightly wavy-edged lobes, buds bright red, with some patches of yellow shading, opening inside vivid yellow (12A), outside yellow, with reddish areas near the margin; sometimes with small, delicate reddish markings at the base of the three dorsal lobes; style and filaments short; stigma green. Lvs elliptic, margins flat, glossy above, hairless. Shrub 0.8 × 0.9m in 10 years. Mid May to late June.

Syn: 'Marathon Gold', 'Ines'. Trade designation: GOLDEN EVEREST

(r) '**Haru-no-yuki**'

cv. Elepidote rhododendron: selection from *makinoi* (s)

G: S. Ijima (2005), N: S. Ijima (2008), REG: S. Ijima (2012)

Fls 6–10 (average 8)/loose ball truss, open funnel-shaped, 50 × 50mm, with 5 slightly wavy-edged and rather narrow lobes (making rather starry flowers), white in bud, opening white inside and out, with no markings; filaments and style white; anthers and stigma pale greenish. Lvs narrowly oblanceolate, 100 × 20mm, acute at apex, cuneate at base, matt green above, with a dense white indumentum at first below which ages to brown. Shrub 0.6 × 0.4m in 15 years. Mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Grown from seed sown by the registrant in 1996)

(v) '**Hendrik's Kers**'

cv. Lepidote rhododendron: 'Hot Tropic' × *saxifragoides* (s) × 'Thai Gold'

H: H. Van Oost (2002), G: H. Van Oost (2005), N: H. Van Oost (2012), I: H. Van Oost, REG: H. Van Oost (2012)

Fls 13–29/compact, rounded truss, tubular funnel-shaped, 44 × 43mm, with 5 flat-edged lobes, red in bud, opening red throughout (the colour changing with light intensity from dark red to dark orange); unmarked; stamens 10, anthers very dark, filaments red. Pedicels brownish red. Lvs elliptic, 50–60 × 20–23mm, margins flat, acute at apex,

cuneate at base, glossy green above, with a scaly indumentum: scales silvery at first, aging to brown. Shrub 0.6 × 0.6m in 5 years. Spring.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: derived from the raiser's first name and a colour description of the flowers ("kers" meaning cherry red))

(r) '**Hinamatsuri**'

cv. Elepidote rhododendron: 'Double Date' (s) × *degronianum* subsp. *heptamerum* (with double white flowers) × *makinoi* (with double pink flowers)

H: T. Takahashi (2000), G: T. Takahashi (2004), N: T. Takahashi (2005), REG: T. Takahashi (2012)

Fls double, 12–14 (usually 12)/ball truss, funnel-shaped, 45 × 60mm, with 16–21 lobes (usually 16), buds pink, opening inside deep purplish pink (68A), outside strong purplish pink (68B), unmarked.

Calyx 1mm long, bright green. Lvs oblong-lanceolate, 115 × 35mm, acuminate at apex, cuneate at base, margins upcurved, with sparse, white hairs when young, which age to brown. Shrub 0.6 × 0.5m in 11 years. Early April.

(Nomenclatural standard: colour print provided by the registrant (WSY))

(r) '**Howie**'

cv. Elepidote rhododendron: parentage unknown

H: N. Sullivan (1991), G: Emu Valley Rhododendron Garden (2000), N: J. Woods (2011), I: Emu Valley Rhododendron Garden (2011), REG: Emu Valley Rhododendron Garden (2012)

Fls 17/ball truss, funnel-campanulate, 40 × 40mm, with 5 wavy-edged lobes, buds strong red (46A), opening strong red (46A) throughout, with blackish spotting over the three dorsal lobes. Lvs elliptic, 110 × 38mm, margins flat, acute at apex, cuneate at base, matt green above, with fawn-coloured hairs below when mature. Shrub 0.8 × 1.0m in 10 years. Late August (Tasmania, Australia).

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a volunteer at the Emu Valley Rhododendron Garden)

(r) **'Inner Glow'**

cv. Elepidote rhododendron: parentage unknown
H: J. Doppel (1996), G: J. Doppel (2000), N: J. Doppel (2012), REG: J. Doppel (2012)
Fls 9/conical truss, funnel-shaped, 48 × 75mm, with 5 flat-edged lobes, vivid red (44A) in bud, opening inside pale yellow-green (4D) at the base, shading to strong purplish red (58B) at the margins, with brilliant yellow (20A) dorsal spotting; outside strong purplish red (58B). Lvs elliptic, 100 × 40mm, rounded at base, flat at margins, dull and mid green above, hairless. Shrub 1.0 × 1.2m in 16 years. Mid May.

(r) **'Jānis Sproģis'**

cv. Elepidote rhododendron: 'Nova Zembla' (s) × 'Panenka'
H: R. Kondratovičs (1985), G: R. Kondratovičs (1991), N: R. Kondratovičs (2005),
I: Rhododendron Nursery Babīte, University of Latvia (2005), REG: Rhododendron Nursery Babīte, University of Latvia (2012)
Fls 13/conical truss, open funnel-shaped, 80–85mm wide, with 5 wavy-edged lobes, buds vivid purplish red (61C), opening inside light purplish pink (73C), with broad, strong purplish pink (73B) margins, outside deep purplish pink (73A); strong greenish yellow (153C) spots on the dorsal lobes; filaments white, anthers yellow; style purple, stigma dark brown. Calyx 1mm long, very light green. Lvs elliptic, 155–160 × 60–65mm, obtuse at base and apex, margins flat, matt green above, hairless. Shrub 2.3 × 3.0m in 20 years. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a famous Latvian singer)

(r) **'Jenny Cox'**

cv. Lepidote rhododendron: *moupinense*, red form × *johnstoneanum* (s) × *moupinense*, red form
H: M. Kupsch (1999), G: Emu Valley Rhododendron Garden (2004), N: M. Kupsch (2010), I: Emu Valley Rhododendron Garden (2011), REG: Emu Valley Rhododendron Garden (2012)
Fls 3/loose truss, open funnel-shaped, 50 × 72mm, with 5 wavy-edged lobes, buds strong red (53D), opening inside light purplish pink (62C), outside

strong purplish pink (62A); with numerous, strong purplish red (61B) spots on the dorsal lobe. Calyx 3mm long, brilliant yellow-green (149B). Lvs elliptic, 60 × 22mm, margins flat, broadly acute to obtuse at apex, cuneate at base, matt green above, with a scaly indumentum. Shrub 0.8 × 0.7m in 10 years. Late August (Tasmania, Australia). (Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a friend of the Emu Valley Rhododendron Garden)

(r) **'Joan Marie'**

cv. Elepidote rhododendron: parentage unknown
H: J.C. Cowles (1960s), G: Heritage Museums & Gardens (c. 1970), N: J.M. O'Connor (2011), I: Briarwood Gardens (c. 1988), REG: Heritage Museums & Gardens (2012)
Fls 14/ball truss, broadly funnel-shaped, 60 × 100mm, with 6 wavy-edged lobes, strong purplish red (60C–D) in bud, opening inside light reddish purple (74C) to moderate purplish pink (74D), unmarked; outside strong purplish red (71C–D). Lvs elliptic, 115 × 45mm, broadly acute at apex, oblique at base, with wavy margins, dull and moderate olive green (147A) to moderate yellow-green (147B) above, hairless. Shrub 2.5 × 1.8m in 20 years. Mid May. (Etymology: Joan Marie O'Connor is the nominant)

(r) **'Joe's Moon Glow'**

cv. Elepidote rhododendron: 'Capistrano' (s) × 'Early Light'
H: J. Doppel (2001), G: J. Doppel (2005), N: J. Doppel (2012), REG: J. Doppel (2012)
Fls 10–12/domed truss, broadly funnel-shaped, 40 × 72mm, with 6 wavy-edged lobes, strong yellowish pink (37A) in bud, opening throughout light greenish yellow (8C), unmarked. Lvs elliptic, 125 × 50mm, obtuse at apex, rounded at base, margins flat, semi-glossy and mid green above, hairless. Shrub 0.75 × 1.2m in 11 years. Early May.

(r) **'Joey Rabbit'**

cv. Elepidote rhododendron: parentage unknown
H: unknown, G: Emu Valley Rhododendron

Garden (2007), N: J. Wood (2011), I: Emu Valley Rhododendron Garden (2011), REG: Emu Valley Rhododendron Garden (2012)
Fls 20/ball truss, broadly-campanulate, 38 × 48mm, with 5 wavy-edged lobes, buds strong red (51A), opening strong red (53C) throughout; unmarked; bracts very pale, strongly hairy, conspicuous as the inflorescence is opening. Lvs elliptic, 85 × 32mm, margins flat, acute at apex, cuneate to obtuse at base, matt green above, with pale yellow hairs (158A when young, aging to 158B). Shrub 1.8 × 1.0m in 10 years. Late August (Tasmania, Australia).
(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named in memory of a small boy, a cousin of the nominant Juanita Wood)

(r) **'John's Lavender Haze'**

cv. Elepidote rhododendron: 'Minnetonka' (s) × 'Smokey'
H: J. Doppel (2006), G: J. Doppel (2010), N: J. Doppel (2012), REG: J. Doppel (2012)
Fls 8/ball truss, broadly funnel-shaped, 40 × 65mm, with 5 wavy-edged lobes, strong purplish red (71B) in bud, opening inside very light purple (76B), with a solid pale yellow-green (149D) blotch on the dorsal lobe; outside light reddish purple (74C). Lvs elliptic, 100 × 40mm, broadly acute at apex, rounded at base, margins downcurved, glossy and olive green above, hairless. Shrub with a dense habit, 0.5 × 0.6m in 5 years. Mid May.

(r) **'Junkō-musume'**

cv. Elepidote rhododendron: *makinoi* (white-flowered) (s) × 'Hotei'
H: H. Suzuki (1995), G: H. Suzuki (2003), N: H. Suzuki (2008), REG: H. Suzuki (2012)
Fls 13–18 (average 15)/hemispherical truss, open campanulate, 40 × 40mm, with 5(–6) slightly wavy-edged lobes, deep yellow in bud, opening light yellow throughout, with yellowish dorsal spotting. A conspicuous orange bract at the base of each pedicel. Lvs narrowly elliptic to oblong-elliptic, 80 × 20mm, margins flat to slightly recurved, acute to broadly acute at apex, cuneate at base, with hairs below white when young, aging to brown. Shrub 50

× 30cm, in 6 years from grafting. Mid May.
(Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **'Kiden'**

cv. Elepidote rhododendron: *makinoi* (white-flowered) (s) × 'Hotei' × 'Lem's Cameo'
H: H. Suzuki (1995), G: H. Suzuki (2003), N: H. Suzuki (2008), REG: H. Suzuki (2012)
Fls 9–13 (average 11)/loose truss, open campanulate, 40 × 40mm, with 5(–6) slightly wavy-edged lobes, buds orange, tipped yellow, opening pale yellow throughout, with inconspicuous greenish dorsal spotting. Lvs narrowly elliptic, 80 × 20mm, margins flat to recurved, cuneate at base, acute at apex, slightly glossy green above, with a white indumentum below when young which ages to brown. Shrub 50 × 40cm, in 6 years after grafting. Mid May.
(Nomenclatural standard: colour print provided by the registrant (WSY))

(v) **'Lab Retriever Webster'**

cv. Lepidote rhododendron: *helligii* hybrid (s) × ('Doctor Sax' × *aurigeranum*) × *leucogigas*
H: P. Sullivan, G: P. Sullivan, S: Pacific Island Nursery (2009), N: R. Kruppa (2009), I: Pacific Island Nursery, REG: R. Kruppa (2012)
Fls 7–9/conical truss, tubular funnel-shaped, 95 × 95mm, with 7 wavy-edged lobes, purple in bud, opening inside vivid red (46C), with a dark red, narrow vein on each lobe, extending from base to tip; outside vivid red (46C); anthers red, with a yellow tip, filaments red; style pink, stigma purple. No calyx apparent. Lvs elliptic, 135–140 × 55–60mm, glossy and dark yellowish green (139A) above, with no evident indumentum. Shrub 1.1 × 0.8m in 9 years. May to June and November to December.

(r) **'Lavender Moonlight'**

cv. Elepidote rhododendron: 'Purple Amethyst' (s) × 'Plum Passion'
H: J. Barlup (2002), G: J. Barlup (2006), N: J. Barlup (2011), REG: J. Barlup (2012)
Fls 13/conical truss, broadly funnel-shaped, 45 × 75mm, with 5 very wavy-edged lobes, dark red (59A)

in bud, opening inside light purple (75B), edged strong purple (77B), with a white (155D) dorsal flare having moderate yellow-green (138B) spots and extending 30mm from the base; outside light purple (77D), the edge and midribs strong purple (77B); not scented. Lvs elliptic, 120 × 45mm, margins upcurved, broadly acute at apex, rounded at base, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.8m in 9 years. Late May.

(r) **‘Lemon Passion’**

cv. Elepidote rhododendron: ‘Tisa’ (s) × ‘Berg’s Yellow’ × *proteoides*

H: J. Barlup (2004), G: J. Barlup (2009), N: J. Barlup (2011), REG: J. Barlup (2012)

Fls 16/domed truss, funnel-shaped, 45 × 45mm, with 5 wavy-edged lobes, deep yellowish pink (39B) in bud, opening inside pale yellow (11C–D) with a flare of discrete, deep red (185A) spots extending upwards c.35mm from the base of the dorsal lobe; outside pale yellow (11C); not scented. Calyx c.10mm long, pale yellow (11C). Lvs elliptic, c.80 × 40mm, margins flat, broadly acute at apex, rounded at base, dull and moderate olive green (147A) above, hairless. Shrub with a dense habit, 0.6 × 0.7m in 7 years. Late April.

(Etymology: name suggested by the flower colour, which resembles that of a ripe lemon)

(r) **‘Lil’ Darlin’**

cv. Elepidote rhododendron: ‘Capistrano’ (s) × ‘September Song’

H: J. Doppel (2005), G: J. Doppel (2007), N: J. Doppel (2012), REG: J. Doppel (2012)

Fls 10/domed truss, broadly funnel-shaped, 48 × 50mm, with 6 flat-edged lobes, strong red (41B) in bud, opening inside light orange-yellow (23C), with dark greenish yellow (152D) speckles on the dorsal lobe; outside deep yellowish pink (39B), changing to light yellowish pink (26D) at the base. Calyx 25mm long, light yellowish pink (26D), shading to light yellowish pink (38D). Lvs elliptic, 100 × 55mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and olive green above, hairless. Dense shrub, 0.4 × 0.75m in 7 years. Early May.

(r) **‘Maddie Mae’**

cv. Elepidote rhododendron: ‘Nancy Evans’ (s) × ‘Pink Petticoats’

H: J. Barlup (1989), G: J. Barlup (1994), N: J. Barlup (2011), REG: J. Barlup (2012)

Fls 16/ball truss, funnel-shaped, 50 × 55mm, with 5 very wavy-edged lobes, strong purplish red (63A) in bud, opening inside pale yellowish pink (159D), with a pale yellow-green (4D) dorsal flare speckled with strong red (153D) spots and extending 40mm from the base; nectar pouches strong red (53B), extending c.6mm from the base; outside pale yellowish pink (159D), the midribs pale purplish pink (65D). Calyx c.20mm long, pale yellow-green (4D), with strong red (53B) flares. Lvs elliptic, 105 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.4 × 2.8m in 22 years. Mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the dog belonging to a friend of the registrant)

(r) **‘Makayla Quinn’**

cv. Elepidote rhododendron: ‘Janet Blair’ (s) × ‘Helen Vieira’

H: J.D. Brim (2002), G: J.D. Brim (2005), N: J.D. Brim (2012), REG: J.D. Brim (2012)

Fls 20/ball truss, open funnel-shaped, 57 × 100mm, with 6 wavy-edged lobes, inside moderate purplish pink (54D), with a deep red (60A) blotch, outside strong pink (54C). Lvs oblong, 150 × 50mm, acute at apex, cuneate at base, margins flat, glossy and dark yellowish green (139A) above, hairless. Shrub 1.2 × 1.7m in 10 years. Midseason.

(Etymology: named for a granddaughter of the hybridiser)

(r) **‘Manamusume’**

cv. Elepidote rhododendron: *degronianum*

subsp. *degronianum* (red-flowered) (s) × ‘America’

H: H. Suzuki (1998), G: H. Suzuki (2008), N: H. Suzuki (2008), REG: H. Suzuki (2012)

Fls 9–13 (average 12)/hemispherical truss, open funnel-shaped, 50 × 50mm, with 5(–6) slightly

wavy-edged lobes, deep red in bud, opening red throughout, with inconspicuous red dorsal spotting. Lvs narrowly elliptic, 100 × 30mm, margins flat, cuneate at base, acute at apex, glossy green above, with few hairs below. Shrub 50 × 40cm, in 5 years after grafting. Early May.

(Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **'Mardi'**

cv. Elepidote rhododendron: 'Coral Blossom' (s) × 'Bambi' × *proteoides*

H: J. Barlup (2002), G: J. Barlup (2008), N: J. Barlup (2009), REG: J. Barlup (2012)

Fls 13/flat truss, broadly funnel-shaped, 45 × 65mm, with 6 wavy-edged lobes, vivid reddish orange (44C) in bud, opening inside pale yellow (20D), the edges and mid veins deep pink (48A), with the base solid deep red (53A) and extending upwards c.5mm, and with spotted flares of the same colour extending approximately an additional 15mm, mainly on the dorsal side; outside pale yellow (20D), with edges and midveins deep pink (48A). Calyx c.15mm long, pale yellow (20D), with deep red (53A) flares. Lvs elliptic, c.95 × 30mm, margins flat, broadly acute at apex, rounded at base, dull and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 9 years. Early June.

(Etymology: named for a friend of the registrant)

(a) **'Mary Jane Cummings'**

cv. Evergreen azalea: 'Brenda Marie' (s) × 'Festive' × 'Satellite'

H: J.F. Klimavicz (2005), G: J.F. Klimavicz (2007), N: J.F. Klimavicz (2012), REG: J.F. Klimavicz (2012)
Fls double, 2/flat truss, broadly funnel-shaped, 35 × 65mm, with 10 wavy-edged lobes, buds strong reddish purple (78B); opening inside white, with strong reddish purple (78B) and light purple (78C) stripes, sectors and occasional solids, with a slight vivid reddish purple (78A) blotch; outside strong reddish purple (78B) and light purple (78C), fading towards a white base. Calyx 10mm long, light yellow-green (145C). Lvs elliptic, 45 × 20mm, broadly acute at apex, cuneate at base, glossy and moderate olive green (146A) above, hairless. Upright

shrub, 0.6 × 0.5m in 5 years. Early April.

(The 'Satellite' used as a parent was the USDA, Glenn Dale plant. Etymology: the selection of the name was auctioned to benefit the National Institute of Health's Children's Inn (Bethesda, Maryland, USA))

(r) **'Mary S. Phipps'**

cv. Elepidote rhododendron: 'Scintillation' × ('Pink Petticoats' × *yakushimanum*) (s) × ('Dumper's Yellow' × 'Phipps Yellow') × 'One Thousand Butterflies'

H: G. Woodard (2003), G: G. Woodard (2005), N: G. Woodard (2012), REG: G. Woodard (2012)

Fls 23/conical truss, broadly funnel-shaped, 100 × 150mm, with 7 very wavy-edged lobes, strong purplish pink (63C) in bud, opening light purplish pink (62C), with a light yellow, basal blotch. Lvs elliptic, 175 × 65mm, obtuse at apex, rounded at base, with flat margins, dull and light green above, hairless. Dense shrub, 1.5 × 1.5m in 9 years. Mid May.

(Etymology: named for the owner of the Phipps Estate (New York, USA))

(r) **'Mikawa-botan'**

cv. Elepidote rhododendron: *hyperythrum* × *aureum* (s) × *degronianum* subsp. *degronianum*

H: Y. Shibata (1999), G: Y. Shibata (2008), N: Y. Shibata (2009), REG: Y. Shibata (2012)

Fls fully double, 10–15 (average 12)/hemispherical truss, open funnel-shaped, 50 × 60mm, with 15–20 (average 18), very wavy-edged lobes, deep pink in bud, opening deep pink inside and out, with red dorsal spotting. Lvs narrowly elliptic, 130 × 45mm, margins flat, acute at apex, cuneate at base, glossy green above, very sparsely hairy. Shrub 30 × 40cm, 3 years after grafting. Early April.

(Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **'Mine-no-hohoemi'**

cv. Elepidote rhododendron: 'Hotei' (s) × 'Nancy Evans'

H: H. Suzuki (1993), G: H. Suzuki (2006), N: H. Suzuki (2006), REG: H. Suzuki (2012)

Fls 7–12 (average 10)/halfball truss, funnel-campanulate, 50 × 70mm, with 5–7 (usually 5) wavy-edged lobes, orange in bud, opening outside flushed pale orange on a yellow ground, inside with a faint flush of orange around the margins of the lobes, otherwise yellow; dorsal lobes with red spots; filaments and style also pale yellow. Lvs obovate to broadly elliptic, 90 × 40mm, obtuse and apiculate at apex, cuneate at base. Shrub 0.8 × 0.8m in 16 years. Mid May. (Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **'Misty Memory'**

cv. *Elepidote rhododendron*: 'Casanova' (s) × 'Goldsworth Orange'
H: C. Trautmann (1994), G: J. Doppel (1995), N: J. Doppel (2011), REG: J. Doppel (2012)
Fls 7/domed truss, broadly funnel-shaped, 40 × 70mm, with 5 flat-edged lobes, deep pink (52B) in bud, opening inside pale purplish pink (56A), fading to pale yellow-green (4D), with strong red (53C) dorsal spotting, which fades to light yellow (160B); outside deep purplish pink (68C) and pale purplish pink (56B); not scented. Calyx insignificant. Lvs elliptic, 115 × 40mm, margins downcurved, broadly acute at apex, rounded at base, semi-glossy and olive green above; hairless. Shrub with a dense habit, 0.9 × 1.5m in 15 years. Early May.

(r) **'Mone-no-mai'**

cv. *Elepidote rhododendron*: 'Hallelujah', open-pollinated (s)
H: H. Suzuki (1995), G: H. Suzuki (2005), N: H. Suzuki (2005), REG: H. Suzuki (2012)
Fls 9–14 (average 11)/truss, open funnel-shaped, 50 × 70mm, with 5(–7) noticeably wavy-edged lobes, deep red in bud, opening red, with a white throat and white along centre line of the lobes especially at the base, the colour paling considerably with age; small, solid red blotch at the base of the dorsal throat; filaments and style white; anthers dark red; stigma pale green. Lvs ovate to broadly elliptic, 140 × 60mm, broadly acute at apex, cuneate at base, hairless. Shrub 1.0 × 0.8m in 16 years. Mid May. (Nomenclatural standard: colour print provided by the registrant (WSY))

(v) **'Mrs Frances Underwood'**

cv. *Lepidote rhododendron*: *konori* (s) × unknown
H: H.R. Malone (1989), G: Emu Valley Rhododendron Garden (2007), N: Emu Valley Rhododendron Garden (2011), I: Emu Valley Rhododendron Garden (2012), REG: Emu Valley Rhododendron Garden (2012)
Fls 9/flat-topped truss, tubular funnel-shaped, 90 × 100mm, with 7 wavy-edged lobes, inside white (155C), with pink tinges at the base of the lobe margins, outside white (155A), with tinges of pink on the tube; unmarked; anthers long, pale beige; strongly scented. Lvs oblanceolate, 140 × 75mm, margins wavy, obtuse at apex, matt green above, with brown scales below. Shrub 1.0 × 1.0m in 10 years. Spring. (Nomenclatural standard: colour print provided by the registrant (WSY))

(a) **'My Li'l Tonto'**

cv. Evergreen azalea: 'Christina' (s) × 'Shin'nyo-no-tsuki'
H: W.L. Clagett (1989), G: W.L. Clagett (1994), N: W.L. Clagett (2012), I: White's Nursery (2012), REG: W.L. Clagett (2012)
Fls double and hose-in-hose, 3/flat truss, broadly funnel-shaped, 25 × 50mm, with 17–18 very wavy-edged lobes, vivid red (44A), with deep yellowish pink (39B) midribs in bud, opening inside vivid red (44A), with varying amounts of deep yellowish pink (39B); outside vivid red (44A), with deep yellowish pink (39B) midribs. Calyx present, but insignificant. Lvs elliptic, 18 × 6mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (147A) to moderate yellow-green (147C) above, slightly hairy below (hairs silvery at first, aging brownish). Shrub 0.36 × 0.6m in 20 years. Late May. (Etymology: named for a grandchild of the raiser. The 'Christina' parent is the Vuyk van Nes plant derived from 'Florida' × 'Louise Gable')

(r) **'Noce Bretonne'**

cv. *Elepidote rhododendron*: 'Halopeanum' (s) × 'Coronation Day'
H: J.F. Saint-Jalm (1996), G: J.F. Saint-Jalm (2003),

N: J.F. Saint-Jalm (2012), REG: J.F. Saint-Jalm (2012)

Fls 12–14/ball to conical truss, broadly funnel-shaped, 65 × 110mm, with 5–8 wavy-edged lobes, vivid purplish red (61C) in bud, opening inside strong purplish pink (68B), with slightly paler mottling and fading to very pale purple (69B), with a dark ring around the very base of the corolla, outside deep purplish pink (N66C) and fading to very pale purple (69A); dorsal lobe with deep purplish red (59B) spots; stamens 12–14, filaments white; style white and glandular below, stigma deep purplish red (59B); ovary green, with white glands. Calyx 3mm long, deep purplish pink (N66C). Lvs elliptic, 130 × 45mm, margins flat, broadly acute at apex, rounded at base, matt green above, hairless. Shrub 3.2 × 2.4m in 16 years. Mid April to early May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named to celebrate the wedding of the registrant's eldest daughter)

(r) **'Noiret'**

cv. Elepidote rhododendron: 'Francesca' (s) × 'Purple Splendour'

H: F.E. Burns (1995), G: F.E. Burns (2008), N: F.E. Burns (2011), REG: F.E. Burns (2012)

Fls 20/domed truss, 65 × 85mm, open funnel-shaped, with 6 very wavy-edged lobes, deep purplish red (61A) in bud, opening inside light purplish pink (65B), with a prominent, spotted, dark red (183A or darker) blotch on two dorsal lobes, and a few scattered spots of the same colour on adjacent lobes, and with a 10mm long, strong purplish red (57A) ray at the base of the other lobes; outside moderate purplish red (64A); stamens 10, filaments strong purplish pink (62A), anthers white; style curved, light purplish pink (65B), stigma deep purplish red (59B). Calyx variable in length, c.2–c.15mm long, pale purplish pink (62D). Lvs elliptic, 160 × 70mm, margins downcurved, acute at apex, oblique at base, dull and moderate olive green (147A) above, hairless. Shrub 1.2 × 0.9m in 15 years. Late May to early June.

(Etymology: Noiret is an American-bred red wine grape)

(v) **'Oogappel'**

cv. Lepidote rhododendron: (*macgregoriae* × *laetum*) × *saxifragoides* hybrid no.2 from Bertelmann (s) × *saxifragoides* hybrid no.2 from Bertelmann

H: H. Van Oost (2002), G: H. Van Oost (2005), N: H. Van Oost (2012), I: H. Van Oost, REG: H. Van Oost (2012)

Fls 4–8/truss, out-facing to pendent, open funnel-shaped, 84 × 110mm, with 5 flat-edged lobes, lobes inside salmon orange rose, shading to yellow in the throat, outside with a yellow tube and lobe bases, shading to yellow salmon rose on the margins of the lobes; unmarked; stamens 10, anthers dark, filaments salmon rose in upper part, yellow towards the base; style pale green, stigma slightly darker green. Pedicels reddish brown. Lvs elliptic, 100 × 40mm, margins flat, acute at apex, cuneate at base, glossy green above, with a scaly indumentum, scales silver when young, aging to brown. Shrub 0.5 × 0.5m in 5 years. Spring (and sometimes in autumn and winter).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the epithet means "eyeball" in Dutch but is also used, as here, to express admiration in the sense of "favourite" or "darling")

(r) **'Penheale Spring'**

cv. Lepidote rhododendron: *edgeworthii* (s) × *moupinense*

H: J.A.R. Moffat (1975), G: J.A.R. Moffat (1979), N: Mr & Mrs J. Colville (2012), REG: Mr & Mrs J. Colville (2012)

Fls 2–6/lax truss, funnel-campanulate, 45–65 × 55–65mm, with 5 wavy-edged lobes, reddish pink in bud, opening white flushed pink inside and usually with dark pink spots below the dorsal lobe (these showing though to the outside); outside white, often flushed pink and with pink stripes down the midveins; filaments white, anthers dark brown; style white, stigma light brown. Calyx 15mm long, flushed pale pink. Lvs elliptic, 70 × 25mm, acute and apiculate at apex, mostly cuneate at base, margins decurved, matt green above and with slightly impressed veins, the scales below light green at first, aging to green. Shrub 1.8 × 1.8m in 25 years. April to May.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: raised at Penheale Manor)

(r) **'Penn-ar-Bed'**

cv. Elepidote rhododendron: 'Halopeanum' (s) × 'Fred Wynniatt' (FCC 1980)
H: L. Le Duigou (2003), G: A. Le Duigou (2011), N: L. & A. Le Duigou (2012), REG: L. & A. Le Duigou (2012)
Fls 13/conical truss, very broadly funnel-shaped, 70 × 125mm, with 7 flat-edged, recurved lobes, buds vivid purplish red (61C), opening inside very pale purple (69C), with light purplish pink (65B) margins; outside pale purplish pink (65D), with light purplish pink (65B) margins; solid dark red (185A) around the whole of the base of the throat, breaking up into spots around its edges; stamens 12–14; style light yellow-green (145C), stigma strong yellow-green (N144C). Calyx 5mm long, moderate red (185B). Lvs oblong, 200 × 60mm, margins flat, broadly acute at apex, rounded at base, glossy green above, bronzed when young, hairless; petioles moderate red (185B). Shrub 1.75 × 1.2m in 8 years. May.

(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY). Etymology: named for one of the four departments of Brittany – Penn-ar-Bed being the Breton name for Finistère)

(r) **'Pilkingtons' Pride'**

cv. Elepidote rhododendron: parentage unknown
H: A. Consolini (c. 1965), G: A. Consolini, N: N. & J. Beaudry (2012), I: Sandwich Club (2006), REG: Sandwich Club (2012)
Fls 9/lax truss, open funnel-shaped, 90 × 115mm, with 6 very wavy-edged lobes, strong yellowish pink (37B) in bud, opening to creamy white (pale yellowish pink, 159D) inside, with a moderate yellow (160A) flare and some spotting on the dorsal lobe and a partial flare on the adjacent lobes, outside pale yellowish pink (27D); stigma strong yellow-green (145A). Lvs oblong, 140 × 50mm, broadly acute at apex, oblique at base, margins flat, moderate olive green (147A) above, hairless. Shrub 5.0 × 4.5m

in 47 years. Midseason.

(Etymology: named for Harold and Eveline Pilkington, who grew the plant from c.1970)

(r) **'Pink Pirouette'**

cv. Elepidote rhododendron: 'Pink Flourish' (s) × 'Princess Mary of Cambridge'
H: J. Doppel (2003), G: J. Doppel (2007), N: J. Doppel (2011), REG: J. Doppel (2012)
Fls 12/ball truss, broadly funnel-shaped, 50 × 65mm with 5 wavy-edged lobes, moderate purplish red (57A) in bud, opening inside deep purplish pink (68A), with a light yellowish brown (199C) dorsal blotch; the central area on each lobe moderate purplish pink (68C), giving a star-like appearance; outside deep purplish pink (68A). Lvs elliptic, 115 × 48mm, broadly acute at apex, rounded at base, with downcurved margins, semi-glossy and mid green above, hairless. Shrub 0.6 × 0.9m in 9 years.

(r) **'Plum Magic'**

cv. Elepidote rhododendron: 'Plum High' (s) × 'Violetta's Song'
H: J. Barlup (1989), G: J. Barlup (2004), N: J. Barlup (2011), I: Hazelwood Gardens (2010), REG: J. Barlup (2012)
Fls 23/ball truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, strong purplish red (72A) in bud, opening inside very pale purple (76C) at centre, blending into light purple (77D) and edged light purple (77C), with a strong yellow (153D) dorsal flare extending 40mm from the base; outside light purple (77C & 77D), the midribs strong purple (77B); not scented. Lvs elliptic, 125 × 50mm, margins flat, broadly acute at apex, rounded at base, dull and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.9m in 6 years. Late May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY))

(r) **'Princess-kei'**

cv. Elepidote rhododendron: selection derived from *yakushimanum* 'Koichiro Wada' (s)
H: H. Suzuki (1994), G: H. Suzuki (2003), N: H. Suzuki (2007), REG: H. Suzuki (2012)
Fls 7–14 (average 10)/hemispherical truss, funnel-

shaped, 50 × 70mm, with 5–7 (usually 5) very wavy-edged lobes, buds pink, opening white inside and out, with yellowish green dorsal spotting; filaments white; style flushed red especially towards the base, stigma red. Lvs elliptic, 80 × 30mm, broadly acute at apex, cuneate at base, margins flat, glossy green above, with grey hairs below. Shrub 0.8 × 0.5m in 13 years. Mid May.

(Nomenclatural standard: colour print provided by the registrant (WSY))

(r) **‘Purple Cape’**

cv. Elepidote rhododendron: ‘Sefton’, open-pollinated (s)

H: J. Doppel (2000), G: J. Doppel (2004), N: J. Doppel (2008), REG: J. Doppel (2012)

Fls 15/conical truss, broadly funnel-shaped, 50 × 75mm, with 5 flat-edged lobes, strong reddish purple (72B) in bud, opening inside strong purple (77B), with strong reddish orange (34B) dorsal spotting, outside strong purple (77B), with vivid reddish purple (74B) midribs. Lvs elliptic, 100 × 40mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and mid green above, hairless. Shrub 1.8 × 1.8m in 12 years. Mid May.

(r) **‘Quiet Elegance’**

cv. Elepidote rhododendron: ‘Sun Blush’ (s) × ‘Tia’

H: J. Barlup (2003), G: J. Barlup (2007), N: J. Barlup (2011), REG: J. Barlup (2012)

Fls 21/ball truss, broadly funnel-shaped, 45 × 70mm, with 6 very wavy-edged lobes, strong purplish red (67A) in bud, opening inside yellowish white (158C), shading to light purplish pink (65B) edges, with deep red (185A) speckling on the dorsal lobe and a six-pointed star of the same colour at the base; outside similar to inside, but unmarked; not scented. Calyx insignificant. Lvs elliptic, 105 × 50mm, margins flat, broadly acute at apex, rounded at base, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.7m in 5 years. Late April.

(a) **‘Rebecca Taffet’**

cv. Evergreen azalea: ‘Elsie Lee’ (s) × ‘Satellite’ (USDA, Glenn Dale)

H: J.F. Klimavicz (1990), G: J.F. Klimavicz (1992), N: J.F. Klimavicz (2012), REG: J.F. Klimavicz (2012)
Fls double, 2–3/ball truss, saucer-shaped when fully opened, 40 × 70mm, with 13–16 very wavy-edged lobes, buds white, shading to light yellow-green (142C) at the base, opening white throughout, with a faint spotted blotch of light yellow-green (142C) at the centre (the blotch may be more pronounced or absent altogether). Calyx 10mm long, strong yellow-green (144A). Lvs elliptic, 50 × 20mm, broadly acute at apex, cuneate at base, with flat margins, glossy and moderate olive green (137A) above, hairless. Dense shrub, 0.5 × 0.5m in 5 years. Mid May.

(Etymology: named at the request of Richard Taffet, as a gift to his daughter)

(r) **‘Rosebery Gold Mine’**

cv. Lepidote rhododendron: parentage unknown

H: unknown, G: Emu Valley Rhododendron Garden (2003), N: MMG Ltd (2012), I: Emu Valley Rhododendron Garden, REG: Emu Valley Rhododendron Garden (2012)

Fls 8/loose truss, broadly tubular-campanulate, 35 × 45mm, with 5 wavy-edged lobes, moderate red (180B) at apex in bud, opening inside light greenish yellow (4B), outside light greenish yellow (4B), lightly suffused strong red (51A) on the lobes; very faint greenish yellow spotting at the base of the dorsal throat; filaments and style pale greenish yellow; anthers pale brown; stigma greenish. Lvs elliptic, 70 × 28mm, margins flat, acute at apex, cuneate at base, matt green above, with scales. Shrub 1.2 × 1.0m in 10 years. Early August.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named in part for the mining town of Rosebery, Australia, partly owned by the Chinese company MMG Ltd (Minerals and Metals Group), who have donated money to the Emu Valley Rhododendron Garden for a Chinese pavilion in the Garden)

(a) **‘Selga’**

cv. Deciduous azalea: ‘Aida’ (s) × ‘Uguns’

H: R. Kondratovičs (1978), G: R. Kondratovičs (1984), N: R. Kondratovičs (2012),

REG: Rhododendron Nursery Babīte, University of Latvia (2012)

Fls single, 13/slightly domed truss, tubular funnel-shaped, 60 × 50mm, with 5 very wavy-edged lobes, buds brilliant orange-yellow (23B), opening inside and out brilliant yellow (14C), with a strong orange-yellow (17A) blotch on the dorsal lobes; filaments, anthers and style yellow, stigmas green; slightly scented. Calyx 2mm long, green. Lvs elliptic, 120–150 × 45–55mm, pointed at base and apex, wavy-edged, matt green above, hairless. Shrub 2.0 × 2.5m in 20 years. Late May to early June.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for one of the Latvian University student corporations, Selga)

(r) **‘Sparkling Stars’**

cv. Elepidote rhododendron: ‘Yaku Sunrise’ × ‘Hansel’ (s) × ‘Lem’s Cameo’

H: F. Fujioka (1983), G: F. Fujioka (1988), N: D.S. Wallace (2011), I: Singing Tree Gardens & Nursery (2011), REG: Singing Tree Gardens & Nursery (2012)

Fls 9/domed truss, open funnel-shaped, c.40 × 65mm, with 6 smooth to very slightly wavy-edged lobes, the lobes almost separated to the base and often with shallowly toothed margins, yellowish white (158D) in bud, opening white inside and out, with a prominent, deep purplish pink (55A) blotch on the dorsal lobe and with some purplish suffusion on the midveins outside. Lvs elliptic to obovate, 150 × 50mm, broadly acute at apex, cuneate at base, margins decurved, glossy and strong green (132C) above, hairless. Shrub 2.1 × 2.1m in 27 years; the shoots with prominent red bracts. Mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). A sister seedling of ‘Starbright Champagne’; the very deep lobes give the flowers a very “spiky” appearance)

(r) **‘Spring Sprite’**

cv. Elepidote rhododendron: ‘Capistrano’ (s) × ‘September Song’

H: J. Doppel (2005), G: J. Doppel (2009), N: J. Doppel (2011), REG: J. Doppel (2012)

Fls 10/domed truss, 40 × 50mm, with 5 wavy-edged lobes, strong red (41B) in bud, opening inside light yellow (20B) at centre, shading outward to light yellow (20D), edged with moderate yellowish pink (31D) and with a strong red (41B), speckled dorsal blotch; outside moderate yellowish pink (31D), shading to strong yellowish pink (37A) at edges. Lvs elliptic, 65 × 32mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and mid green above, hairless. Dense shrub, 0.4 × 0.75m in 7 years. Mid May.

(r) **‘Stordagull’**

cv. Elepidote rhododendron: ‘September Song’ (s) × ‘Berg’s Yellow’

H: M. Chernet (1995), G: K.E. Grov (2008), N: P.M. Jørgensen (2012), I: Gimle Planteskule, REG: P.M. Jørgensen (2012)

Fls 6–10/loosely domed truss, open funnel-shaped, 30–40 × 35–45mm, with 5–6 wavy-edged lobes, pinkish orange in bud, opening inside orangey pink, with yellow in the centre, outside pinkish orange, unmarked; reported to be a particularly glowing colour. Calyx 1–2mm long, green, with a reddish margin. Lvs lanceolate, 60–100 × 30–40mm, cuneate at base, obtuse at apex, margins flat, hairless. Shrub 0.75 × 0.5m in 10 years. Early May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). First published and illustrated in *Lapprosen* 2: 4–6 (2012). Etymology: “Stord” is the place in Norway where the plant first flowered, “gull” means “gold” in Norwegian and refers to the colour of the flowers)

(r) **‘Sun Music’**

cv. Elepidote rhododendron: ‘Sun Quest’ (s) × ‘Peach Recital’

H: J. Barlup (2004), G: J. Barlup (2008), N: J. Barlup (2010), REG: J. Barlup (2012)

Fls 18/ball truss, broadly funnel-shaped, 45 × 75mm, with 5 wavy-edged lobes, deep pink (48A) in bud, opening inside pale yellow (11D), with dorsal lobes a slightly darker shade of light yellow (11B) starting c.15mm from the base and extending

upwards, and with deep red (53A) dorsal spotting beginning c.15mm from the base and extending c.20mm; outside pale yellow (11D), with light yellow (11B) midveins; not scented. Calyx c.20mm long, pale yellow (11D), tipped with light yellow (11B). Lvs elliptic, 105 × 45mm, margins flat, broadly acute at apex, rounded at base, dull and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 7 years. Mid May.

(r) **‘Sun Spirit’**

cv. Elepidote rhododendron: ‘Recital’ (s) × ‘Goldprinz’
H: J. Barlup (2002), G: J. Barlup (2007), N: J. Barlup (2011), REG: J. Barlup (2012)
Fls 13/domed truss, funnel-campanulate, 35 × 75mm, with 6 wavy-edged lobes, strong yellowish pink (34D) in bud, opening inside light yellow-green (150D), the dorsal lobes a slightly darker shade of light yellow-green (154D), with two deep red (53A) dorsal flares extending c.5mm from the base and speckling of the same colour extending c.20mm above the flares; outside similar to the inside, but without the marking; not scented. Calyx insignificant. Lvs elliptic, 95 × 45mm, margins flat, broadly acute at apex, rounded at base, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.9m in 9 years. Mid May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY))

(r) **‘Teton Twilight’**

cv. Elepidote rhododendron: ‘Coral Blossom’ (s) × ‘Christina Dee’
H: J. Barlup (2005), G: J.D. Winberg (2010), N: J. Barlup (2011), REG: J. Barlup (2012)
Fls 16/domed truss, broadly funnel-shaped, 65 × 90mm, with 5 wavy-edged lobes, deep pink (52B) in bud, opening inside light purplish pink (63D), edged strong purplish red (63A), lightly tinged pale yellow (18D) at the base, and with two c.10mm long, solid, dark red (187B), dorsal flares, changing to small spots and spreading upwards; outside light purplish pink (63D), with strong purplish red (63B) midribs; not scented. Calyx 25mm long, pale yellow (18D), with dark red (187B) flares and

moderate red (183D) spotting. Lvs elliptic, 120 × 55mm, margins flat, broadly acute at apex, rounded at base, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 1.2m in 6 years. Late May.

(r) **‘Tomasz Wojciech’**

cv. Elepidote rhododendron: open-pollinated seedling from ‘Scintillation’ (s)
H: P. Muras (1982), G: P. Muras (1987), N: P. Muras (1996), REG: Mrs M. Muras (2012)
Fls 12–15/ball truss, open funnel-shaped, 50–55 × 60–80mm, with 5 wavy-edged lobes, buds light purple, opening light purple (75B) both inside and out, with light olive to dark greenish yellow (152B/C) spots on the dorsal lobes. Calyx 3mm long, green. Lvs elliptic, 75–110 × 30–45mm, obtuse at apex, rounded at base, margins flat, slightly glossy and moderate green above, hairless. Shrub 1.5 × 2.5m in 20 years. Mid May (in southern Poland).
(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a friend of the registrant’s family and his former teacher. First published in *Szkółkarstwo* 2: 54 (2005))

(r) **‘Tropical Gem’**

cv. Elepidote rhododendron: ‘Mindy’s Love’ (s) × (‘Lem’s Cameo’ × *forrestii* subsp. *forrestii*) × (‘Lem’s Cameo’ × ‘Brandt’s Tropicana’)
H: J. Barlup (1988), G: J. Barlup (1994), N: B. Johnson (2004), I: Hazelwood Gardens, REG: J. Barlup (2012)
Fls 17/domed truss, broadly funnel-shaped, 55 × 90mm, with 7 wavy-edged lobes, strong red (53B) in bud, opening inside pale yellowish pink (27D), blending to pale greenish yellow (13D) towards the base, with an c.6mm wide, strong pink (50C), central stripe on each lobe, extending from the base to the edge of the corolla, and moderate red (181B) dorsal spotting extending 40mm from the base; outside similar except stripes are deep pink (50B) and there is no spotting; not scented. Calyx c.20mm long, pale greenish yellow (13D), blending into strong pink (50C). Lvs elliptic, 100 × 50mm,

margins upcurved, broadly acute at apex, rounded at base, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.8m in 5 years. Early May.

(a) **'TTT'**

cv. Deciduous azalea: 'Uguns' (s) × Exbury azalea seedling (Ex-5)

H: R. Kondratovičs (1976), G: R. Kondratovičs (1982), N: R. Kondratovičs (2006),

REG: Rhododendron Nursery Babīte, University of Latvia (2012)

Fls single, 10/slightly domed truss, tubular funnel-shaped, 55mm wide, with 5–7 wavy-edged lobes, buds vivid reddish orange (N30A), opening inside brilliant yellow (21C) and strong orange (25A), outside light orange-yellow (22C); dorsal lobe strong orange (N25C); filaments yellow-orange, anthers orange; style yellow-orange, stigma green. Calyx 2mm long, green. Lvs elliptic, 100 × 35mm, pointed at base and apex, margins wavy, matt green above, hairless. Shrub 2.0 × 1.7m in 15 years. Early June.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a famous Latvian basketball team (TTT))

(r) **'Twilight Touch'**

cv. Elepidote rhododendron: 'Light Years' (s) × 'Percy Wiseman'

H: J. Barlup (2002), G: J. Barlup (2007), N: J. Barlup (2011), REG: J. Barlup (2012)

Fls 16/ball truss, open funnel-shaped, 45 × 60mm, with 5 wavy-edged lobes, vivid red (46B) in bud, opening inside light purplish pink (55C) on ventral lobes, and light purplish pink (55C) blending to pale yellow (19C) on dorsal lobes, with strong purplish pink (55B) dorsal spotting; outside similar except for the lack of spotting and the addition of strong purplish pink (55B) midribs on all lobes; not scented. Calyx c.15mm long, light yellowish pink (19B), edged strong purplish pink (55B). Lvs elliptic, 100 × 50mm, margins upcurved, broadly acute at apex, rounded at base, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 9 years. Mid May.

(r) **'Vilhelmīne Petkevičs'**

cv. Elepidote rhododendron: *yakushmanum*

'Koichiro Wada' (s) × 'Purple Splendour'

H: R. Kondratovičs (1974), G: R. Kondratovičs (1980), N: R. Kondratovičs (2010),

REG: Rhododendron Nursery Babīte, University of Latvia (2012)

Fls 12/slightly domed truss, open funnel-shaped, 50–55mm wide, with 5 wavy-edged lobes, buds deep purplish pink (N66C), opening inside very pale purple (76D), with very light purple (76C) margins, outside deep purplish pink (N66D), dorsal lobe with several relatively inconspicuous spots of strong greenish yellow (153B); filaments and anthers white, style light purple, stigma cerise. Calyx 1mm long, light green. Lvs oblanceolate, 110 × 45mm, pointed at apex, rounded at base, margins flat, matt green above, with light brown hairs below which age to brown. Shrub 2.0 × 2.5m in 20 years. Late May to early June.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: a Latvian female christian name)

(r) **'Vizma'**

cv. Elepidote rhododendron: 'Catawbiense Grandiflorum' (s) × 'Panenka'

H: R. Kondratovičs (1978), G: R. Kondratovičs (1987), N: R. Kondratovičs (2005),

I: Rhododendron Nursery Babīte, University of Latvia (2005), REG: Rhododendron Nursery Babīte, University of Latvia (2012)

Fls 11/slightly domed truss, open funnel-shaped, 80–85mm wide, with 5–7 wavy-edged lobes, buds deep purplish pink (66C), opening inside very pale purple (75D), with a light purple (75B) margin, outside very pale purple (75C), dorsal lobes with strong yellowish green (N144A) spots; filaments, anthers and style white, stigmas light yellow-green. Calyx 2mm long, very light green. Lvs elliptic, 125–140 × 50–55mm, pointed at apex, rounded at base, margins flat, matt green above, hairless. Shrub 3.0 × 2.5m in 20 years. Early June.

(Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: a Latvian female christian name)

(r) **'Vyron Minor'**

cv. Elepidote rhododendron: 'Janet Blair' × *decorum* (yellow form) (s) × 'Helen Vieira'

H: J.D. Brim (2002), G: J.D. Brim (2005), N: J.D.

Brim (2012), REG: J.D. Brim (2012)

Fls 20/ball truss, open funnel-shaped, 57 × 100mm,

with 6 wavy-edged lobes, strong purplish red (54A)

in bud, opening inside moderate purplish pink

(54D), with a deep red (60A) blotch, outside strong

pink (54C). Lvs obovate, 180 × 63mm, broadly

acute at apex, cuneate at base, semi-glossy and

moderate olive green (137B) above, hairless. Shrub

1.3 × 1.7m in 10 years. Midseason.

(Etymology: named for a daughter of the hybridiser)

(r) **'White Marsh'**

cv. Elepidote rhododendron: parentage unknown (s)

H: J. Doppel (2006), G: J. Doppel (2007), N: J.

Doppel (2011), REG: J. Doppel (2012)

Fls 9/ball truss, open funnel-shaped, 90 × 75mm,

with 5 wavy-edged lobes, vivid purplish red (57D) in

bud, opening inside pale purplish pink (56A),

shading to white; lightly spotted vivid purplish red

(57D), changing to moderate orange-yellow (168D)

on the lower half of the dorsal lobe; outside deep

purplish pink (68C); not scented. Calyx

insignificant. Lvs elliptic, 115 × 66mm, margins

downcurved, broadly acute at apex, rounded at base,

semi-glossy and mid green above, hairless. Shrub 0.6

× 0.8m in 5 years. Mid May.

Names and addresses

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

* = deceased

Barlup, J.
Bellevue, Washington, USA
Beaudry, N. & J.
Bethesda, Maryland, USA
Briarwood Gardens,
East Sandwich, Massachusetts, USA
Brim, J.D.
Mount Airy, North Carolina, USA
Burns, F.E.
Vida, Oregon, USA
*Chernut, M.
USA
Clagett, W.L.
Derwood, Maryland, USA
Colville, Mr & Mrs J.
Launceston, Cornwall, England
*Consolini, A.
Sandwich, Massachusetts, USA
Coombes, L.
Burnie, Tasmania, Australia
*Cowles, J.C.
Stow, Massachusetts, USA
Delano, J.
Duxbury, Massachusetts, USA
Doppel, J.
Lenhartsville, Pennsylvania, USA
Emu Valley Rhododendron Garden
Burnie, Tasmania, Australia
Fujioka, F.
Freeland, Washington, USA
Gilberg, G.
Bergen, Norway
Gimle Planteskule
Nordheimsund, Norway
Gro, K.R.
Stord, Norway
Hachmann, H. (Hans Hachmann)
Baumschule H. Hachmann, Barmstedt, Germany
Hachmann, Holger
Baumschule H. Hachmann, Barmstedt, Germany

Hazelwood Gardens
Newcastle, Washington, USA
Heritage Museums and Gardens
Sandwich, Massachusetts, USA
Hübbers, K.-H. & Mrs L.
Kranenburg, Germany
Ijima, S.
Fukushima-city, Fukushima-prefecture, Japan
Johnson, B.
Newcastle, Washington, USA
Jørgensen, P.M.
Det Norske Arboret, Hjeljestad, Norway
Klimavicz, J.F.
Vienna, Virginia, USA
Kline, H.B.
Newmanstown, Pennsylvania, USA
Kondratovičs, R.K.
Riga, Latvia
Kruppa, R.
Volcano, Hawaii, USA
Kupsch, M.
Burnie, Tasmania, USA
Le Duigou, A. & L.
Raden, Scaer, France
Lilly, C.
Falmouth, Massachusetts, USA
Malone, H.R.
Lapoinya, Tasmania, Australia
Meerkerk, A.
Greenbank, Washington, USA
Meerkerk Rhododendron Gardens
Greenbank, Washington, USA
MMG Ltd
Melbourne, Victoria, Australia
*Moffat, J.A.R.
Launceston, Cornwall
Muras, P. & Mrs M.
Wieliczka, Poland
O'Connor, J.M.
Osterville, Massachusetts, USA
Oppenrieder, M.
Baumschule H. Hachmann, Barmstedt, Germany

Pacific Island Nursery
Kea'au, Hawaii, USA
*Rhein, W.L.
Mechanicsburg, Pennsylvania, USA
Rhein Registration Committee, ARS Susquehanna
Valley Chapter, USA
Rhododendron Nursery Babite, University of Latvia
Riga, Latvia
Saint-Jalm, J.F.
Guémené-sur-Scorff, France
Sandwich Club
Sandwich, Massachusetts, USA
Scaer Town Council
Scaer, France
Shibata, Y.
Okazaki-city, Aichi-prefecture, Japan
Singing Tree Gardens
McKinlayville, California, USA
Société Bretonne du Rhododendron
Scaer, France
Sullivan, N.
Burnie, Tasmania, Australia
*Sullivan, P.
San Francisco, California, USA
Suzuki, H.
Koriyama-city, Fukushima-prefecture, Japan

Takahashi, T.
Chikusei-city, Ibaraki-prefecture, Japan
Trautmann, C.
Florence, Oregon, USA
Van Oost, H.
Kruishouten, Belgium
Wallace, D.S.
Singing Tree Gardens, McKinlayville, California,
USA
Westhoff, J.
Bremen, Germany
White, R.H.
Sandwich, Massachusetts, USA
White's Nursery
Germantown, Maryland, USA
Winberg, J.D.
Fall City, Washington, USA
Woodard, G.
Westbury, New York, USA
Wood, J.
c/o Emu Valley Rhododendron Garden
Woods, J.
Burnie, Tasmania, Australia

Corrections to the International Rhododendron Register and Checklist

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to the seven subsequent Supplements.

The Registrar is grateful to all those who have brought such errors to his notice and welcomes knowledge of further amendments that may need to be made. Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version. The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the *Register and Checklist* (or one of its Supplements) on which the entry concerned is to be found.

‘Bernadette Chirac’ (p.130)

The roles have been omitted and should read as follows: H: E de Rothschild, 1985, G: E. de Rothschild, 1992, N: E de Rothschild, REG: E de Rothschild, 1996

‘Dartmoor Rose’ (p.316)

This Checklist entry is the plant subsequently incorporated as a Register entry in the Second Supplement, p.11 (2007)

‘Goldinetta’ (p.531)

This epithet was subsequently shown to be a registered trade mark in Germany and should thus be treated here as a trade designation (GOLDINETTA). The name ‘Hachgold’ replaced it as the cultivar epithet, but this was subsequently used for another plant which was granted EU PBR as ‘Hachgold’, and is sold under the trade mark GOLDEN EVEREST. The original ‘Goldinetta’ should now be distinguished under the accepted cultivar epithet ‘Hachinetta’

‘Leitmotif’ (Seventh Supplement, p.17)

The parentage was supplied in error: the plant was raised from seed of unknown parentage, supplied by Hobbie

‘Le Vainqueur’ (p.783)

The spelling should be corrected to ‘Le Vainqueur’

‘Our Kate Marise’ (Fifth Supplement, p.24)

The plant registered under this name in 2008 is now considered to be the same as that registered as ‘Our Kate’ in 1963 and must be treated as a synonym. The confusion arose because Exbury had thought that the AM plant (named as ‘Our Kate’) was a form of the Our Kate Group with pink-flushed flowers (see comments in *The Rothschild Rhododendrons*, 1979), but the published AM description and the registration from 1963 make it clear this was not the case. A second plant from this Group (description not recorded) was exhibited at the same time in 1963 and may have been this pink-flushed plant, which is now thought to have been lost

‘Promenade Home’ (p.1113)

This was listed incorrectly as an evergreen azalea; it is correctly an elepidote rhododendron (as the parentage listed implies)

‘Schöne von Gressen’ (p.1233)

This should read ‘Schöne von Giessen’. It was raised by F. Rehelt in Giessen, Germany, pre1916; it also has *occidentale* in its parentage

‘Special Occasion’ (p.1294)

The spelling should be corrected to ‘Special Occasion’

‘Vainqueur’ (p.1402)

Both entries under this name should be corrected to read ‘Vainqueur’ and the cross reference under the first entry to ‘Le Vainqueur’ should read ‘Le Vainqueur’

ISBN 9781907057427



9 781907 057427