



Royal  
Horticultural  
Society

Sharing the best in Gardening

# The International Rhododendron Register & Checklist (2004)

*Ninth Supplement*



© 2014 The Royal Horticultural Society  
80 Vincent Square, London SW1P 2PE, United Kingdom  
[www.rhs.org.uk](http://www.rhs.org.uk)

International Rhododendron Registrar:  
Dr A C Leslie

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057519

Printed and bound in the UK by Page Bros, Norwich

The previous supplement (Eighth Supplement) was published on 17 February 2014

Cover: *Rhododendron* 'Mrs Charles E. Pearson', by Alfred John Wise  
Image courtesy of RHS Lindley Library

# The International Rhododendron Register and Checklist (2004)

## *Ninth Supplement*

Names registered, January–December 2013	page 2
Names and addresses	page 24
Corrections	page 26

## Introduction

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first eight Supplements included names registered from 1 January 2003 to 31 December 2012. This ninth Supplement includes all those registered in 2013.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7–10 of the Introduction.

It may be helpful here to repeat that the following abbreviations are employed:

(a)	azalea
(az)	azaleodendron
cv.	cultivar
G	grown to first flower by
gp	Group
H	hybridized by
I	introduced by
N	named by
(r)	rhododendron
R	raised by
REG	registered by
S	selected by
(s)	seed parent
(v)	vireya

Where possible colour references are given to the *RHS Colour Chart* (1966, 1986, 1995 & 2001) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website ([www.rhs.org.uk/plants/pdfs/plant-registration-forms/rhododendron-name-registration](http://www.rhs.org.uk/plants/pdfs/plant-registration-forms/rhododendron-name-registration)) or from one of the national or regional registrars as follows:

**North America** Mr Michael Martin Mills, 632 Burnham Road, Philadelphia, Pennsylvania PA19119-3510, USA

**Australia** Mrs Lesley Eaton, PO Box 597, Snug, Tasmania 7054, Australia

**New Zealand** Mr Brian J. Coker, Stonecrop, 8B Barnsley Crescent, West Melton, New Zealand 7618

**Japan** Dr Katsuhisa Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

**Dr A.C. Leslie, International Rhododendron Registrar, 109 York Street, Cambridge CB1 2PZ, UK**  
(email: [alanleslie@rhs.org.uk](mailto:alanleslie@rhs.org.uk))

# Names registered January–December 2013

## (r) 'Amber Waves'

cv. *Elepidote rhododendron*: 'Amber Touch' (s) × 'Janet Blair'

**H:** J. Barlup (1996), **G:** J. Barlup (2002), **N:** J. Barlup (2011), **I:** Whitney Gardens (2013), **REG:** J. Barlup (2013)

Fls 22/ball truss, broadly funnel-shaped, 57 × 89mm, with 5 very wavy-edged lobes, strong purplish red (54A) in bud, opening inside pale yellow (12D) at centre of flower, the apical half of each lobe light yellow (12C), with a 6mm wide margin of strong pink (54C), moderate orange (166D) spotting on dorsal lobes, starting 13mm from the base and extending 32mm; outside pale yellow (12D), the apical portion of the lobes light yellow (12C), with a 6mm wide margin of strong pink (54C); not scented. Calyx insignificant. Lvs elliptic, 140 × 44mm, rounded at base, broadly acute at apex, with wavy margins, dull and moderate olive green (147A) above, hairless. Shrub 1.2 × 0.9m in 9 years. Midseason.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the name is descriptive of the flower)

## (r) 'Anne Laure de Beau Vallon'

cv. *Elepidote rhododendron*: 'Nancy Evans' (s) × *fortunei*

**H:** J.-F. Saint-Jalm (1999), **G:** J.-F. Saint-Jalm (2006), **N:** J.-F. Saint-Jalm (2013), **REG:** J.-F. Saint-Jalm (2013)

Fls 12/ ball truss, open funnel-shaped, 60 × 87mm, with 7 wavy-edged lobes, deep pink (52C) in bud, opening pale yellow (12D) inside and out, with a pale pink flush around the margins when young, with a small, strong purplish red (60B) blotch at the base of the dorsal throat; stamens 14, filaments yellow; style yellow, flushed pink, glandular, stigma green, becoming yellow. Calyx 18mm long, pale yellow (12D), edged purplish red. Lvs elliptic to obovate, 110 × 46mm, cuneate at base, broadly acute and mucronate at apex, greyish olive green (N137B) above, hairless. Shrub 2.2 × 1.9m in 13 years. Early to mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a daughter of the registrant (Anne Laure) and the village in France where they live (Beau Vallon))

## (r) 'Annie Fish'

cv. *Elepidote rhododendron*: parentage unknown  
**H:** J.C. Cowles (c. 1970), **G:** J.C. Cowles (1970s), **N:** A. Morrissey (2013), **REG:** Heritage Museums and Gardens (2013)

Fls 11/ball truss, open funnel-shaped, 57 × 89mm, with 7 wavy-edged lobes, deep purplish pink (73A) in bud, opening inside pale purplish pink (65D), with a light yellow-green (145B) pistil; outside light purplish pink (65B). Calyx insignificant. Lvs oblong, 127 × 57mm, oblique at base, broadly acute at apex, margins downcurved, matt and moderate olive green (137B)

above, hairless. Shrub 3.7 × 3.7m in 40 years.

Midseason (May in Cape Cod, Massachusetts, USA). (Nomenclatural standard: colour print of an electronic image (WSY). Etymology: named for the mother-in-law of Cyndy Fish, the latter a supporter of the Heritage Museums and Gardens (Sandwich, Massachusetts, USA))

## (a) 'Antoinette Martin'

cv. Evergreen azalea: 'Festive' (s) × 'Satellite' (Klupinger)

**H:** J.F. Klimavicz (1997), **G:** J.F. Klimavicz (1999), **N:** T. Burnham & S. Visner (2013),

**REG:** J.F. Klimavicz (2013)

Fls 2/truss, broadly funnel-shaped, 35 × 70mm, with 5 wavy-edged lobes, deep red (60A) in bud, with some white and deep purplish pink (58D) striping, opening inside with white on the sides of the lobes and a deep purplish pink (58D) central stripe extending from the centre of the flower to the lobe apex (and roughly covering one third of the lobe width); base of the throat deep red (60A), this colour extending on the 3 dorsal lobes as wide rays almost to the margin (irregularly centred within the pink stripe and thus forming a prominent rayed blotch; some irregular deep red (60A) flecks; outside deep red (60A), with deep purplish pink stripes down the centre of each lobe to the tip, the sides of the lobes white. Calyx 10mm long, strong yellow-green (144B). Lvs elliptic, 40 × 22mm, cuneate at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Dense shrub, 0.5 × 0.5m in 5 years. Early (April in northern Virginia).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the late Antoinette Martin (formerly of Washington, D.C.), a friend of the nominants; the latter won the naming rights in a benefit auction for the US National Institutes of Health Children's Inn in Bethesda, Maryland, USA. Hybridizer's ref. no. FT-97-4)

## (v) 'April Vows'

cv. *Lepidote rhododendron*: 'Neesa' (s) × *konori* var. *phaeocephalum*

**H:** G.L.S. Snell (2004), **G:** N. Puddey (2010),

**N:** K. Puddey, **I:** N. Puddey (2012),

**REG:** N. Puddey (2013)

Fls 8–12/truss, tubular funnel-shaped, 80 × 75mm, with 5–6 flat-edged lobes, dark red (187A) in bud, opening yellowish white (155B), deepening to pale yellow (11D) in the centre of the flower, outside yellowish white (155B), deepening to pale yellow (11D) along the tube; unmarked; scented. Calyx less than 1mm long, dark yellowish pink (181D). Lvs broadly elliptic, 100 × 50mm, broadly tapering at base, obtuse at apex, margins decurved, glossy green above, scales pale yellow (158B) at first, later moderate reddish brown (174A). Shrub 1.4 × 0.6m in 8 years. Flowering irregularly, 2–3 times a year.

(Nomenclatural standard: colour print provided by the registrant (WSY0119004). Etymology: considered to be a fitting component of bouquets for April weddings)

(r) **'Arnulf'**

cv. Elepidote rhododendron: *aureum* (s) × *caucasicum*

**H:** A. Ringstad (c. 1990), **G:** A. Ringstad (c. 1995),  
**N:** P.M. Jørgensen (2013), **I:** A. Ringstad (c. 2000),  
**REG:** P.M. Jørgensen (2013)

Fls 5–7/flat truss, broadly campanulate, 15–20 × 30mm, with 5 wavy-edged lobes, greenish yellow in bud, opening inside pale yellow, outside creamy yellow when fully developed; numerous greenish dots over the dorsal lobe (including the throat) and on parts of the adjacent lobes. Calyx 2mm long, green, pilose. Lvs narrowly ovate, 40–50 × 15–25mm, cuneate at base, apiculate at apex, margins flat, with a brown indumentum below, which becomes brownish and more scattered with age. Shrub 0.3–0.5 × 1.0m in 10 years. Early May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: christian name of the breeder)

(a) **'Auseklitis'**

cv. Semi-evergreen azalea: 'Adventglöckchen' (s) × 'Esmeralda'

**H:** R. Kondratovičs (1972), **G:** R. Kondratovičs (1975), **N:** R. Kondratovičs (1975), **REG:** Botanical Garden, University of Latvia (2013)

Fls single, 2–3/truss, funnel-shaped, 30 × 40–45mm, with 5 slightly wavy-edged lobes, deep pink (52B) in bud, opening deep purplish pink (55A) inside and out, with inconspicuous, strong purplish red (54A), dorsal spotting; style white, stigma light green. Calyx 3–4mm long, light green. Lvs elliptic, 19–40 × 12–22mm, pointed at base, rounded at apex, margins flat, matt green above, hairless. Shrub 0.8 × 0.8m in 15 years. February to March (under glass).

(Nomenclatural standard: colour print provided by the registrant (WSY0119338). Etymology: described as the folklore name of the morning star)

(r) **'Beni-Shujaku'**

cv. Elepidote rhododendron: selection from *degronianum* subsp. *heptamerum* var. *hondoense*  
**S:** Y. Yamamoto (2011), **N:** Y. Yamamoto (2013),  
**REG:** Y. Yamamoto (2013)

Fls 3–10 (average 7)/loose, rounded truss, open funnel-shaped, 35 × 50mm, with 7 flat-edged lobes, deep pink in bud, opening deep pink both inside and out, non-fading, unmarked. Calyx 5mm long, brown. Lvs elliptic, 80 × 30mm, narrowed at base, acute at apex, margins flat, with dense light brown hairs below when young, which age to a darker brown. Shrub 10 × 12cm in 2 years after grafting. Early May.

(Nomenclatural standard: colour print provided by the registrant (WSY0119070). Raised from seed sown by the registrant in 2007)

(a) **'Birdie Mac'**

cv. Evergreen azalea: 'Elsie Lee' × 'Dorsett' (s) × 'Helen Curtis'

**H:** W.L. Clagett (1996), **G:** W.L. Clagett (2001),  
**N:** W.L. Clagett (2013), **I:** White's Nursery,  
**REG:** W.L. Clagett (2013)

Fls hose-in-hose, 3/truss, broadly funnel-shaped, 25 × 32mm, with 5 + 5 wavy-edged lobes, buds strong red

(39A), with off-white along the midribs, inside opening deep yellowish pink (39B) fading to off-white in the centre of each lobe, with heavy strong purplish red (63A-B) spotting on the 3 dorsal lobes; outside deep yellowish pink (39B), with off-white on the midribs. Calyx 5mm long, pale green, with white hairs. Lvs elliptic, 16 × 6mm, cuneate at base, broadly acute at apex, margins flat, semi-glossy and dark yellowish green (139A) above, with white hairs below when young. Floriferous shrub, 1.5 × 0.9m in 17 years. Midseason (May in Washington, D.C., USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: nickname of the registrant's older sister – derived from the American comic strip "Snuffy Smith". Hybridizer's ref. no. 96-2-15)

(a) **'Bob Stewart'**

cv. Evergreen azalea: un-named hybrid (ref. no. CB-1) × 'Elsie Lee' (s) × 'James Stewart'

**H:** J.F. Klimavicz (2001), **G:** J.F. Klimavicz (2003),  
**N:** J.F. Klimavicz (2013), **REG:** J.F. Klimavicz (2013)

Fls fully double, 2–3/truss, saucer-shaped, 35 × 70mm, with 16–18 wavy-edged lobes (the outer corolla whorl 5-lobed), buds deep purplish pink (72D), opening deep purplish pink (72D), fading to very pale purple (73D), with strong purplish red (59D) dorsal spotting; outside deep purplish pink (72D); occasional deformed stamens or pistil present, but these not petaloid. Calyx 10mm long, strong yellow-green (144B). Lvs elliptic, 40 × 20mm, cuneate at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A), hairless. Dense shrub 0.5 × 0.5m in 5 years. Early (April in northern Virginia).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: named for an azalea breeder and friend of the registrant. Hybridizer's reference no. L2-01-6. The parent CB-1 was a plant discarded from the University of Maryland's breeding programme in the 1990s: its parentage is not known)

(r) **'Bontebok'**

cv. Elepidote rhododendron: 'Ring of Fire' (s) × 'Yaku Angel'

**H:** F. Schnupper (2004), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls 8–9/domed truss, funnel-campanulate, 45–60 × 50–70mm, with 5 wavy-edged lobes, vivid reddish orange (33B) in bud, opening inside strong yellowish pink (33C), outside vivid reddish orange (33B); brown spotting on dorsal lobe. Calyx 5mm long, green. Lvs elliptic, 80–100 × 30–40mm, cuneate at base, broadly acute at apex, margins flat and decurved, matt green above, hairless. Shrub 0.8 × 1.0m in 10 years. Mid to late May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in South Africa)

(a) **'Cape Fear Gold'**

cv. Deciduous azalea: *austrinum* (seed parent grown from seed collected at Brookgreen Gardens, South Carolina, USA) (s) × *calendulaceum* (pollen collected

on Roan Mountain, North Carolina, USA)

**H:** L.M. Player (1978), **G:** L.M. Player (c. 1982),

**N:** L.M. Player (2013), **REG:** L.M. Player (2013)

Fls 7–11/bud (with 1–4 buds/shoot), tubular funnel-shaped, 44 × 38mm, with 5 wavy-edged lobes, buds vivid yellow (15A), opening vivid yellow (16A), with a strong orange-yellow (17A) blotch covering almost all of the dorsal lobe; outside light orange-yellow (16B); stamens and style extend 40mm beyond the corolla; moderately scented. Calyx 1mm long, moderate yellow-green (146C). Lvs elliptic, 50 × 19mm, cuneate at base, broadly acute at apex, margins flat, matt and strong yellow-green (144A) above, hairless. Shrub 2.4 × 2.4m in 15 years. April (Fayetteville, North Carolina, USA). Reported to be very heat tolerant.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the Cape Fear River (North Carolina, USA), in whose valley the registrant lives. Hybridizer's no. Player-1-79-1)

(v) 'Captain Ollie'

cv. *Lepidote rhododendron*: parentage unknown

**H:** N. Puddey (2010), **G:** N. Puddey, **N:** N. Puddey

(2012), **I:** N. Puddey (2012), **REG:** N. Puddey (2013)

Fls 10–12/full truss, funnel-shaped, 60 × 55mm, with 5–6 flat-edged lobes, dark red (187A) in bud, opening vivid reddish orange (41A) on most of each lobe, light yellow (17D) towards the base of each lobe and in the throat, outside vivid reddish orange (41A) towards the apex of each lobe, otherwise pale yellow (18D) and this colour extending onto the tube; unmarked. Calyx less than 1mm long, strong yellowish pink (33C). Lvs broadly elliptic, 110 × 65mm, rounded at base, obtuse at apex, margins flat, glossy green above, scales greenish white (192D) at first, not evident later. Shrub 0.6 × 0.3m in 3 years, with dark reddish brown internodes and leaf midribs. Flowering at irregular intervals.

(Nomenclatural standard: colour print provided by the registrant (WSY0119042). Etymology: named for the grandson of the registrant)

(r) 'Carol's Candy'

cv. *Elepidote rhododendron*: 'Lem's Tangerine' (s) × 'Tia'

**H:** J. Barlup (2000), **G:** J. Barlup (2005), **N:** J. Barlup

(2010), **REG:** J. Barlup (2013)

Fls 14/domed truss, open funnel-shaped, 51 × 70mm, with 7 wavy-edged lobes, moderate yellowish pink (38B) in bud, opening inside pale orange-yellow (159C), with strong red (53D) midveins from base to apex, margins strong pink (54C), dorsal lobes deep purplish pink (54B), with a strong red (53B) flare on the dorsal lobe (32mm long) and with 6mm long, strong red (53B) stripes on each lobe; outside pale orange-yellow (159C), with strong red (53D) midveins and deep purplish pink (54B) dorsal lobes; not scented. Calyx insignificant. Lvs elliptic, 120 × 50mm, rounded at base, broadly acute at apex, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 1.1 × 1.1m in 12 years. Midseason (April, in Bellevue, Washington, USA). (Nomenclatural standard: colour print of an

electronic image provided by the registrant (WSY).

Etymology: named for a friend of the hybridizer, who likened the truss to candy)

(r) 'Cemetery Coral'

Syn. of 'Howard J'

(r) 'Chobe'

cv. *Elepidote rhododendron*: *smirnowii* (s) × unknown

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F.

Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)

Fls 14–15/conical truss, broadly funnel-shaped, 25–35 × 60mm, with 5–6 wavy-edged lobes, strong reddish purple (70B) in bud, opening moderate purplish pink (70D) inside, deep purplish pink (70C) outside; with strong reddish purple (70B) spots on the dorsal lobe; white stamens; very prominently exerted, pale style. Calyx 2mm long, white. Lvs elliptic, 120–150 × 40mm, rounded at base, broadly acute at apex, margins wavy and decurved, matt green above, hairless. Broadly erect shrub, 1.1 × 1.1m in 10 years. Late May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Botswana)

(r) 'Ciscoe Morris'

cv. *Elepidote rhododendron*: 'The Honourable Jean

Marie de Montague' (s) × unknown

**H:** D.A. Griswold (1969), **G:** Griswold Nursery

(1975), **N:** Griswold Nursery (2013), **I:** Griswold

Nursery (2013), **REG:** Griswold Nursery (2013)

Fls 4/lax truss, broadly funnel-shaped, 38 × 51mm, with 5–7 wavy-edged lobes, vivid red (45A) in bud, opening inside light purplish pink (55C), outside deep pink (52B). Lvs oblong, 150 × 38mm, rounded at base, acute at apex, margins upcurved, glossy and moderate olive green (137A) above, hairless. Shrub 1.1 × 1.2m in 40 years. Midseason (Late May in Seattle, Washington, USA).

(Etymology: named for a garden author and television personality in Seattle, Washington, USA)

(v) 'Coco Williams'

cv. *Lepidote rhododendron*: *suaveolens* (s) × *laetum*

**H:** A. Rouse (2002), **G:** A. Rouse (2009), **N:** A. Rouse

(2013), **REG:** A. Rouse (2013)

Fls 7–12/open truss, trumpet-shaped, 45–50 × 40mm, with (4–) 5 very slightly wavy-edged lobes, inside yellowish white (158C), outside with the tube yellowish white to pale yellowish pink (36D); unmarked; not scented. Pedicels to 30mm long, red. Lvs broadly elliptic, 30–70 × 18–38mm, broadly tapering at base, apiculate at apex, margins flat to upcurved, glossy green above, very sparsely scaly. Stiff-branched, spreading shrub, 1.2 × 1.2m in 11 years. No regular flowering period. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a friend of the registrant)

(r) 'Copper Canyon'

cv. *Elepidote rhododendron*: 'Amber Sun' (s) ×

((Jalisco Group × *yakushimanum*) × Joanita Group) × (Goldfort Group × 'Odee Wright')

**H:** J. Barlup (2000), **G:** J. Barlup (2005), **N:** J. Barlup

(2012), **REG:** J. Barlup (2013)

Fls 12/domed truss, open funnel-shaped, 60 × 89mm, with 5 wavy-edged lobes, strong red (53B) in bud, opening inside light orange-yellow (16C), fading to pale orange-yellow (16D) towards the margins, with light orange-yellow (16B) on the dorsal lobe and with a 13mm long, strong red (53C) flare, extending for 38mm above the flare on the 3 dorsal lobes as strong red (53C) spotting; outside pale orange-yellow (16D), with light orange-yellow (16B) on the dorsal lobe fading to pale orange-yellow (16D) on the lobe margin; not scented. Calyx 38mm long, light orange-yellow (16C) with strong red (53C) flaring. Lvs elliptic, 127 × 57mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.0 × 1.2m in 11 years. Midseason (May, in Bellevue, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the name is suggestive of the flower colour)

(r) **'Craig Holmes'**

cv. *Elepidote rhododendron*: 'Gift' (s) × 'Samantha Savers'

**H:** D.J. Dosser (1995), **G:** D.J. Dosser (1999),

**N:** D.J. Dosser, **REG:** D.J. Dosser (2013)

Fls 11/ball truss, open funnel-shaped, with 7 wavy-edged lobes and with partially petaloid stamens, vivid red (57A) in bud, opening deep purplish pink (66C) inside and out, with a diffuse white stripe down the centre of each lobe and strong purplish red (59D) spots on the dorsal lobe; slightly scented. Lvs elliptic, 125 × 40mm, obtuse and mucronate at apex, margins decurved, matt green above, hairless. Shrub 0.9 × 1.2m in 15 years. November (Australia).

(Nomenclatural standard: colour print provided by the registrant (WSY0119346). Etymology: named for a friend of the registrant)

(r) **'Cuyabeno'**

cv. *Elepidote rhododendron*: parentage unknown

**H:** F. Schnupper (2005), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)

Fls 13–14/conical truss, funnel-campanulate, 45–50 × 50–60mm, with 5 flat to wavy-edged lobes, deep purplish pink (73A) in bud, opening light purplish pink (73C) inside, deep purplish pink (73A) outside; with light orange-yellow (22B) spots on the dorsal lobe; dark filaments and style; white anthers. Calyx 2mm long, white. Lvs elliptic, 120–140 × 30–40mm, rounded at base, acute at apex, margins flat, matt green above, hairless. Shrub 1.0 × 1.1m in 10 years. Late May to early June.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Ecuador)

(r) **'Dalmatian'**

cv. *Elepidote rhododendron*: 'Madame Jules Porgès' (s) × 'White Peter'

**H:** J. Doppel (1994), **G:** J. Doppel (1998), **N:** J. Doppel (1998), **REG:** J. Doppel (2013)

Fls 15/ball truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, very pale purple (69C) in bud,

opening inside white, with a slight bluish cast and with a dark purplish red (N79C) blotch and spots occupying most of the dorsal lobe; outside very pale purple (69C), shading to white at margins. Calyx negligible. Lvs elliptic, 125 × 50mm, rounded at base, broadly acute at apex, margins flat, olive green and semi-glossy above, hairless. Shrub 1.2 × 1.5m in 19 years.

Midseason (early May in eastern Pennsylvania, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the blotch is reported to be evocative of the marking of the breed of dogs called Dalmatians)

(r) **'Dawn Star'**

cv. *Elepidote rhododendron*: 'Capistrano' (s) × 'Skipper' **H:** J. Doppel (2005), **G:** J. Doppel (2009), **N:** D. Mac Mullan (2010), **REG:** J. Doppel (2013)

Fls 12/ball truss, broadly funnel-shaped, 50 × 65mm, with 5 wavy-edged lobes, buds moderate reddish orange (171A) at base, shading to moderate yellowish pink (173D) at apex, opening inside strong yellowish pink (37A) in throat, blending to pale yellowish pink (29D) at margins; outside strong yellowish pink (37A) at base, blending to pale yellowish pink (29D) at margins. Calyx negligible. Lvs elliptic, 100 × 40mm, rounded at base, broadly acute at apex, margins slightly downcurved, olive green above, hairless. Shrub 0.75 × 0.75m in 8 years. Midseason (early May in eastern Pennsylvania). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY))

(a) **'Doctor Douglas J. Fraser Junior'**

cv. Evergreen azalea: 'Mrs Nancy Dipple' (s) × 'Satellite' (Klupenger)

**H:** R. Stewart (1997), **G:** R. Stewart (1999),

**N:** R. Stewart (2012), **REG:** R. Stewart (2013)

Fls double, 2/truss, broadly funnel-shaped, 25 × 76mm, with 20 very wavy-edged lobes, strong reddish purple (72C) in bud, opening inside strong reddish purple (72C), with strong purplish red spots on the basal part of the dorsal lobes; outside strong reddish purple (72C). Calyx 10mm long, strong yellow-green. Lvs elliptic, 57 × 25mm, cuneate at base, broadly acute at apex, with upcurved margins, moderate olive green (137A) and semi-glossy above. Shrub 0.9 × 0.9m in 10 years. Midseason.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136119). Etymology: named for the registrant's ophthalmologist. Hybridizer's ref. nos: RBS 96 and 2-97)

(r) **'Dodori'**

cv. *Elepidote rhododendron*: 'Hotei' (s) × Volker Group

**H:** F. Schnupper (2004), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)

Fls hose-in-hose, 11–13/domed truss, campanulate, 50mm wide, with 5–7 wavy-edged lobes, light greenish yellow (8C) and light yellowish pink (38D) in bud, opening inside light greenish yellow (8C), outside light greenish yellow (8C) and light yellowish pink (38D); light yellowish pink (38D) spots on the dorsal lobe; white filaments; style yellowish green, stigma greenish; scented. Calyx 10–20mm long, light greenish yellow (8C) and light yellowish pink (38D).

Lvs elliptic, 60–70 × 20–30mm, cuneate at base, broadly acute at apex, margins flat, matt to glossy green above, with brown hairs below when mature. Dense shrub, 0.5 × 0.8m in 10 years. Mid to late May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Kenya)

(r) **'Doktors Babarikins'**

cv. Elepidote rhododendron: 'Scandinavia' (s) × 'Mrs R.G. Shaw'

**H:** R. Kondratovičs (1974), **G:** R. Kondratovičs (1980), **N:** R. Kondratovičs (2013), **REG:** Rhododendron Nursery Babite, University of Latvia (2013)

Fls 17/strongly domed truss, open funnel-shaped, 90mm wide, with 5 wavy-edged lobes, moderate purplish red (58A) in bud, opening inside very pale purple (69A) in the centre and strong purplish pink (68B) at the margins, outside deep purplish pink (68A); conspicuous blotch of dark red (187A) spots on the dorsal lobes; filaments light purple, anthers grey-yellow; style purple, stigma dark cherry red. Calyx 1mm long, green. Lvs elliptic, 160 × 65mm, pointed at base and apex, margins wavy, matt green above, hairless. Shrub 3 × 4m in 25 years. Early June.

(Nomenclatural standard: colour print provided by the registrant (WSY0119330). Etymology: named for a Dr Babarikin)

(r) **'Dzanga Sangha'**

cv. Elepidote rhododendron: 'Queen Anne's' (s) × 'Capriccio'

**H:** F. Schnupper (2004), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013) Fls 10–12/conical truss, broadly funnel-shaped, 25–30 × 50–60mm, with 5 flat to wavy-edged lobes, and a short, star-shaped ring of separate, regularly formed, petaloid stamens, coloured the same as the corolla, except for a strong purplish pink streak down the centre of each, corolla pale purplish pink (55D) in bud and throughout when open; unmarked. Calyx 1–2mm long, green. Lvs elliptic, 100–110 × 30–40mm, rounded at base, broadly acute at apex, margins flat to wavy, hairless. Dense shrub, 0.9–1.0 × 0.8–0.9m in 10 years.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in the Central African Republic)

(r) **'Dzidra'**

cv. Elepidote rhododendron: 'Catawbiense Grandiflorum' (s) × unknown

**H:** R. Kondratovičs (1978), **G:** R. Kondratovičs (1984), **N:** R. Kondratovičs (2012), **REG:** Rhododendron Nursery Babite, University of Latvia (2013)

Fls 19/strongly domed truss, funnel-shaped, 70mm wide, with 5–7 wavy-edged lobes, light purple (75A) in bud, opening inside very pale purple (69C) in the centre and very pale purple (69B) at the margins, outside very pale purple (69B); strong greenish yellow (151A) spots on the dorsal lobes; filaments and anthers white; style white, with a light yellow stigma. Calyx 1mm long, light green. Lvs elliptic, 160 × 65mm, rounded at base, pointed at apex, margins wavy, matt green above,

hairless. Shrub 1.6 × 2.5m in 25 years. Late May. (Nomenclatural standard: colour print provided by the registrant (WSY0119332). Etymology: a Latvian woman's christian name)

(a) **'Dzige'**

cv. Semi-evergreen azalea: 'Palestrina' (s) × 'Esmeralda' **H:** R. Kondratovičs (1972), **G:** R. Kondratovičs (1976), **N:** R. Kondratovičs (1976), **REG:** Botanical Garden, University of Latvia (2013)

Fls single, 2–3/truss, broadly funnel-shaped, 35 × 40–50mm, with 5 flat to slightly wavy-edged lobes, buds strong purplish red (67A), opening inside very pale purple (76D) in centre, light purple (75A) at margins; outside very pale purple (76D); with deep purplish red (61A), non-contiguous spots over the dorsal lobes; style white, stigma light green. Calyx 3–4mm long, light green. Lvs elliptic, 19–50 × 11–36mm, pointed at base, rounded at apex, margins flat, matt green above, hairless. Shrub 0.7 × 0.8m in 15 years. February to March (under glass).

(Nomenclatural standard: colour print provided by the registrant (WSY0119337). Etymology: derived from the Latvian female name Dzintra Getliņa)

(a) **'Dzirkstelīte'**

cv. Semi-evergreen azalea: 'Adventglöckchen' (s) × 'Esmeralda'

**H:** R. Kondratovičs (1972), **G:** R. Kondratovičs (1975), **N:** R. Kondratovičs (1975), **REG:** Botanical Garden, University of Latvia (2013)

Fls single, 2–4/truss, funnel-shaped, 35 × 40–50mm, with 5 more-or-less flat-edged lobes, deep yellowish pink (47C) in bud, opening strong pink (49A) inside and out, with inconspicuous, non-contiguous, moderate purplish red (58A), dorsal spotting; style white, stigma light green. Calyx 3–4mm long, light green. Lvs obovate, 20–50 × 10–25mm, pointed at base, rounded at apex, margins flat, matt green above, hairless. Shrub 0.8 × 0.8m in 15 years. February to March (under glass).

(Nomenclatural standard: colour print provided by the registrant (WSY0119340). Etymology: Latvian word for "sparklet")

(r) **'Early Winters'**

cv. Elepidote rhododendron: 'Independence Day' (s) × 'Snow Candle'

**H:** J. Barlup (1999), **G:** J. Barlup (2006), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)

Fls 18/ball truss, broadly funnel-shaped, 57 × 83mm, with 5 wavy-edged lobes, light purplish pink (65B) in bud, opening inside white (155D), with a dark red (185A), spotted blotch starting at the base of the dorsal lobe and extending for 45mm; outside white (155D). Calyx insignificant. Lvs elliptic, 210 × 70mm, rounded at base, broadly acute at apex, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.2m in 8 years. Midseason (May in Seattle area, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the Early Winters Spires, twin peaks in the Cascade Mountains (USA))

(r) 'Elise Ann'

cv. *Elepidote rhododendron*: parentage unknown  
**H:** C. O. Dexter (c. 1940), **G:** Heritage Museums and Gardens (1940s), **N:** A. Morrissey (2012), **REG:** Heritage Museums and Gardens (2013)  
Fls 6/lax truss, broadly funnel-shaped, 50 × 82mm, with 7 wavy-edged lobes, buds moderate purplish pink (68C), opening inside very pale purple (69A), with a pale greenish yellow throat, outside pale yellow-green (4D), with subtle, light purplish pink (65B) veining. Lvs elliptic, 114 × 50mm, rounded at base, obtuse at apex, margins flat, matt and moderate olive green (137A) above. Floriferous shrub, 5.5 × 4.5m in 65 years. Late.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: named for Elise Ann Morrissey, daughter of Alyce Morrissey. The latter is a member of the Heritage Museums and Gardens Board of Trustees)

(r) 'Ethan'

cv. *Elepidote rhododendron*: 'Black Magic' (s) × 'Queen Souriya'  
**H:** J.-F. Saint-Jalm (2005), **G:** J.-F. Saint-Jalm (2012), **N:** J.-F. Saint-Jalm (2013), **REG:** J.-F. Saint-Jalm (2013)  
Fls 11/ball truss, open funnel-shaped, 60 × 85mm, with 7 wavy-edged lobes, strong purplish red (61B) in bud, opening inside strong purplish red (71C), fading to strong purplish red (71D), outside strong purplish red (71C); deep purplish red (61A) blotch at base of dorsal throat, with a few spots above; stamens 14, filaments white; style white, glandular, stigma dark red (59A); slightly scented. Calyx insignificant. Lvs elliptic, 110 × 40mm, cuneate at base, broadly acute at apex, margins wavy, hairless. Shrub 1.7 × 0.8m in 7 years. Early to mid May.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: named for the grandson of the registrant)

(a) 'Evelyn Potter'

cv. Evergreen azalea: un-named hybrid (ref. no. CB-1) × 'Elsie Lee' (s) × 'James Stewart'  
**H:** J. F. Klimavicz (2001), **G:** J. F. Klimavicz (2003), **N:** J. F. Klimavicz (2013), **REG:** J. F. Klimavicz (2013)  
Fls 2/truss, double, saucer-shaped, 35 × 70mm, with 15–18 wavy-edged lobes, strong purplish pink (62A) in bud, opening very pale purple (69A) at margins, fading lighter in the centre of the lobes, with some lobes entirely strong purplish pink (62A), with a spotted, strong purplish pink (63C) blotch on the 3 dorsal lobes; outside very pale purple (69A), with occasional strong purplish pink (62A) marks; no stamens; pistil usually deformed. Calyx 10mm long, strong yellow-green (144B). Lvs elliptic, 50 × 20mm, cuneate at base, broadly acute at apex, margins flat, semi-glossy and moderate yellow-green (146B) above. Dense shrub, 0.5 × 0.5m in 5 years. Midseason (May, in northern Virginia, USA).  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: named for a newly born daughter of a business associate of the registrant)

(r) 'Fancy That'

cv. *Elepidote rhododendron*: 'Kera' (s) × 'Christina Dee'  
**H:** J. Barlup (2006), **G:** J. Barlup (2010), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)  
Fls 13/ball truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, deep pink (52B) in bud, opening inside pale yellow (11C) fading to pale yellow (11D) towards margins and with a pale yellowish pink (36D) margin line; 6mm long, strong red (53C) flaring on all lobes, changing to spots and extending 35mm from the base on the dorsal lobes; outside pale yellow (11C), fading to pale yellow (11D) towards the margins; not scented. Calyx insignificant. Lvs elliptic, 108 × 44mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.8m in 5 years. Midseason.  
(Nomenclatural standard: colour print provided by the registrant (WSY).)

(r) 'Farewell Bend'

cv. *Elepidote rhododendron*: 'Mrs Furnivall' (s) × 'One Thousand Butterflies'  
**H:** J. Barlup (1989), **G:** J. Barlup (1995), **N:** J. Barlup (1999), **REG:** J. Barlup (2013)  
Fls 15/domed truss, saucer-shaped, 70 × 102mm, with 6 wavy-edged lobes, deep purplish pink (68A) in bud, opening inside very pale purple (69D) in the centre of the flower, fading to very pale purple (75D) towards the margins, with very light purple (75C) dorsal lobes and moderate red (182A) spotting on the dorsal lobe, extending 48mm from the base; outside very pale purple (75D), with very light purple (75C) mid veins and very light purple (75C) margins; not scented. Calyx insignificant. Lvs elliptic, 127 × 54mm, rounded at base, broadly acute at apex, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.1 × 0.9m in 7 years. Midseason (May, in Bellevue, Washington, USA).  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: named for the Farewell Bend Park, on the Snake River, Oregon, USA)

(v) 'Feria'

cv. *Lepidote rhododendron*: parentage unknown  
**H:** N. Puddey (2008), **G:** N. Puddey (2010), **N:** N. Puddey (2011), **I:** N. Puddey (2012), **REG:** N. Puddey (2013)  
Fls 8–10/truss, funnel-shaped, 65 × 70mm, with 5–7 wavy-edged lobes, buds pale yellow-green (194D), opening both inside and out light yellowish pink (36A), streaked with yellowish white (158D). Calyx less than 1mm long, strong reddish orange (31B). Lvs broadly elliptic, 75 × 45mm, broadly tapering at base, obtuse at apex, margins flat, glossy green above, scales greenish white (155C) when young, aging to dark reddish orange (172B). Shrub 0.6 × 0.4m in 4 years. Irregular flowering period.  
Syn: 'Fiera'  
(Nomenclatural standard: colour print provided by the registrant (WSY0119006).  
Etymology: named after a Spanish festival in Seville; the short pedicels and wavy-edged lobes giving the inflorescence the impression of a Spanish dancers ruffled dress. Originally introduced with the incorrect spelling 'Fiera')

(v) **'Fiera'**  
Syn. of 'Feria'

(r) **'Feuer und Flamme'**

cv. Elepidote rhododendron: *dichroanthum* subsp. *scyphocalyx* (s) × *yakushimanum* 'Koichiro Wada'  
**H:** F. Schnupper (2004), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls 10/domed to lax truss, funnel-shaped, 40–50 × 40–50mm, with 5 flat to wavy-edged lobes, bicoloured, strong purplish red (58C) on the lobes, shading to brilliant yellow (21C) in the throat (described as pink to orange), with moderate orange-yellow (168D) spots on the dorsal lobe. Calyx 5mm long, yellow-green. Lvs elliptic, 60–80 × 25–35mm, rounded at base, broadly acute at apex, margins flat to wavy, matt green above. Dense shrub, 0.9 × 0.9m in 10 years. Flowering over a long period from early to mid May.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: named for the colours of the flowers. The seed parent is reported to have hairless leaves and thus requires confirmation)

(r) **'Frosted Lemon'**

cv. Elepidote rhododendron: 'Scintillation' (s) × 'Recital'  
**H:** J. Barlup (1996), **G:** J. Barlup (2001), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)  
Fls 19/ball truss, broadly funnel-shaped, 44 × 70mm, with 6 very wavy-edged lobes, brilliant greenish yellow (4A) in bud, opening inside white (155D), blending to light greenish yellow (4C) on the dorsal lobes, with brilliant yellow-green (154C) rays on the dorsal lobe extending 13mm from the base and a moderate red (185B) flare on the dorsal lobe changing to brilliant yellowish green (154C) with age; outside white (155D), blending to light greenish yellow (4C) on the dorsal lobes; not scented. Calyx 25mm long, white (155D) and brilliant yellowish green (154C). Lvs elliptic, 108 × 51mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.9m in 7 years. Midseason (May, in Bellevue, Washington, USA).  
(Nomenclatural standard: colour print provided by the registrant (WSY0136172).  
Etymology: the name is suggestive of the flower colour)

(r) **'Funny Mask'**

cv. Elepidote rhododendron: 'Mondnacht' (s) × 'Hachkaba'  
**H:** K.-H. Hübbers (2001), **G:** K.-H. Hübbers (2008), **N:** K.-H. Hübbers (2013), **REG:** J. Westhoff (2013)  
Fls c.12/rounded, domed truss, open saucer-shaped, 75mm wide, with 5 wavy-edged, recurved lobes, strong purplish red (63B) in bud, opening inside light to pale purplish pink (62C/D), outside moderate to light purplish pink (65A/B); a 3-lobed, dark red to moderate purplish red (59A-C), dorsal blotch, extends onto the adjacent lobes and breaks up into dots. Calyx 5mm long, light green. Lvs ovate-oblong, 100 × 50mm, tapered at base, broadly acute and mucronate at apex, margins flat, matt green and slightly rugose above, with tiny brown hairs near the midrib below when

young. Shrub 0.6 × 0.7m in 5 years. Early to mid May.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136171).  
Etymology: the flower markings considered reminiscent of the smirking smile on a Guy Fawkes mask. The pollen parent is better known under the trade designation KABARETT or HACHMANN'S KABARETT)

(r) **'Garden Show'**

cv. Elepidote rhododendron: un-named yellow-orange seedling (s) × 'Pucine'  
**H:** K.-H. Hübbers (1999), **G:** K.-H. Hübbers (2012), **N:** K.-H. Hübbers (2012), **REG:** J. Westhoff (2013)  
Fls up to 9/domed truss, open funnel-shaped, 110mm wide, with 7 wavy-edged lobes, buds light yellow (10C), opening inside brilliant yellow-green (154C), outside light yellow (12C); with faint, strong greenish yellow (151A) spots on the 3 dorsal lobes (except at the margins). Calyx 5mm long, green, with a dark red rim. Lvs oblong, 170 × 70mm, tapered at base, acute at apex, margins slightly upcurved, matt green above, hairless. Shrub 1.5 × 1.0m in 10 years. Mid May.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136170))

(v) **'Golden Dunes'**

cv. Elepidote rhododendron: 'Carillon Bells' (s) × 'Desert Song'  
**H:** N. Puddey (2008), **G:** N. Puddey (2010), **N:** N. Puddey (2011), **I:** N. Puddey (2012), **REG:** N. Puddey (2013)  
Fls 8–10/open truss, tubular funnel-shaped, 50 × 20mm, with 5 flat-edged lobes, moderate reddish brown (174A) in bud, opening brilliant orange-yellow (23B) throughout, with strong orange (26A) along the basal margins of the lobes. Calyx less than 1mm long, light greenish yellow. Lvs broadly elliptic, 50 × 20mm, broadly tapering at base, acute at apex, margins decurved, glossy green above, with greenish white (155C) scales when young (scales not visible with age). Shrub 1.0 × 0.5m in 4 years. Irregular flowering period.  
(Nomenclatural standard: colour print provided by the registrant (WSY0119046).  
Etymology: flower colour reminiscent of the registrant's local sands dunes)

(r) **'Grace Louise'**

cv. Elepidote rhododendron: parentage unknown  
**H:** J.C. Cowles (c. 1960), **G:** Heritage Museums and Gardens (1960s), **N:** S.J. Hamilton (2013), **REG:** Heritage Museums and Gardens (2013)  
Fls 5/lax truss, open funnel-shaped, 60 × 90mm, with 7 wavy-edged lobes, moderate yellowish pink (38B) in bud, opening inside pale greenish yellow (2D), with 2 light yellow-green flares on the dorsal lobe, extending 25mm from the base over a light salmon throat; outside pinkish white (N155B). Calyx insignificant. Lvs elliptic, 82 × 64mm, rounded at base, acute at apex, margins downcurved, matt and dark green (136A) above, hairless. Shrub 3.6 × 3.0m in 40 years. Midseason (May on Cape Cod, Massachusetts, USA).  
(Nomenclatural standard: colour print of an electronic image (WSY0136169).  
Etymology: named for Grace Louise Cooke, mother of Susan J. Hamilton, the

latter a Trustee and supporter of the Heritage Museums and Gardens (Sandwich, Massachusetts, USA). Heritage ref. no. 134-2002 (143-02); Sandwich Club ref. no. JG-1)

(a) ‘**Grācija**’

cv. Evergreen azalea: sport from ‘Schloss Pillnitz’  
**G:** R. Kondratovičs (1989), **N:** R. Kondratovičs (1989),  
**REG:** Botanical Garden, University of Latvia (2013)  
Fls single, 2–4/truss, funnel-shaped, 30 × 50mm, with 5 flat-edged lobes, buds white (155C), opening white (155C) throughout; unmarked; style white, stigma light green. Calyx 3mm long, light green. Lvs elliptic, 25 × 12mm, pointed at base, rounded at apex, margins flat, glossy green above, hairless. Shrub 0.7 × 0.7m in 15 years. February to March (under glass). (Nomenclatural standard: colour print provided by the registrant (WSY0119339). Etymology: Latvian female christian name)

(v) ‘**Graham’s Gift**’

cv. *Lepidote rhododendron*: ‘Neesa’ (s) × *konori* var. *phaeocephalum*  
**H:** G. L. S. Snell (2004), **G:** N. Puddey (2010),  
**N:** N. Puddey & W. Snell, **I:** N. Puddey (2012),  
**REG:** N. Puddey (2013)  
Fls 5–8/open truss, tubular funnel-shaped, 80 × 75mm, with 5–6 flat-edged lobes, buds dark red (187A), opening inside white (155B), outside white (155B) deepening to pale yellow (11D) along the tube; with a narrow pink suffusion evident on both the inside and outside at the base of the lobe edges; slightly scented. Calyx less than 1mm long, dark yellowish pink (181D). Lvs broadly elliptic, 100 × 50mm, broadly tapering at base, obtuse at apex, with wavy margins, glossy green above, scales pale yellow (158B) at first, aging to moderate reddish brown (174A). Upright shrub, 1.2 × 0.6m in 8 years. Flowering irregularly, 2–3 times a year.  
(Nomenclatural standard: colour print provided by the registrant (WSY0119040). Etymology: named in honour of the breeder, the late Graham Snell)

(r) ‘**Great Bear**’

cv. *Elepidote rhododendron*: *brachycarpum* (s) × unknown  
**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls hose-in-hose, 11–12/conical truss, funnel-campanulate, 35–40 × 50mm, with 5 flat lobes, pink in bud, opening white throughout, with light greenish yellow (3D) spots on the 3 dorsal lobes; filaments white; style white, stigma yellow. Calyx 10–15mm long, white. Lvs elliptic, 70–100 × 30–40mm, cuneate at base, broadly acute at apex, margins flat and upcurved, matt green above, indumentum white when young, aging to brown. Dense shrub, 0.5 × 0.7m in 10 years. Early May.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the Great Bear Rainforest in Canada)

(r) ‘**Hamburger Deern**’

cv. *Elepidote rhododendron*: ‘Mooreule’ × ‘Mrs T.H. Lowinsky’ (s) × ‘Extraordinaire’  
**H:** H. Hesse (2003), **G:** H. Hesse (2008),  
**N:** H. Hesse, **REG:** H. Hesse (2013)  
Fls 6–11/full, rounded truss, broadly funnel-shaped, 45–65 × 65–100mm, with 5–7 flat-edged, reflexed lobes, dark pink in bud, opening inside strong yellow, with a pale pink margin, outside matt pink, the basal third matt yellowish pink; striking dark red markings on the 3 dorsal lobes. Calyx 1–2mm long, dark green. Lvs elliptic, 51–120 × 28–50mm, rounded at apex, margins flat to decurved, matt green above, hairless. Shrub 1.25 × 0.7m in 10 years. Mid May.

(a) ‘**Hampton Ruby Red**’

cv. Evergreen azalea: (‘Ripples’ × ‘Girard’s Hot Shot’) × ‘Girard’s Hot Shot’ (s) × ‘Karafune’  
**H:** S.F. McDonald (1992), **G:** S.F. McDonald (1995),  
**N:** S.F. McDonald (2013), **REG:** S.F. McDonald (2013)  
Fls nearly fully double, open funnel-shaped, 22 × 35mm, with typically more-or-less 15 flat-edged lobes, strong red (46A) in bud, opening inside vivid red (45A), with a few, inconspicuous, strong red (46A) spots; outside vivid red (45B); occasional stamens and pistil present. Calyx 4mm long, strong yellow-green (145A). Lvs obovate, 20 × 8mm, cuneate at base, apiculate at apex, margins flat, glossy and moderate olive green (137A) above, with sparse, pale beige hairs above and below. Shrub with a dense habit, 1.2 × 0.7m in 20 years. Midseason (late April to early May, at south end of Chesapeake Bay, Virginia, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136168). Etymology: Hampton is where the registrant lives)

(v) ‘**Happy Cam**’

cv. *Lepidote rhododendron*: ‘Kisses’, selfed (s) × ‘Beejay Bay’  
**H:** N. Puddey (2008), **G:** N. Puddey (2011),  
**N:** N. Puddey (2011), **I:** N. Puddey (2012),  
**REG:** N. Puddey (2013)  
Fls 5/open truss, funnel-shaped, 65 × 55mm, with 5 flat-edged lobes, pale yellow-green (193B) in bud, opening inside vivid reddish orange (30A), with light yellow (21D) in the tube, outside vivid reddish orange (30A) fading to light yellow (21D) at the base of the lobe and on the tube; unmarked. Calyx less than 1mm long, light greenish yellow (1C). Lvs broadly elliptic, 115 × 55mm, narrowly tapering at base, obtuse at apex, margins decurved, glossy green above, scales greenish white (192D) at first, lost when mature. Shrub 1.0 × 0.5m in 3 years. Flowering irregularly.  
(Nomenclatural standard: colour print provided by the registrant (WSY0119044). Etymology: named for Cameron, the son of the registrant, and described as a ‘happy person’)

(r) ‘**Hell’s Gate**’

cv. *Elepidote rhododendron*: *williamsianum* × *orbiculare* (s) × ‘Ring of Fire’  
**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls 5–7/lax truss, funnel-campanulate, 25–30 ×

30–40mm, with 5 flat to wavy-edged lobes, strong purplish red (58B) in bud, opening strong purplish red (58C) on the margins of the lobes and brilliant yellow (21C) in the throat and base of the lobes – both inside and out, with slight greenish spotting on the dorsal lobe; anthers white; stigma green. Calyx 1mm long, green. Conspicuous, long, pale pedicels, held so the flowers are out or slightly down-facing. Lvs orbicular, 60–70 × 30–40mm, cordate at base, broadly acute at apex, margins flat, glossy green above, hairless. Shrub 0.6 × 1.0m in 10 years. Early May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Kenya)

(r) **'Honey Gold'**

cv. *Elepidote rhododendron*: 'Percy Wiseman' (s) × 'Orange Prelude'

**H:** J. Barlup (2006), **G:** J. Barlup (2011), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)

Fls 14/domed truss, broadly funnel-shaped, 57 × 83mm, with 5 wavy-edged lobes, strong red (53C) in bud, opening inside pale yellow (11C), blending to moderate yellowish pink (38B), with deep pink (47D) lobe tips and with 2 strong yellow (153D) flares on the dorsal lobe, extending 19mm from the base; outside pale yellow (11C), blending to moderate yellowish pink (38B), with strong yellowish pink (38A) midveins and deep pink (47D) lobe tips. Calyx 41mm long, moderate yellowish pink (38B), with strong yellow (153D) flaring. Lvs elliptic, 127 × 54mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.7 × 0.9m in 7 years. Midseason (May in the Seattle area, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136162))

(r) **'Howard J'**

cv. *Elepidote rhododendron*: parentage unknown

**H:** J.C. Cowles (1960s), **G:** Heritage Museums and Gardens (1960s), **N:** D. McGraw (2012),

**REG:** Heritage Museums and Gardens (2013)

Fls 12/lax truss, open funnel-shaped, with 5 wavy-edged lobes, vivid purplish red (61C) in bud, opening inside pale orange-yellow (159B) at base of corolla and in centre of each lobe, blending to strong purplish red (63B) at margins, with brownish orange to moderate orange-yellow (164A-B) twin rays on the 3 dorsal lobes extending 25mm from the base; outside vivid purplish red to deep purplish pink (61C-D); stamens rudimentary. Calyx insignificant. Lvs elliptic, 90 × 38mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and dark yellowish green (139A) above, hairless. Shrub 2.0 × 2.5m in 25 years. Late. Syn: 'Cemetery Coral'

(Nomenclatural standard: colour print of an electronic image provided with the registration (WSY0136161). Etymology: named for Howie McGlynn, father-in-law of the nominant David McGraw (the latter a member of the Heritage Museums and Gardens Board of Trustees). Also referred to under the numbers: 46-1994 and 46-94)

(r) **'Ilka'**

cv. *Elepidote rhododendron*: 'Berliner Liebe' (s) × unknown

**H:** F. Schnupper (1990), **G:** F. Schnupper, **N:** F.

Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)

Fls 14–15/conical truss, broadly funnel-shaped, 25–40 × 60mm, with 5 wavy-edged lobes, deep purplish pink (68A) in bud, opening inside moderate purplish pink (68C), outside strong purplish pink (68B); with black nectar pouches; filaments and style pale purplish pink. Calyx 2mm long, green. Lvs elliptic, 100–120 × 40mm, rounded at base, broadly acute at apex, margins flat and decurved, glossy green above, hairless. Dense shrub, 1.1 × 1.3m. Mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the registrant's wife, Ilka Schnupper)

(r) **'Indriķis'**

cv. *Elepidote rhododendron*: 'Carawbiense Grandiflorum' (s) × unknown

**H:** R. Kondratovičs (1975), **G:** R. Kondratovičs (1981), **N:** R. Kondratovičs (2013), **REG:** Rhododendron

Nursery Babite, University of Latvia (2013)

Fls 19/strongly domed truss, open funnel-shaped, 75 × 70mm, with 5 very wavy-edged lobes, deep purple (N81A) in bud, opening inside very light purple (76B) in the centre, light purple (76A) at the margins, outside light purple (75A); conspicuous greyish purple (N77A) markings on the dorsal lobe (some of which show through to the outside); filaments white, anthers light violet; style white, stigma dark cerise. Calyx 1mm long, light green. Lvs elliptic, 135 × 55mm, rounded at base, pointed at apex, margins flat, matt green above, hairless. Shrub 1.5 × 2.5m in 25 years. Early June.

(Nomenclatural standard: colour print provided by the registrant (WSY0119334). Etymology: a Latvian male christian name)

(r) **'Invitation's Gift'**

cv. *Elepidote rhododendron*: 'Invitation' (s) × (*yakushimanum* × *lacteum*) × 'Powder Snow'

**H:** J. Barlup (2000), **G:** J. Barlup (2006), **N:** J. Barlup (2009), **REG:** J. Barlup (2013)

Fls 18/domed truss, funnel-campanulate, 38 × 51mm, with 5 wavy-edged lobes, light greenish yellow (4B) in bud, opening inside pale greenish yellow (2D), with the apical half of the lobes light yellow-green (2C), and with a semicircle of strong red (46A) at the base of the dorsal throat, extending 10mm from the base; outside pale greenish yellow (2D), with dorsal lobes light yellow-green (2C); not scented. Calyx insignificant. Lvs elliptic, 102 × 44mm, rounded at base, broadly acute at apex, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.8 × 1.2m in 11 years. Midseason (April, in Bellevue, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136160). Etymology: derived from the name of the seed parent)

(r) **'Ivory Dream'**

cv. *Elepidote rhododendron*: *yakushimanum* × *lacteum* (s) × *yakushimanum* × *macabeanum*

**H:** J. Barlup (1992), **G:** J. Barlup (1998), **N:** J. Barlup

(2011), **REG:** J. Barlup (2013)

Fls 18/domed truss, funnel-campanulate, 44 × 57mm, with 6 slightly wavy-edged lobes, deep pink (48B) in bud, opening inside pale yellow-green (4D), with light greenish yellow (4C) ridges on each segment, the apical 13mm of lobes white (155D) and with a moderate red (181C), semicircular dorsal blotch, extending 10mm from the base; outside pale yellow-green (4D) at base, changing to white (155D) at the tip of the lobes; not scented. Calyx insignificant. Lvs elliptic, 120 × 57mm, rounded at base, broadly acute at apex, margins downcurved, moderate olive green (147A) above. Shrub 0.9 × 1.4m in 19 years. Early (April, in Bellevue, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136159)).  
Etymology: name suggestive of the flower colour)

(r) '**Jeremiah William**'

cv. *Elepidote rhododendron*: parentage unknown  
**H:** C.O. Dexter (c. 1930), **G:** C.O. Dexter (1930s),  
**N:** Mrs J.M. O'Connor (2013), **REG:** Heritage  
Museums and Gardens (2013)

Fls 10/ball truss, broadly funnel-shaped, 51 × 89mm, with 7 wavy-edged lobes, buds deep purplish pink (N57D), opening yellowish white (N155D) in the centre, blending to deep purplish pink (N74C) at the margins, with twin deep purplish red (71A) flares on the dorsal lobe; outside deep purplish pink (N74C), with prominent central veins on each lobe; stamens 13mm long; pistil 51mm long, prominent. Calyx insignificant. Lvs obovate, 127 × 44mm, rounded at base, acute at apex, margins flat, matt and moderate olive green (137A) above, hairless. Shrub with an open habit, 2.5–3.0 × 3.0m wide in 40 years.

Midseason (May on Cape Cod, Massachusetts, USA).  
(Nomenclatural standard: colour print of an electronic image (WSY0136158)). Etymology: named for Jeremiah William O'Connor, the late husband of Joan O'Connor, the latter a Trustee and supporter of the Heritage Museums and Gardens)

(v) '**Joan McClelland**'

cv. *Lepidote rhododendron*: *viriosum* (s) ×  
*rarelepidotum*

**H:** A. Rouse (2001), **G:** A. Rouse (2008), **N:** A. Rouse (2013), **REG:** A. Rouse (2013)

Fls 10–15/open truss, campanulate, 40 × 45mm, with 5 very slightly wavy-edged lobes, strong reddish orange (31A) towards the base, strong reddish orange (32B) throughout the rest of the corolla; unmarked. Calyx less than 1mm long, strong reddish orange (32B). Pedicels c.40mm long, vivid reddish orange (32A), some acutely angled at point of attachment to the flower. Lvs broadly elliptic, 70–120 × 32–57mm, broadly tapering at the base, shortly acuminate to obtuse at the apex, margins wavy-edged, mostly upturned, glossy green above, very sparsely scaly on underside of leaves and on new growth. Shrub 1.2 × 1.0m in 12 years. Flowering irregularly.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136157)).

Etymology: named for a friend of the registrant on the occasion of her 100th birthday)

(r) '**Joanne Holmes**'

cv. *Elepidote rhododendron*: parentage unknown

**H:** D.J. Dosser, **G:** D.J. Dosser, **N:** D.J. Dosser,

**REG:** D.J. Dosser (2013)

Fls 8/ball truss, broadly funnel-shaped, 115mm wide, with 7 slightly wavy-edged lobes, strong purplish red (64C) in bud, opening inside moderate purplish pink (65A), outside strong purplish red (63A), with a moderate purplish red (64A) marking all around the base of the throat. Lvs 115 × 40mm, truncate at base, acute at apex, margins decurved, matt green above, hairless. Shrub 2 × 2m in 10 years. October to November (Australia).

(Nomenclatural standard: colour print provided by the registrant (WSY0119350)). Etymology: named for a friend of the registrant)

(r) '**John Kinssies's Waterfall**'

cv. *Elepidote rhododendron*: 'The Honourable Jean Marie de Montague' (s) × 'Suzy Lou'

**H:** D.A. Griswold (1969), **G:** Griswold Nursery (1987),

**N:** Griswold Nursery (2013), **I:** Griswold Nursery

(2013), **REG:** Griswold Nursery (2013)

Fls 10/domed truss, funnel-campanulate, 38 × 51mm, with 5 wavy-edged lobes, vivid red (45A) in bud, opening inside light purplish pink (55C), becoming paler toward the centre, with a small, pale orange-yellow (159C) blotch in the centre of the dorsal lobe; outside deep pink (52B). Calyx 38mm long, split into 2 recurved segments held over the dorsal side of the corolla, pale orange-yellow (159C); no apparent calyx around the rest of the corolla. Lvs lanceolate, 100 × 38mm, cuneate at base, acuminate at apex, with wavy, decurved margins, matt and moderate yellow-green (146B) above, hairless. Shrub with an open habit, 1.5 × 1.2m in 40 years. Midseason.

(Etymology: named for a noted landscape designer in Seattle, Washington, USA)

(r) '**Joker's Wild**'

cv. *Elepidote rhododendron*: 'Sun Stream' (s) ×  
'Hill's Low Red'

**H:** J. Barlup (2005), **G:** J. Barlup (2012), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)

Fls 17/ball truss, broadly funnel-shaped, 50 × 83mm, with 6 wavy-edged lobes, strong red (53B) in bud, opening inside pale orange-yellow (16D), blending to vivid red (52A) towards the margins, with 9–12mm long, strong red (53C) nectar pouches and two 13mm long, dark red (187A) flares on the dorsal lobe; outside strong red (53B), blending though deep pink (52B) to vivid red (52A) on the edges. Calyx 38mm long, pale orange-yellow (16D) changing to vivid red (52A), with a 13mm long, strong red (53C) flare. Lvs elliptic, 102 × 53mm, rounded at base, broadly acute at apex, margins downcurved, matt and moderate olive green (147A) above, hairless. Shrub 0.7 × 0.8m in 8 years. Midseason (May, in the Seattle area, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136156))

(v) **'Joy Lenore'**

cv. *Lepidote rhododendron*: 'Kisses' (s) × 'Charming Valentino'

**H**: N. Puddey (2008), **G**: N. Puddey (2009),

**N**: N. Puddey (2010), **I**: N. Puddey (2012),

**REG**: N. Puddey (2013)

Fls 8–10/domed truss, funnel-shaped, 45 × 50mm, with 5 flat-edged lobes, pale yellow-green (194D) in bud, opening light yellowish pink (38C) and fading to light yellowish pink (27A) both inside and out; unmarked. Calyx less than 1mm long, strong yellowish pink (33C). Bract margins dark reddish orange (173A). Lvs broadly elliptic, 60 × 35mm, rounded at base, obtuse at apex, margins decurved, glossy green above, scales white (155C) at first, not evident later. Shrub 1.0 × 0.5m in 4 years. Flowering irregularly, 2–3 times a year. (Nomenclatural standard: colour print provided by the registrant (WSY0119010). Etymology: named for the sister of the registrant)

(r) **'Judy Judy Judy'**

cv. *Lepidote rhododendron*: 'Peach Charm' (s) × 'Pirouette'

**H**: J. Barlup (1992), **G**: J. Barlup (1998),

**N**: R. Blackmore (2012), **REG**: J. Barlup (2013)

Fls 21/ball truss, broadly funnel-shaped, 57 × 95mm, with 5 wavy-edged lobes, strong pink (52D) in bud, opening inside pale purplish pink (62D), fading soon after opening to white (155D), with 2 pale greenish yellow (2D) flares extending 9mm from the base; outside pale purplish pink (62D), fading to white (155D). Calyx insignificant. Lvs elliptic, 146 × 48mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Dense shrub, 0.5 × 0.7m in 6 years. Midseason (May, in the Seattle area, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136155).)

Etymology: named for the registrant's wife and evoking the apocryphal story that Cary Grant uttered "Judy Judy Judy" in the 1939 movie *Only Angels Have Wings*)

(r) **'Judy's Choice'**

cv. *Lepidote rhododendron*: 'Hachmann's Charmant' (s) × 'Tia'

**H**: J. Barlup (2002), **G**: J. Barlup (2006), **N**: J. Barlup (2009), **REG**: J. Barlup (2013)

Fls 13/ball truss, broadly funnel-shaped, 51 × 73mm, with 5 wavy-edged lobes, moderate purplish pink (65A) in bud, opening pale yellow (8D) at the base, extending up for 13mm from the base and blending into white (155D) and with a deep yellowish pink (47C) flare on the dorsal lobes, extending 38mm from the base, outside pale yellow (8D) at the base, extending up 13mm from the base, then blending into moderate purplish pink (65A), changing to white (155D); not scented. Calyx insignificant. Lvs elliptic, 140 × 57mm, rounded at base, broadly acute at apex, margins upcurved, dull and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.2m in 10 years. Midseason (April, in Bellevue, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136154))

(r) **'Kaza'**

cv. *Elepidote rhododendron*: *yakushmanum*

'Koichiro Wada' (s) × 'Manderley'

**H**: F. Schnupper (2004), **G**: F. Schnupper, **N**: F.

Schnupper, **I**: F. Schnupper, **REG**: F. Schnupper (2013)

Fls 8–9/domed truss, funnel-campanulate, 40–50 × 50–60mm, with 5 flat-edged lobes, deep pink (52C) in bud, opening inside light greenish yellow (8C) to white, outside deep pink (52C) to pale yellowish pink (36D); moderate orange (170C) spots on the dorsal lobe. Calyx 2–3mm long, with a grey indumentum. Lvs elliptic, 80–100 × 30mm, cuneate at base, acute at apex, margins flat and decurved, glossy green above, indumentum fawn when young, aging to brown. Shrub 0.7 × 0.9m in 10 years. Mid May. (Etymology: named for the transfrontier Kavango-Zambezi [Kaza] national park (covering parts of Botswana, Angola, Namibia, Zambia and Zimbabwe))

(v) **'Kerry Gold'**

cv. *Lepidote rhododendron*: 'Great Scent-sation' (s) × 'Frosted Candy'

**H**: E. Lyon (2004), **G**: E. Lyon (2007), **N**: E. Lyon

(2013), **REG**: E. Lyon (2013)

Fls 8/open-topped truss, tubular funnel-shaped, 80 × 70mm, with 5–6 flat-edged lobes, cream in bud, opening rich cream inside and yellow cream outside; unmarked; prominent orange anthers. Lvs oblanceolate, 95 × 35mm, attenuate at base, acuminate at apex, margins wavy, matt green above, with no evident scales. Shrub 1.2 × 0.6m in 9 years. Flowering all year round. (Nomenclatural standard: colour print provided by the registrant (WSY0119353))

(r) **'Kranenlicht'**

cv. *Elepidote rhododendron*: 'Lem's Cameo' (s) × 'Deenah'

**H**: K.-H. Hübbers (1992), **G**: K.-H. Hübbers (1999),

**N**: K.-H. Hübbers (2013), **REG**: J. Westhoff (2013)

Fls up to 12/loose, flattened ball truss, funnel-shaped, 50 × 70mm wide, with 5 wavy-edged lobes, opening inside strong purplish pink (63C), shading paler towards the centre, outside deep to pale purplish pink (55A-D); a small, vivid to strong reddish orange (42A/B) blotch in the centre of the corolla. Calyx 2–3mm long, greenish, with a small red rim. Lvs oblong, 130 × 55mm, tapered at base, apiculate at apex, margins flat, matt green above, with a few brownish hairs below when young. Shrub 1.0 × 1.1m in 10 years. Early to mid May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136153) Etymology: named in part for the city where it was raised (Kranenburg, Germany) )

(r) **'Kui Buri'**

cv. *Elepidote rhododendron*: *yakushmanum*

'Koichiro Wada' (s) × unknown

**H**: F. Schnupper (2004), **G**: F. Schnupper, **N**: F.

Schnupper, **I**: F. Schnupper, **REG**: F. Schnupper (2013)

Fls 8–10/conical truss, open funnel-shaped, 40–45 × 60mm, with 5 wavy-edged lobes, vivid purple (81B) in bud opening light purple (81C) to white inside, light purple (81C) outside; yellow-green spots on the

dorsal lobe; filaments and style more-or-less white, anthers very pale. Calyx 2mm long, green. Lvs elliptic, 100–120 × 30–40mm, cuneate at base, broadly acute at apex, margins flat and decurved, matt green above. Dense shrub, 0.8 × 1.0m in 10 years. Mid May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Thailand)

(r) **'Lady Bug'**

cv. *Elepidote rhododendron*: 'Terra' (s) × 'Berg's Yellow' × *proteoides*  
**H:** J. Barlup (2004), **G:** J. Barlup (2009), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)  
Fls 20/ball truss, open funnel-shaped, 51 × 70mm, with 5 wavy-edged lobes, light greenish yellow (4B) in bud, opening inside pale yellow-green (4D), deeper on the dorsal lobe (light greenish yellow, 4C), and with a strong red (53B) flare on the dorsal lobe, which extends 17mm from the base and is surmounted by strong red (53B) spotting which reaches 38mm from the base; outside pale yellow-green (4D), deeper on the dorsal lobes (light greenish yellow (4C)); not scented. Calyx 10mm long, light greenish yellow (4B), with strong red (53B) flaring. Lvs elliptic, 83 × 38mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A), hairless. Dense shrub, 0.6 × 0.75m in 8 years. Midseason (April, in Bellevue, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136151))

(a) **'Laima'**

cv. Semi-evergreen azalea: 'Adventglöckchen' (s) × 'Esmeralda'  
**H:** R. Kondratovičs (1972), **G:** R. Kondratovičs (1975), **N:** R. Kondratovičs (1975), **REG:** Botanical Garden, University of Latvia (2013)  
Fls single, 2–4/truss, funnel-shaped, 40 × 47–60mm, with 5 flat-edged lobes, white (155A) in bud, opening white (155C) inside and out; with moderate red (185B), non-contiguous, dorsal spots; style white, stigma light pink. Calyx 3–6mm long, light green. Lvs elliptic, 24–53 × 12–25mm, pointed at base, rounded at apex, margins flat, matt green above, hairless. Shrub 1.2 × 1.3m in 15 years. February to March (under glass). (Nomenclatural standard: colour print provided by the registrant (WSY0119342). Etymology: Latvian female christian name)

(r) **'Late Lady'**

cv. *Elepidote rhododendron*: 'Admiral Vanessa' (s) × 'Welse'  
**H:** K.-H. Hübbbers (2000), **G:** K.-H. Hübbbers (2009), **N:** K.-H. Hübbbers (2013), **REG:** J. Westhoff (2013)  
Fls 11/flat-domed truss, funnel-shaped, 50 × 60mm, with 6 wavy-edged lobes, buds vivid purplish red (61C), opening inside pale purplish pink (62D), outside strong purplish pink (63C), the lobes pale purplish pink (62D); dotted, moderate purplish red (59C) blotch on dorsal lobe and half of each adjacent lobe. Calyx 1mm long, greenish. Lvs oblong, 100 × 45mm, rounded at base, apiculate at apex, rugose and

matt green above, hairless. Shrub 1.0 × 0.7m in 10 years. June.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136150). Etymology: named to mark the late flowering time)

(a) **'Latvijas Itera'**

cv. Deciduous azalea: un-named Exbury (i.e. Knapp Hill) seedling (Ex-5) (s) × 'Uguns'  
**H:** R. Kondratovičs (1976), **G:** R. Kondratovičs (1983), **N:** R. Kondratovičs (2002), **REG:** Rhododendron Nursery Babīte, University of Latvia (2013)  
Fls single, 9/slightly domed truss, tubular funnel-shaped, 80 × 70mm, with 5 wavy-edged lobes, strong red (41B) in bud, opening inside strong reddish orange (31A) in the centre, strong red (41B) at the margins; outside strong red (41B); vivid orange-yellow (23A) blotch on the dorsal lobes; filaments red, anthers cream; style red, stigma green. Calyx 2mm long, green. Lvs elliptic, 110 × 35mm, pointed at base and apex, margins wavy, glossy green above, hairless. Shrub 2 × 2m in 20 years. Late May to early June. (Nomenclatural standard: colour print provided by the registrant (WSY0119329). Etymology: Itera is the name of a gas supply company in Latvia)

(r) **'Lemon Prelude'**

cv. *Elepidote rhododendron*: 'Recital' (s) × 'Bud's Yellow'  
**H:** J. Barlup (2000), **G:** J. Barlup (2005), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)  
Fls 13/domed truss, 48 × 73mm, with 5 wavy-edged lobes, strong red (51A) in bud, opening inside pale yellow (18C), with light yellow (18A) midveins and dorsal lobes, and with twin, deep red (53A) flares on the dorsal lobes, with spotting extending 13mm above the flares; outside pale yellow (18C), with light yellow (18A) midveins and dorsal lobes; not scented. Calyx 17mm long, light yellow (18A), with deep red (53A) spotting. Lvs elliptic, 95 × 51mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 11 years. Midseason (May, in Bellevue, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136149))

(a) **'Li'l Sister'**

cv. Evergreen azalea: 'Elsie Lee' × 'Dorsett' (s) × 'Helen Curtis'  
**H:** W.L. Clagett (1996), **G:** W.L. Clagett (2001), **N:** W.L. Clagett (2013), **I:** White's Nursery (2005), **REG:** W.L. Clagett (2013)  
Fls semi-double, 3/truss, broadly funnel-shaped, 32 × 44mm, with 5 + 5–7 wavy-edged lobes, strong purplish red (63A) in bud, opening inside strong purplish red (63A), with moderate purplish red (70A) spotting on the dorsal lobes; stamens rare; outside strong purplish red (63A). Lvs elliptic, 35 × 13mm, cuneate at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Very floriferous shrub, 1.2 × 0.8m in 17 years. Early (April in Washington, D.C., USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant

(WSY0136148). Etymology: named for the registrant's youngest sister. Originally introduced by White's Nursery under the ref. no. 96-2-12)

(r) '**Łobéké**'

cv. Elepidote rhododendron: *smirnowii* (s) × unknown  
**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls 12/conical truss, open funnel-shaped, 40–50 × 60–80mm, with 5–7 wavy-edged lobes, strong reddish purple (72C) in bud, opening inside and out deep purplish pink (72D); with a dark greenish yellow (152C) blotch on the dorsal lobe; filaments white to pale pink, anthers white; style purplish pink, darker at apex. Calyx 1–2mm long, green. Lvs elliptic, 120–200 × 40–80mm, rounded at base, broadly acute at apex, margins flat to wavy, upcurved, glossy green above, hairless; new foliage yellowish. Dense shrub, 1.2 × 1.2m in 10 years. Mid to late May.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Cameroon)

(r) '**Lois Blackmore**'

cv. Elepidote rhododendron: 'Fantastica' (s) × 'Rocky Point'  
**H:** J. Barlup (1996), **G:** J. Barlup (2001), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)  
Fls 15/ball truss, broadly funnel-shaped, 44 × 76mm, with 6 very wavy-edged lobes, strong purplish red (60D) in bud, opening inside white (155D), light greenish yellow (4C) on apical part of the dorsal lobe, with inconspicuous pale yellow-green (4D) midveins on all lobes, and a moderate red (182A) flare on the dorsal lobe, extending 35mm from the base; outside white (155D), blending to pale purplish pink (65C) on dorsal lobes at 32mm from the base, with pale yellow-green (4D) midveins on all lobes; not scented. Calyx insignificant. Lvs elliptic, 120 × 54mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A), hairless. Shrub 1.5 × 2.0m in 16 years. Midseason (May, in Bellevue, Washington, USA).  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136147))

(a) '**Louise Alice**'

cv. Evergreen azalea: parentage unknown  
**H:** W.G.T. Hyde (1970s), **G:** W.G.T. Hyde (1970s), **N:** Mrs R. Legrand (2013), **I:** Mrs R. Legrand, **REG:** Mrs R. Legrand (2013)  
Fls single, open trumpet-shaped, 35 × 21mm, with 5 wavy-edged lobes, vivid red (57A) in bud and throughout on opening, with faint strong purplish red (60B) speckling on the 3 dorsal lobes. Calyx irregular, 5–8mm long, strong yellow-green (144C). Lvs elliptic, 25 × 11mm, cuneate at base, acute at apex, margins upcurved, matt green above, with no obvious indumentum. Shrub 1.2m × 61cm in 30–40 years. Very prolific in flower. May.  
(Etymology: named for a young mother fighting terminal cancer)

(a) '**LU Solveiga**'

cv. Deciduous azalea: 'Aida' (s) × 'Uguns'  
**H:** R. Kondratovičs (1978), **G:** R. Kondratovičs (1985), **N:** R. Kondratovičs (2013), **REG:** Rhododendron Nursery Babite, University of Latvia (2013)  
Fls single, 11/slightly domed truss, tubular funnel-shaped, 70mm wide, with 5–6 wavy-edged lobes, deep yellowish pink (46D) in bud, opening deep purplish pink (55A) both inside and out; with a brilliant orange-yellow (23B) blotch covering most of the dorsal lobe; small stamens, with orange anthers; style pink, stigma green. Calyx 5mm long, light green. Lvs elliptic, 130 × 40mm, pointed at base and apex, with wavy margins, glossy above, hairless. Shrub 2 × 2m in 15 years. Early June.  
(Nomenclatural standard: colour print provided by the registrant (WSY0119331). Etymology: LU represents the Latvian University, Solveiga is a Latvian female christian name)

(r) '**Madidi**'

cv. Elepidote rhododendron: 'Fuju-kaku-no-matsu' (s) × 'Fuju-kaku-no-matsu'  
**H:** F. Schnupper (2005), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls 10–12/conical truss, funnel-shaped, 40–50 × 50mm, with 5 wavy-edged lobes, moderate purplish pink (62B) in bud, opening inside pale purplish pink (62D), outside light purplish pink (62C); red/orange spots on the dorsal lobe; filaments and style purplish pink. Calyx 3mm long, with a grey indumentum. Lvs linear, 100–200 × 10mm, cuneate at base, acute at apex, margins flat and decurved, the leaf lamina curved longitudinally, with a brown indumentum below when mature. Dense shrub, 0.7 × 0.7m in 10 years. Early to mid May.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Bolivia)

(r) '**Main Street**'

Syn. of 'Thea Ann'

(r) '**Mansandra**'

cv. Elepidote rhododendron: 'Petra's Debut' (s) × unknown  
**H:** D.J. Dosser, **G:** D.J. Dosser (2012), **N:** D.J. Dosser, **REG:** D.J. Dosser (2013)  
Fls 14/ball truss, funnel-shaped, 90mm wide, with 6 wavy-edged lobes, moderate purplish red (58A) in bud, opening deep purplish pink (64D), unmarked; slightly scented. Lvs oblanceolate, to at least 150 × 45mm, cuneate at base, broadly acute and apiculate at apex, margins very narrowly recurved, hairless. Shrub to at least 1.5 × 1.0m. October to November (Tasmania).  
(Nomenclatural standard: colour print provided by the registrant (WSY0119015). Etymology: an amalgamation of the names of two towns in Victoria, Australia (Mansfield and Alexandra), in an area where the registrant grew up)

(a) '**Marie Susan Melkovitz**'

cv. Evergreen azalea: 'Mrs Nancy Dipple' (s) × 'Satellite' (Klupenger)

**H:** R. Stewart (1997), **G:** R. Stewart (1999),  
**N:** R. & S. Bauer (2012), **REG:** R. Stewart (2013)  
Fls hose-in-hose, 2/truss, open funnel-shaped, 57 × 70mm, with 10 very wavy-edged lobes, deep purplish pink (73A) in bud, opening inside deep purplish pink (73A), with a blotch of strong purplish red spots on the dorsal lobe; outside deep purplish pink (73A). No normal calyx present. Lvs elliptic, 57 × 22mm, cuneate at base, broadly acute at apex, with upcurved margins, moderate olive green (137A) and semi-glossy above. Dense shrub, 0.9 × 1.4m in 10 years. Midseason. (Nomenclatural standard: colour print of an electronic image (WSY0136146). Etymology: named in honour of the mother of Susan Bauer, who bought the naming rights in an auction held by the Northern Virginia Chapter of the American Azalea Society. Stewart ref. nos. RBS 42 and 2-97-1)

(v) **'Martin John'**

cv. *Elepidote rhododendron*: 'Saxon Glow' (s) × 'The Star'

**H:** N. Puddey (2008), **G:** N. Puddey (2009),  
**N:** N. Puddey (2010), **I:** N. Puddey (2012),  
**REG:** N. Puddey (2013)

Fls 3–4/open truss, funnel-shaped, 50 × 55mm, with 6 flat-edged lobes, moderate reddish brown (174A) in bud, opening inside vivid reddish orange (30B), fading to pale orange-yellow (24D) at the centre of the lobes and on the inside of the tube, outside vivid yellowish orange (30C) on the lobe tips, fading quickly to pale orange-yellow (24D) on the tube; unmarked. Calyx less than 1mm long, vivid reddish orange (33B). Lvs narrowly elliptic, narrowly tapering at base, acute at apex, glossy green above, scales dark reddish orange (175B) at first, not evident late. Shrub 0.5 × 0.25m in 4 years. Flowering irregularly. (Nomenclatural standard: colour print provided by the registrant (WSY0119008). Etymology: named for the brother of the registrant)

(r) **'Mary Collins'**

cv. *Elepidote rhododendron*: parentage unknown  
**H:** J.C. Cowles (1960s), **G:** Heritage Museums and Gardens (c. 1970), **N:** E.W. Collins (c. 2003),  
**REG:** Sandwich Club (2013)

Fls 12–14/ball truss, broadly funnel-shaped, 65 × 110mm, with 7 wavy-edged lobes, strong red (51A) in bud, opening inside strong purplish pink (63C) at the margins, centre white (155C), throat deep red (53A) extending 20mm from the base on the dorsal side, with strong red (53D) spotting and streaking on all lobes and with colourless streaking up the centre of all lobes; outside strong pink to moderate purplish pink (54C–54D), with strong red (51A) stripes. Calyx 2mm long, greyish red (182B). Lvs elliptic, 130 × 60mm, rounded at base, broadly acute at apex, margins upcurved, dull and moderate olive green (146A) above, hairless. Shrub 0.8 × 1.5m in 7 years. Late (May in Massachusetts, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136145). Etymology: named for the daughter of American Rhododendron Society member Edward W. Collins)

(r) **'Mary Margaret Haugen'**

cv. *Elepidote rhododendron*: 'Seaview Sunset' (s) × 'Elya' × (*yakushimanum* × 'C.I.S')

**H:** F. Fujioka (1997), **G:** F. Fujioka (2000),  
**I:** Meerkerk Rhododendron Gardens (2013),  
**REG:** F. Fujioka (2013)

Fls 18/domed truss, funnel-campanulate, 50 × 64mm, with 5 flat-edged lobes, vivid red (45C) in bud, opening inside pale yellowish pink (27C), with a deep yellowish pink (43C) picotee band (6mm wide) at the edge of all the lobes and a few deep yellowish pink (43C) spots on the dorsal lobes; outside vivid reddish orange (44C). Calyx 38mm long, vivid reddish orange (44C), with pale yellowish pink stripes. Lvs elliptic, 100 × 50mm, cuneate at base, acute at apex, margins downcurved, semi-glossy and dark bluish green (131A) above, hairs below moderate yellow (162A) at first, aging to moderate orange-yellow (164C). Dense shrub, 0.9 × 0.9m in 10 years. Midseason (April in Puget Sound, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136144). Etymology: named by several of her friends for a member (1993–2013) of the Washington State Senate, USA)

(r) **'Masai Mara'**

cv. *Elepidote rhododendron*: *bureauvii* (s) × *microgynum*

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls 8–9/domed truss, broadly funnel-shaped, 25–35 × 70–80mm, with 5 wavy-edged lobes, vivid purplish red (66A) in bud, opening inside moderate purplish pink (66D) to white, outside deep purplish pink (66C), unmarked; filaments white; style white, pink at apex, stigma greenish. Calyx 8mm long, brown-red. Lvs broadly elliptic, 90–100 × 50–60mm, cordate at base, broadly acute at apex, margins flat and decurved, glossy and very dark green above, indumentum white below at first, aging to cinnamon. Shrub 0.4–0.5 × 0.8m. Mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a national park in Kenya. Published in *Rhododendron und Immergrüne* 15 (2013))

(a) **'Meri'**

cv. *Evergreen azalea*: sport from 'Schloss Pillnitz'

**G:** R. Kondratovičs (1989), **N:** R. Kondratovičs (1989),  
**REG:** Botanical Garden, University of Latvia (2013)

Fls 1–4/truss, funnel-shaped, 30 × 45–50mm, with 5 flat-edged lobes (occasionally with some partially petaloid stamens), strong purplish red (58B), with white tips, opening inside deep purplish pink (N57C) in the centre, white at margins, outside strong purplish red (58B), with white margins; very weak markings; style red-purple, stigma sandy. Calyx 3mm long, light green. Lvs elliptic, 25 × 15mm, pointed at base and apex, margins flat, glossy green above, hairless. Shrub 0.9 × 1.1m in 15 years. February to March (under glass). (Nomenclatural standard: colour print provided by the registrant (WSY0119341). Etymology: Latvian female christian name)

(r) **'Mother'**

cv. *Elepidote rhododendron*: 'Annie Dosser' (s) × 'Red Gate'

**H:** D.J. Dosser (c. 1990), **G:** D.J. Dosser (2000), **N:** D.J. Dosser, **REG:** D.J. Dosser (2013)

Fls 15/domed truss, broadly campanulate, 50 × 40mm, with 6 wavy-edged lobes, strong purplish pink (68B) in bud, opening white inside and out, with deep purplish red (59B) spots all around the inside of the corolla.

Lvs lanceolate, 125 × 40mm, acute at apex, margins decurved, glossy green above, hairless. Shrub to at least 1.5 × 1.5m. September to November (Australia). (Nomenclatural standard: colour print provided by the registrant (WSY0119347))

(r) **'Murrindindi'**

cv. *Elepidote rhododendron*: 'Petra's Debut' (s) × unknown

**H:** D.J. Dosser, **G:** D.J. Dosser (2012), **N:** D.J. Dosser, **REG:** D.J. Dosser (2013)

Fls 14/conical truss, funnel-shaped, 100mm wide, with 5 wavy-edged lobes, vivid purplish red (57D) in bud, opening inside white, with broad vivid purplish red (57D) margins (which fade with age), outside vivid purplish red (57D); dorsal blotch of moderate purplish red (58A) spots (extending over 25mm); slightly scented. Lvs broadly elliptic, to at least 150 × 55mm, broadly cuneate at base, broadly acute and apiculate at apex, margins flat, hairless. Shrub to at least 1.0 × 1.0m. October to November (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0119011). Etymology: named for the shire to the northeast of Melbourne, Australia, which includes the registrant's home town of Alexandra)

(v) **'Noah's Gold'**

cv. *Lepidote rhododendron*: parentage unknown

**H:** N. Puddey (2008), **G:** N. Puddey (2010),

**N:** N. Puddey (2010), **I:** N. Puddey (2012),

**REG:** N. Puddey (2013)

Fls 5–8/flat truss, tubular funnel-shaped, 70 × 60mm, with 5 flat-edged lobes, light yellowish green (135D) in bud, opening vivid yellow (9B), unmarked; stigma and style light yellow-green (142C). Calyx less than 1mm long, light yellowish green (140D). Lvs narrowly elliptic, 110 × 35mm, rounded at base, acute at apex, margins upcurved, glossy green above, scaly at least at first. Shrub 1.5 × 0.8m in 4 years. Flowering irregularly, 2–3 times a year.

(Nomenclatural standard: colour print provided by the registrant (WSY0119034). Etymology: named after the registrant's first grandson)

(a) **'Norman's Choice'**

cv. *Deciduous azalea*: parentage unknown

**H:** W.G.T. Hyde (1970s), **G:** W.G.T. Hyde,

**N:** Mrs R. Legrand (2013), **I:** Mrs R. Legrand,

**REG:** Mrs R. Legrand (2013)

Fls 7–9/loose ball truss, flat open trumpet-shaped, 52 × 26mm, with 5–7 very wavy-edged lobes, vivid reddish orange (42A) in bud, opening inside vivid reddish orange (32A), outside shaded vivid to strong reddish orange (34A-C) and strong yellowish pink (32C); 2 dorsal lobes shaded dark greenish yellow

(152C) at the base to vivid yellow (17B) at the apex; scented. Calyx irregular, up to 5mm long, strong yellow-green (144A). Lvs broadly oblanceolate, 100 × 51mm, attenuate at base, narrowly acute at apex, margins upcurved, matt and rich bronze-coloured above. Shrub 1.8 × 1.02m in 30 years. May.

(Nomenclatural standard: colour print provided by the registrant (WSY0119345). Etymology: named for retired nurseryman Norman Standbrook)

(r) **'Noyo Star'**

cv. *Elepidote rhododendron*: 'Grandma's Hat' (s) × 'Red Eye'

**H:** K.D. Jones (2006), **G:** K.D. Jones (2012),

**N:** K.D. Jones (2013), **I:** Forest Lane Nursery,

**REG:** K.D. Jones (2013)

Fls 22/ball truss, broadly funnel-shaped, 81mm wide, with 5 very wavy-edged lobes, deep purplish red (71A) in bud, opening inside white (155C), with a picotee-like band on all margins of deep purplish red (71A), fading to very pale purple (69D), and with a prominent strong greenish yellow (151A) blotch covering the basal half of the dorsal lobe and extending slightly onto the adjacent lobes; outside deep purplish red (71A). Calyx insignificant. Lvs oblong, 169 × 48mm, rounded at base, acute and slightly twisted at apex, margins downcurved, semi-glossy and dark yellowish green (139A) above, hairless. Shrub 0.9 × 0.8m in 7 years. Midseason (May in Mendocino County, California, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136143))

(r) **'Orange Amber'**

cv. *Elepidote rhododendron*: 'Dazzler' (Elliott) (s) × 'Hill's Low Red'

**H:** J. Barlup (2006), **G:** J. Barlup (2010), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)

Fls 14/domed truss, 64 × 70mm, with 5 wavy-edged lobes, moderate purplish red (64A), opening inside brilliant yellow (11A) blending to deep yellowish pink (41D), with 5 deep red (53A), 3mm long nectar guides at the base and with deep purplish pink (54B) spotting on the dorsal lobes, 38mm from the base; not scented. Calyx 10mm long, deep yellowish pink (41D). Lvs elliptic, 114 × 51mm, rounded at base, broadly acute at apex, margins flat, dull and moderate olive green (147A) above. Shrub 0.6 × 0.8m in 6 years. Midseason (May, in Bellevue, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136142))

(r) **'Painted Bride'**

cv. *Elepidote rhododendron*: 'Mrs T.H. Lowinsky' (s) × *calophyllum*

**H:** J. Doppel (1996), **G:** J. Doppel (1999), **N:** T. Ahern (2013), **REG:** J. Doppel (2013)

Fls 11/ball truss, broadly funnel-shaped, 58 × 75mm, with 5 wavy-edged lobes, strong pink (48C) in bud, opening inside white, with a vivid red (45A) blotch and spots on the dorsal lobe, outside pale pink (49D); stamens almost absent. Calyx insignificant. Lvs obovate, 165 × 50mm, cuneate at base, rounded at apex, with downcurved margins, olive green above,

hairless. Shrub 1.8 × 3.0m in 17 years. Early midseason (mid April in eastern Pennsylvania, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136141))

(r) **'Patricia Hansen'**

cv. *Elepidote rhododendron*: 'Burgundy Mist' (s) × 'Holli's Hope'

**H:** J. Barlup (2006), **G:** J. Barlup (2010), **N:** J. Barlup (2013), **REG:** J. Barlup (2013)

Fls 17/ball truss, broadly funnel-shaped, 44 × 64mm, with 5 wavy-edged lobes, dark red (187A) in bud, opening inside strong purple (83D) and blending towards edges into strong purple (83C), with dark greenish yellow (152D) spotting on the dorsal lobe extending 29mm from the base; outside strong purple (83D), fading to deep purple (83B) towards the margins and with strong purplish red (71B) midveins from base to apex; not scented. Calyx insignificant. Lvs elliptic, 127 × 57mm, rounded at base, broadly acute at apex, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 6 years. Midseason (May, in Bellevue, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136140))

(r) **'Peucine'**

cv. *Elepidote rhododendron*: 'Feuervogel' (s) × 'New Comet'

**H:** K.-H. Hübbers (1986), **G:** K.-H. Hübbers (1994), **N:** K.-H. Hübbers (1997), **REG:** J. Westhoff (2013)

Fls up to 10/domed truss, open funnel-shaped, 50 × 95mm, with 7–8 wavy-edged lobes, opening inside pale yellow-green (4D), shading into pale greenish yellow (1D) in the throat, outside pale greenish yellow (2D); a few, small, strong yellow-green (145A) spots deep inside the corolla and forming a stream up to the dorsal lobe. Calyx 1mm long, greenish. Lvs oblong, 150 × 70mm, tapering at base, apiculate at apex, margins flat, glossy green above, hairless. Shrub 1.5 × 1.4m in 10 years. Mid May.

(Nomenclatural standard: colour print of an electronic image provided by the hybridizer (WSY0136139)).

Etymology: the original plant grows under a *Pinus peuce* in the registrant's garden.)

(a) **'Phil Legrand'**

cv. Deciduous azalea: parentage unknown

**H:** W.G.T. Hyde (1970s), **G:** W.G.T. Hyde, **N:** Mrs R. Legrand (2013), **I:** Mrs R. Legrand, **REG:** Mrs R. Legrand (2013)

Fls 13/compact, ball truss, open trumpet-shaped, 62 × 33mm, with 6 very wavy-edged lobes, vivid red (44A) in bud, opening inside vivid red (44A), shaded with vivid reddish orange (32A) on the dorsal lobe, outside vivid red (44A); long stamens. Calyx irregular, 5–10mm long, moderate yellow-green (146C). Lvs narrowly elliptic, 102 × 37mm, cuneate at base, acute at apex, margins upcurved, glossy and bronze-coloured above. Shrub 1.55 × 1.5m in 30 years. May. (Nomenclatural standard: colour print provided by the registrant (WSY0119343)). Etymology: named for the son of the registrant)

(a) **'Phil Louer'**

cv. Evergreen azalea: 'Desiree' (Shammarello) (s) × 'Painted Tips'

**H:** R. Stewart (1996), **G:** R. Stewart (1998), **N:** L. & D. Nanney (2012), **REG:** R. Stewart (2013)

Fls hose-in-hose, 2/truss, broadly funnel-shaped, 38 × 57mm, with 10 wavy-edged lobes, strong purplish red (67A) in bud, opening strong purplish red (67A), with moderate red spotting on the dorsal lobe; outside strong purplish red (67A). No normal calyx present. Lvs elliptic, 32 × 16mm, cuneate at base, broadly acute at apex, with upcurved margins, dark yellowish green (139A) and semi-glossy above. Dense shrub, 0.6 × 1.1m in 10 years. Midseason.

(Nomenclatural standard: colour print of an electronic image sent by the registrant (WSY0136138)).

Etymology: named to mark Phil Louer's long service to the Northern Virginia Chapter of the American Azalea Society. Naming rights acquired in a Chapter auction)

(r) **'Polonez'**

Syn. of 'Polonez Chopina'

(r) **'Polonez Chopina'**

cv. *Elepidote rhododendron*: *aureum* (s) × 'Tomasz Wojciech'

**H:** P. Muras (1995), **G:** P. Muras (2000), **N:** P. Muras (2011), **REG:** Mrs M. Muras (2013)

Fls 12–15/ovate truss, open funnel-campanulate, 35–40 × 55–60mm, with 5 wavy-edged lobes, strong pink (50C) in bud, opening inside creamy white (159B/D), with a dorsal blotch of vivid to brilliant yellow-green (154A/B) spots; outside creamy yellow (159B/D), with tinges of light pink on the midveins. Calyx 2mm long, reddish. Lvs oblong-elliptic, 60–110 × 25–40mm, cuneate at base, obtuse and apiculate at apex, margins flat, dark green and semi-glossy above, hairless. Rounded shrub, with a dense habit, 1.1 × 1.2m in 15 years. Late April to early May (southern Poland). Syn: 'Polonez'

(Nomenclatural standard: colour print provided by the registrant (WSY0119001)). Etymology: Polonez is a Polish folk dance, a form much used by the Polish composer Fryderyk Chopin)

(r) **'Ria'**

cv. *Elepidote rhododendron*: 'Graf Lennart' (s) × 'Blewbury'

**H:** K.-H. Hübbers (1987), **G:** K.-H. Hübbers (1992), **N:** Mrs L. Hübbers (2012), **REG:** J. Westhoff (2013)

Fls 7–10/flat, loose, uneven truss, campanulate, 30 × 30mm, with 5 wavy-edged lobes, vivid purplish red (57B) in bud, opening inside light to pale purplish pink (65B-D), outside moderate purplish pink (68C), with stripes of deep purplish pink (68A); dorsal lobe with a strong purplish red (67A), spotted blotch. Calyx 2mm long, light greenish. Lvs oblong, 65 × 35mm, truncate at base, acuminate at apex, margins flat to slightly upcurved, matt green above, hairless; new shoots bronze-coloured. Shrub 1.0 × 1.0m in 15 years. Early to mid April.

(Nomenclatural standard: colour print of an electronic image provided by the hybridizer (WSY0136137)).

Etymology: named for a niece of the raiser's wife)

(a) **'Ringla'**

cv. Deciduous azalea: 'Aida' (s) × 'Uguns'  
**H:** R. Kondratovičs (1978), **G:** R. Kondratovičs (1995),  
**N:** R. Kondratovičs (2012), **REG:** Rhododendron  
Nursery Babite, University of Latvia (2013)  
Fls single, 9/1 slightly domed truss, tubular funnel-  
shaped, 55mm wide, with 5 very wavy-edged lobes,  
vivid reddish orange (34A) in bud, with light orange-  
yellow (16C) midribs, opening strong red (39A) both  
inside and out, unmarked; filaments red, anthers light  
brown; style red, stigma green. Calyx 1mm long,  
green. Lvs elliptic, 120 × 45mm, pointed at base and  
apex, with wavy margins, matt green above, hairless.  
Shrub 2.5 × 3.0m in 25 years. Early June.  
(Nomenclatural standard: colour print provided by  
the registrant (WSY0119333). Etymology: Latvian  
female christian name)

(r) **'Rita May'**

cv. Elepidote rhododendron: parentage unknown  
**H:** J.C. Cowles (c. 1970), **G:** Heritage Museums and  
Gardens (mid 1970s), **N:** D.W. McGraw (2011),  
**I:** Briarwood Gardens (1988), **REG:** Heritage  
Museums and Gardens (2013)  
Fls 10/ball truss, broadly funnel-shaped, 50 × 80mm,  
with 6–7 wavy-edged lobes, strong purplish red (63A)  
in bud, opening inside pale purplish pink (62D),  
blending to deep purplish pink (66C) towards the  
margins, with pale greenish yellow (1D), twin rays,  
extending 10mm from the base; outside vivid  
purplish red (57D); stigma moderate purplish red  
(70A). Calyx insignificant. Lvs elliptic, 150 × 60mm,  
rounded at base, obtuse at apex, with flat margins,  
moderate olive green (137A) and matt above, hairless.  
Shrub 4 × 4m in 45 years.  
(Nomenclatural standard: colour print of an electronic  
image sent with registration (WSY0136136).  
Etymology: named for Rita May McGlynn, mother-  
in-law of David McGraw, a member of the Heritage  
Museums and Gardens Board. Briarwood ref. no.:  
Briarwood 508-69 (0508-1969))

(r) **'Robert Sutherland'**

cv. Elepidote rhododendron: 'Ronald Otto Delp' (s)  
× 'Golden Star' × 'Mezitt's Hardy Yellow'  
**H:** J. Looye (1994), **G:** J. Looye (1997), **N:** Amherstburg  
Rhododendron Committee Inc. (2012), **I:** Rhodo Land  
Nursery (2012), **REG:** Amherstburg Rhododendron  
Committee Inc. (2013)  
Fls 12/flattened dome truss, open campanulate, 64 ×  
76mm, with 5 wavy-edged lobes, buds light yellow  
(12C), flushed pink, opening light yellow (12C), with  
a moderate red (179A) blotch extending from the base  
over three quarters of the dorsal lobe; outside light  
yellow (12C). Calyx 6mm long, light yellow (12C).  
Lvs ovate-elliptic, 125 × 50mm, rounded at base,  
obtuse at apex, margins flat, moderate olive green  
(137B) above, hairless. Shrub 1.1 × 1.2m in 5 years.  
Midseason (May in southern Ontario, Canada).  
(Nomenclatural standard: colour print of an electronic  
image provided by the registrant (WSY0136135).  
Etymology: named for the founder of the  
Amherstburg Rhododendron Committee Inc.  
(Amherstburg, Canada))

(a) **'Rose Normandy'**

cv. Evergreen azalea: 'Elsie Lee' (s) × 'Satellite'  
(Klupenger)  
**H:** R. Stewart (1991), **G:** R. Stewart (1993),  
**N:** J.E. Gutierrez (2012), **REG:** R. Stewart (2013)  
Fls fully double, 2/truss, broadly funnel-shaped, 44 ×  
76mm, with 10–15 wavy-edged lobes, buds light  
reddish purple (74C) at the base, paler towards the  
apex, opening inside light reddish purple (74C), with  
a vivid reddish purple (74B) blotch on the dorsal  
lobes; outside light reddish purple (74C). Calyx  
10mm long, strong yellow-green (144A). Lvs elliptic,  
32 × 13mm wide, cuneate at base, broadly acute at  
apex, margins upcurved, semi-glossy and dark  
greenish yellow (136B) above. Shrub 0.9 × 1.2m in  
10 years. Midseason.  
(Nomenclatural standard: colour print of an electronic  
image provided by the registrant (WSY0136134).  
Etymology: named for the mother of Joseph  
E. Gutierrez (who purchased the naming rights at  
an American Azalea Society auction). Hybridizer's  
ref. no. RBS 93 and 1-91)

(a) **'Rožainā Zvaigznite'**

cv. Semi-evergreen azalea: 'Palestrina' (s) × 'Esmeralda'  
**H:** R. Kondratovičs (1972), **G:** R. Kondratovičs (1976),  
**N:** R. Kondratovičs (1976), **REG:** Botanical Garden,  
University of Latvia (2013)  
Fls single, 2–3/truss, funnel-shaped, 30 × 40–50mm,  
with 5 flat-edged lobes, deep purplish pink (N57C) in  
bud, opening inside strong purplish pink (68B),  
outside light purplish pink (68D), with very  
inconspicuous, non-contiguous markings; style white,  
stigma light pink (almost white). Calyx 3mm long,  
light green. Lvs elliptic, 20–40 × 10–15mm, pointed  
at base and apex, margins flat, matt green above,  
hairless. Shrub 1.0 × 0.9m in 15 years. February to  
March (under glass).  
(Nomenclatural standard: colour print provided by  
the registrant (WSY0119344). Etymology: Latvian  
for "pink little star")

(r) **'Ruth's Sister'**

cv. Elepidote rhododendron: 'Tolstead's Ma Belle' (s)  
× 'Virginia Delp'  
**H:** J. Doppel (2000), **G:** J. Doppel (2005), **N:** J. Doppel  
(2005), **REG:** J. Doppel (2013)  
Fls 13/ball truss, saucer-shaped, 50 × 75mm, with 5  
largely separated, wavy edged lobes, dark purplish red  
(N79C) in bud, opening inside white, with a broad,  
strong reddish purple, picotee-like edge and a  
prominent near black blotch occupying two-thirds of  
the dorsal lobe, outside strong reddish purple (72B);  
stigma white. Calyx insignificant. Lvs oblong, 150 ×  
40mm, rounded at base, acute at apex, with flat  
margins, mid green and glossy above, hairless. Shrub  
1.2 × 1.5m in 13 years. Midseason (early May in  
eastern Pennsylvania, USA).  
(Nomenclatural standard: colour print of an electronic  
image provided by the registrant (WSY0136133).  
Etymology: sister seedling of 'Ruth's Showstopper')

(r) **'Sammy Smyth'**

cv. Elepidote rhododendron: 'Countess of Lockington', open-pollinated (s)

**H:** D.J. Dosser (1990), **G:** D.J. Dosser (2000),

**N:** D.J. Dosser, **REG:** D.J. Dosser (2013)

Fls 10/ball truss, funnel-shaped, 115mm wide, with 7 wavy-edged lobes, buds pale yellow (11D), with a very slight pink tinge, opening pale yellow (11D) both inside and out, with a 13mm long, moderate purplish red (59C) blotch at the base of the dorsal throat; slightly scented. Calyx petaloid, almost as long as the corolla and a similar colour. Lvs narrowly obovate, to at least 100 × 30mm, cuneate at base, broadly acute and apiculate at apex, margins flat, hairless. Shrub 1.0 × 1.0m in 12 years. November to December (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0119013). Etymology: named for a friend of the registrant)

(a) **'Sandra's Pink Sparkles'**

cv. Evergreen azalea: probably 'Dainty Rose' × perhaps *kiusianum*

**H:** S.F. McDonald (1981), **G:** S.F. McDonald (1985),

**N:** S.F. McDonald (2013), **REG:** S.F. McDonald (2013)

Fls single, 3–4/truss, open funnel-shaped, 32 × 38mm, with 5 flat-edged lobes, moderate purplish pink (62B) in bud, opening light purplish pink (62C), with occasional vivid red (57A) or white stripes or sectors; outside pale purplish pink (62D). Calyx 5mm long, light yellow-green (145C). Lvs obovate 28 × 12mm, cuneate at base, apiculate at apex, margins flat, glossy and moderate olive green (137A) above, very sparsely hairy above and below, the hairs pale tan. Dense shrub, 1.2 × 0.9m in 30 years. Midseason (late April at south end of Chesapeake Bay, Virginia, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136132). Etymology: derived in part from the name of the registrant)

(r) **'Sandstone'**

cv. Elepidote rhododendron: 'Coral Blossom' (s) × 'Bambi' × *proteooides* (seedling no.8)

**H:** J. Barlup (2001), **G:** J. Barlup (2005), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)

Fls 16/domed truss, funnel-campanulate, 44 × 64mm, with 6 wavy-edged lobes, dark red (59A) in bud, opening inside pale yellow (11D), fusing with strong pink (50C) streaks, the margins strong pink (50C) and the 3 dorsal lobes with a solid flare, which is dark red (183A) at the base and changing to moderate red (183C) after 13mm and extending for 38mm from the base; outside pale yellow (11D) fusing with strong pink (50C), with strong red (50A) midveins; not scented. Calyx 28mm long, pale yellow (11D), with moderate red (183C) flaring. Lvs elliptic, 127 × 48mm, rounded at base, broadly acute at apex, margins downcurved, dull and moderate olive green (147A) above, hairless. Dense shrub, 0.6 × 0.9m in 11 years. Midseason (May, in Bellevue, Washington, USA).

(a) **'Santa'**

cv. Deciduous azalea: 'Aida' (s) × 'Uguns'

**H:** R. Kondratovičs (1978), **G:** R. Kondratovičs (1994),

**N:** R. Kondratovičs (2011), **REG:** Rhododendron

Nursery Babīte, University of Latvia (2013)

Fls 8–13/slightly domed truss, tubular funnel-shaped, 30 × 45mm wide, with 5 wavy-edged lobes, buds deep yellowish pink (43C), with pale yellow (18D) midribs, opening inside pale purplish pink (62D) in the centre, deep purplish pink (N57D) at the margins, outside deep purplish pink (N57D); a large vivid orange-yellow (21A) blotch covering much of the dorsal lobe; filaments pink, anthers orange; style pink, stigma green; slightly scented. Calyx 1mm long, light green. Lvs elliptic, 105 × 45mm, pointed at base and apex, with wavy margins, matt green above, hairless. Shrub 2.5 × 3.0m in 25 years. Early June.

(Nomenclatural standard: colour print provided by the registrant (WSY0119336). Etymology: Latvian female christian name)

(v) **'Sarah Ormiston'**

cv. Lepidote rhododendron: *macgregoriae* (s) × *luraluense*

**H:** A. Rouse (2001), **G:** A. Rouse (2009), **N:** A. Rouse (2013), **REG:** A. Rouse (2013)

Fls 6–9/open truss, with some flowers held upright, tubular funnel-shaped, 30 × 30–35mm, with 5 very slightly wavy-edged lobes, buds light yellow-green (145D), the bracts with deep purplish pink (58D) tips, opening both inside and out greenish white (157D) on the tube and with light yellowish pink (36C) lobes; unmarked; not scented. Pedicels pinkish, some angled at the point of attachment of the flower. Calyx less than 1mm long. Lvs elliptic, 50–60 × 25–35mm, tapering at base, apiculate at apex, margins slightly wavy, glossy green above, laxly scaly below. Shrub with an open habit, 1.2 × 1.0m in 12 years. June (Australia).

(Etymology: named for a friend of the registrant)

(r) **'Schaumperle'**

Syn. of 'Tindecor'

(r) **'Schneewalzer'**

cv. Elepidote rhododendron: 'Ahrensburg' (s) × 'Elizabeth de Rothschild'

**H:** H. Hesse (1997), **G:** H. Hesse (2002), **N:** H. Hesse, **REG:** H. Hesse (2013)

Fls 6–12/compact, rounded truss, broadly funnel-shaped, 45–65 × 80–105mm, with 5–6 wavy-edged lobes, buds pink lilac, opening white inside and out, with light green to yellow-green at the base of the throat and light green markings on the dorsal lobe. Calyx 1–2mm long, light green. Lvs elliptic, 50–135 × 20–60mm, rounded at base and apex, margins flat, matt green above, hairless. Compact shrub 1.1 × 1.2m in 10 years. Early to mid May.

(r) **'Serengeti'**

cv. Elepidote rhododendron: 'Witchery' (Henny) (s) × 'Lem's Monarch'

**H:** F. Schnupper (2005), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)

Fls 15–17/conical truss, funnel-campanulate, 40–50 × 60–70mm, with 5 wavy-edged lobes, dark red (59A) in bud, opening inside deep purplish red (59B), dark red (59A) outside; with dark red (59A) spots on the 3 dorsal lobes; filaments and style red. Calyx 3–4mm long, green to red. Lvs elliptic, 100–200 × 30–35mm, cuneate at base, broadly acute at apex, margins flat, matt green above, with brown hairs below. Shrub 1.0 × 1.0m in 10 years. Late May to early June.  
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY).  
Etymology: named for a national park in Tanzania)

(r) **'Sommertanz'**

cv. Elepidote rhododendron: 'Golden Splendour' (s) × 'Edward Dunn' × *auriculatum*

**H:** K.-H. Hübberts (2000), **G:** K.-H. Hübberts (2008), **N:** K.-H. Hübberts (2011), **REG:** J. Westhoff (2013)  
Fls 10/lax domed truss, narrow funnel-shaped, 80 × 90mm, with 7 very wavy-edged lobes, buds brilliant orange (29A), with yellow and red patches, opening inside brilliant yellow (13B), with pale yellow (12D) margins; outside brilliant yellow (13B/C), with some reddish patches; slightly scented. Calyx 2–15mm long, vivid reddish orange (30B/C). Lvs oblong-elliptic, 160 × 60mm, cuneate at base, broadly acute at apex, margins slightly decurved, matt green above, hairless; new shoots with bright red bracts. Shrub 2.2 × 1.5m in 10 years. June.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136129))

(r) **'Sonning'**

cv. Elepidote rhododendron: *hyperythrum* (s) × *aberconwayi*

**H:** Crown Estate Commissioners (1956), **G:** Crown Estate Commissioners, **N:** Crown Estate Commissioners (2013), **REG:** Crown Estate Commissioners (2013)

Fls 11–13/rounded and slightly flattened truss, 50 × 60mm, with 5 wavy-edged lobes, white in bud (between 155A & B, fading to 155B & C), opening pure white (155D) inside and out, except for a sprinkling of strong purplish red (58A-C) spots, mainly on the dorsal lobe but extending to adjacent lobes. Calyx 1–2mm long, pale whitish green. Lvs lanceolate, 70–90 × 25–30mm, cuneate at base, apiculate at apex, margins decurved, hairless. A 'well-clothed', domed shrub, 4–5 × 3m in 50 years; consistently floriferous. Mid to late April.

(Nomenclatural standard: herbarium specimen provided by the registrant (WSY0119002).  
Etymology: named after a Berkshire (UK) village and part of a series of hybrids raised by Hope Findlay

(working for the Crown Estate Commissioners), for which the names of Berkshire towns and villages have been used as the epithets)

(r) **'Summer Sunrise'**

cv. Elepidote rhododendron: 'Orange Prelude' (s) × 'Berg's Yellow' × *proteoides*

**H:** J. Barlup (2004), **G:** J. Barlup (2009), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)

Fls 15/ball truss, broadly funnel-shaped, 48 × 83mm,

with 7 wavy-edged lobes, strong yellowish pink (31C) in bud, opening inside pale yellowish pink (27C), with pale yellow-green (4D) towards the centre and a tinge of pale yellowish pink (36D) at the margins; strong red (53B) circle at the base, extending 10 mm from the base on all lobe axes; outside a blend of pale yellowish pink (27C) and pale yellow-green (4D), with tinges of pale yellowish pink (36D) on the dorsal lobes; not scented. Calyx 17mm long, pale yellow-green (4D), with two, 6mm long, strong red (53B) flares. Lvs elliptic, 120 × 46mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.9m in 7 years. Midseason (May, in Bellevue, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136128))

(r) **'Sun Blossom'**

cv. Elepidote rhododendron: 'Terra' (s) × 'Trina'  
**H:** J. Barlup (2005), **G:** J. Barlup (2012), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)

Fls 16/ball truss, broadly funnel-shaped, 51 × 77mm, with 7 wavy-edged lobes, deep yellowish pink (39B) in bud, opening inside pale yellow-green (4D), with dorsal lobe light greenish yellow (4B); deep red (185A) flare at base of dorsal throat, extending 19mm from the base and then extending as spots for 13mm; outside pale yellow-green (4D), with midveins and dorsal lobe light greenish yellow (4B). Calyx 10mm long, pale yellow-green, with a deep red (185A) flare. Lvs elliptic, 90 × 50mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.8m in 8 years. Midseason (May, in Seattle area, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136127).)

(v) **'Susan Marie'**

cv. Lepidote rhododendron: 'Kisses' (s) × 'Charming Valentino'

**H:** N. Puddey (2008), **G:** N. Puddey (2008), **N:** N. Puddey (2010), **I:** N. Puddey (2012), **REG:** N. Puddey (2013)

Fls 6–8/open truss, funnel-shaped, 45 × 50mm, with 5 flat-edged lobes, dark reddish orange (173A) in bud, opening inside vivid orange (28B) on the lobes and fading to light yellow (15D) in the tube, outside similarly coloured; unmarked. Calyx less than 1mm long, pale greenish yellow (10D). Lvs broadly elliptic, 110 × 55mm, rounded at base, acute at apex, margins decurved, glossy green above, scales white (155C) at first, not evident later. Shrub 1.0 × 0.5m in 4 years. Flowering irregularly, 2–3 times a year.

(Nomenclatural standard: colour print provided by the registrant (WSY0119038).  
Etymology: named for the sister of the registrant)

(r) **'Tangerine Charm'**

cv. Elepidote rhododendron: 'Cimarron Sun' (s) × 'September Song' × ('Bambi' × *proteoides*)

**H:** J. Barlup (2006), **G:** J. Barlup (2010), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)

Fls 11/domed truss, broadly funnel-shaped, 44 × 57mm, with 5 wavy-edged lobes, moderate red (47A) in bud, opening inside pale orange-yellow (16D) at base, blending into strong yellowish pink (34D) and with deep yellowish pink (47C) margins; no blotch or spots; outside pale orange-yellow (16D) at base, fading to vivid reddish orange; not scented. Calyx 17mm long, pale orange-yellow (16D), with a touch of strong yellowish pink (34D). Lvs elliptic, 102 × 38mm, rounded at base, broadly acute at apex, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.4 × 0.4m in 6 years. Midseason.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136126))

(r) **'Taroko'**

cv. *Elepidote rhododendron*: *yakushimanum*

'Koichiro Wada' (s) × *aberconwayi*

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** F. Schnupper, **REG:** F. Schnupper (2013)  
Fls 15–16/domed truss, campanulate, 45–50 × 50mm, with 5 flat to wavy-edged lobes, strong purplish pink (55B) in bud, opening inside light purplish pink (55C) to white, outside pale purplish pink (55D); slight orange spotting on the dorsal lobe; filaments and style very pale pink; anthers white; stigma yellowish. Calyx 2–3mm long, white. Lvs elliptic, 100–130 × 20–30mm, cuneate at base, acute at apex, margins flat and upcurved, matt green above, with white hairs below when young, aging to brown; lvs held upright so the indumentum is visible. Dense shrub, 0.45 × 0.8m in 10 years. Mid to late May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)). Etymology: named for a national park in Taiwan

(r) **'Teresa Smyth'**

cv. *Elepidote rhododendron*: 'Countess of Lockington', open-pollinated (s)

**H:** D.J. Dosser (1990), **G:** D.J. Dosser (2000), **N:** D.J. Dosser, **REG:** D.J. Dosser (2013)  
Fls 15/ball truss, funnel-shaped, 100mm wide, with 5 wavy-edged lobes, moderate yellowish pink (37C) in bud, opening pale greenish yellow (2D) both inside and out, aging to white, with moderate purplish red (59C) spots on the dorsal lobes; scented. Calyx 13mm long, pale greenish yellow (2D), flushed pink. Lvs elliptic, to at least 135 × 40mm, cuneate at base, narrowly obtuse at apex, margins recurved, hairless. Shrub 0.5 × 1.0m in 12 years. November (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0119009)). Etymology: named for a friend of the registrant

(r) **'Thea Ann'**

cv. *Elepidote rhododendron*: parentage unknown  
**H:** J.C. Cowles (c. 1970), **G:** Heritage Museums and Gardens (mid 1970s), **N:** J. & T. Stoneman (2011), **REG:** Heritage Museums and Gardens (2013)  
Fls 6–7/lax truss, broadly funnel-shaped, 65 × 100mm, with 6–7 wavy-edged lobes, light purplish pink (62C) in bud, opening inside white (155C), with 2–4 rays of deep pink (180D) spots on the 3 dorsal lobes,

extending from the base for one third of the corolla; white petaloid stamens mixed with normal stamens, 13mm long; stigma and style moderate yellow-green; outside white (155C), with light purplish pink (62C) vein stripes. Calyx insignificant. Lvs elliptic, 140 × 60mm, rounded at base, broadly acute at apex, with margins upcurved, moderate yellow-green (137C) and semi-glossy above, hairless. Shrub 1.5 × 1.5m in 12 years. Late.

Syn: 'Main Street'

(Nomenclatural standard: colour print of an electronic image sent with the registration (WSY0136125)). Etymology: named for Thea Ann Stoneman, a friend and supporter of the Heritage Museums and Gardens. Collector ref. no. 15-86; breeder ref. no. 0056-1994)

(a) **'Tidewater Firecracker'**

cv. Evergreen azalea: ('Ripples' × 'Girard's Hot Shot') × 'Girard's Hot Shot' (s) × 'Karafune'

**H:** S.F. McDonald (1992), **G:** S.F. McDonald (1995), **N:** S.F. McDonald (2013), **REG:** S.F. McDonald (2013)

Fls double, 2/truss, open funnel-shaped, 32 × 50mm, with more-or-less 15 wavy-edged lobes, vivid red (45A) in bud, opening inside vivid red (45A), with inconspicuous strong red (46A) spotting; outside vivid red (45A); occasional stamens present; pistil consistently present, vivid red (45A). Calyx 4mm long, vivid red (45C) in some flowers, strong yellow-green (145A) in others. Lvs obovate, 28 × 11mm, cuneate at base, apiculate at apex, with flat margins, glossy and moderate olive green (137A) above, in some winters strong red (46A), and with sparse, pale beige hairs above and below. Shrub 1.5 × 0.9m in 20 years. Midseason (Late April, south end of Chesapeake Bay, Virginia, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136124)). Hybridizer's ref. no. A93-1-4)

(v) **'Tijuana Bell'**

cv. *Lepidote rhododendron*: parentage unknown

**H:** N. Puddey (2010), **G:** N. Puddey (2010),

**N:** N. Puddey (2011), **I:** N. Puddey (2013),

**REG:** N. Puddey (2013)

Fls 4–6/lax truss, funnel-shaped, 75 × 80mm, with 5–6 wavy-edged lobes, dark reddish orange (173A) and pale yellow-green (193A) in bud, opening inside strong yellowish pink (43D) on lobes, fading to pale yellow (11D) in the tube, outside strong yellowish pink (43D) on lobes, fading to pale yellow (11D) at the base of the lobes and on the tube; unmarked; with a good scent. Calyx less than 1mm long, light greenish yellow (1C). Lvs ovate, 100 × 80mm, rounded at base, obtuse at apex, margins decurved, matt green above, scales white (155C) at first, moderate reddish brown (174A) later. Shrub 0.6 × 0.3m in 2 years. Flowering irregularly, twice a year.

(Nomenclatural standard: colour print provided by the registrant (WSY0119036)). Etymology: "Bell" is the nickname of the registrant's daughter and Tijuana Cartel is her favourite Australian band)

(r) **'Tinndecor'**

cv. Elepidote rhododendron: seedling from *decorum*

**R:** D.G. Hobbie (pre 1974), **S:** W. Wüstemeyer (1986), **N:** W. Wüstemeyer, **I:** H. Hachmann (2009), **REG:** J. Westhoff (2013)

Fls up to 19/loose, rounded truss, open funnel-shaped, up to 70mm wide, with up to 7 wavy-edged lobes, buds strong pink to moderate purplish pink (54C/D), opening inside pure white, outside white with pinkish stripes; with a few brown spots on the dorsal lobe. Calyx 2mm long, light green. Lvs oblanceolate, 120 × 35mm, tapered at base, acuminate at apex, margins flat, matt green above, hairless. Shrub 1.1 × 0.9m in 10 years. May.

Syn: 'Schaumperle'

(First published (as 'Tinndecor') in Schmalscheidt, W. *Rhododendron- und Azaleenzüchtung in Deutschland*, Teil II (1930–1988), 1989. Introduced by Hachmann as 'Schaumperle'. Etymology: the original plant was given to a rhododendron grower called H. Tinnefeld and is a *decorum* hybrid. The pollen parent may have been either *degronianum* subsp. *heptamerum* or *smirnowii*)

(r) **'Valdis'**

cv. Elepidote rhododendron: 'Uldis' (s) × unknown

**H:** R. Kondratovičs (1980), **G:** R. Kondratovičs (1986), **N:** R. Kondratovičs (2013), **REG:** Rhododendron Nursery Babite, University of Latvia (2013)

Fls 19/slightly domed truss, funnel-shaped, 70 × 60mm, with 5 wavy-edged lobes, deep purplish red (71A) in bud, opening inside strong purple (N80B) in the centre, strong reddish purple (N78B) at the margins; outside strong reddish purple (N78A); conspicuous blotch of greyish purple (N77A) spots on the dorsal lobe; filaments light purple, anthers light brown; style light purple, stigma dark purple. Calyx 2mm long, purple. Lvs elliptic, 110 × 55mm, pointed at base and apex, margins flat, glossy green above, hairless. Shrub 2.5 × 2.0m in 25 years. Early June.

(Nomenclatural standard: colour print provided by the registrant (WSY0119335). Etymology: Latvian male christian name)

(r) **'Velvet Touch'**

cv. Elepidote rhododendron: 'Terra' (s) × 'Berg's Yellow' × *proteoides*

**H:** J. Barlup (2004), **G:** J. Barlup (2011), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)

Fls 16/ball truss, broadly funnel-shaped, 41 × 64mm, with 7 wavy-edged lobes, strong yellowish pink (37A) in bud, opening inside pale greenish yellow (9D), with light greenish yellow (8C) on the dorsal lobe, blending on dorsal lobe margin to a 6mm wide band of light yellowish pink (36C) and brilliant yellow (9C); with a deep red (53A), 13mm long, solid circle around base of corolla and with deep red (53A) spots on the dorsal lobe; outside pale greenish yellow (9D), blending to light yellowish pink (36C) at the margins. Calyx 3mm long, deep red (53A), with pale greenish yellow (9D) edges. Lvs elliptic, 102 × 48mm, rounded at base, broadly acute at apex, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Dense shrub, 0.6 × 0.8m in 8 years. Midseason (May, in the Seattle area, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136123))

(v) **'Verona Star'**

cv. Lepidote rhododendron: parentage unknown

**H:** S. Saperstein, **G:** S. Saperstein, **S:** N. Puddey (2006), **N:** N. Puddey (2009), **I:** N. Puddey (2012), **REG:** N. Puddey (2013)

Fls 5–6/flat truss, funnel-shaped, 75 × 75mm, with 5 flat-edged lobes, moderate reddish brown (174A) in bud, opening deep pink (47D) at the lobe edges, fading to pale yellow (18C) at the centre and in the tube, outside deep pink (47D) at lobe edges and quickly fading to pale yellow (11D) on the basal part of the lobes and on the tube; unmarked. Calyx less than 1mm long, strong reddish orange (31B). Lvs broadly elliptic, 110 × 75mm, rounded at base, obtuse at apex, matt green above, scales white (155C) at first, not evident later. Shrub 1.2 × 0.5m in 5 years. Flowering irregularly, 2–3 times a year.

(Nomenclatural standard: colour print provided by the registrant (WSY0119012). Etymology: the plant was derived from a cross made by Sylvia Saperstein, and a Sylvia is one of the female characters in Shakespeare's play *The Two Gentlemen of Verona*)

(v) **'Vicky Griffith'**

cv. Lepidote rhododendron: *luraluense* (s) × *R. × psammogenes* var. *inundatum*

**H:** A. Rouse (2005), **G:** A. Rouse (2010), **N:** A. Rouse (2013), **REG:** A. Rouse (2013)

Fls 5–10/upright truss, shortly trumpet-shaped, 43 × 35mm, with 5 wavy-edged, reflexed lobes, pale yellow-green (149D) in bud, opening white inside and out, unmarked. The corolla is distinctly fluted and the face of the flower angled. Calyx insignificant. Pedicels 20mm long. Lvs 5–8 in pseudowhorls, broadly elliptic, 40–60 × 18–30mm, tapering at base, obtuse at apex, margins slightly wavy, glossy green above. Shrub 1.0 × 0.8m in 8 years. March (in Victoria, Australia).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the wife of the registrant)

(r) **'Violet Magic'**

cv. Elepidote rhododendron: 'Purple Amethyst' × 'Plum Passion' (s) × 'Holli's Hope'

**H:** J. Barlup (2006), **G:** J. Barlup (2010), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)

Fls 15/ball truss, broadly funnel-shaped, 51 × 76mm, with 5 very wavy-edged lobes, dark purple (79A) in bud, opening strong purplish red (72A), with deep purplish red (71A) margins and pale orange-yellow (23D) spotting on the 3 dorsal lobes; outside strong purplish red (72A), with deep purplish red (71A) midveins and margins; not scented. Calyx insignificant. Lvs elliptic, 146 × 51mm, rounded at base, broadly acute at apex, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Dense shrub, 0.46 × 0.76m in 5 years. Midseason (May, in Bellevue, Washington, USA). (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0136122))

(r) **'White Ginger'**

cv. *Elepidote rhododendron*: 'Phyllis Korn' (s) × 'Trude Webster'

**H:** J. Barlup (1993), **G:** J. Barlup (1999), **N:** J. Barlup (2011), **REG:** J. Barlup (2013)

Fls 13/ball truss, saucer-shaped, 64 × 102mm, with 5 wavy-edged lobes, moderate purplish pink (68C) in bud, opening inside white (155D), with a split or twinned flare of strong purplish red (60B) on the dorsal lobe, extending 38mm from the base; outside white (155D), with pale purplish pink (65D) midribs on the dorsal lobes; not scented. Calyx insignificant. Lvs elliptic, 178 × 95mm, rounded at base, broadly acute at apex, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.5 × 2.0m in 18 years. Midseason (May, in Bellevue, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the hybridizer (WSY0136121).

Etymology: name descriptive of the flower colour)

(r) **'Wilmor's Red'**

cv. *Elepidote rhododendron*: 'Cornubia' (s) × 'Coronation Day'

**H:** D.J. Dosser (1992), **G:** D.J. Dosser (1998), **N:** D.J. Dosser, **REG:** D.J. Dosser (2013)

Fls 12/ball truss, funnel-shaped, 125mm wide, with 5 wavy-edged lobes, deep red (60A) in bud, opening strong purplish red (58B) both inside and out, unmarked. Lvs broadly oblanceolate, 140 × 65mm, broadly cuneate at base, obtuse at apex, margins flat, hairless. Shrub 1.75 × 1.75m in 20 years. October (Tasmania).

(Nomenclatural standard: colour print provided by the registrant (WSY0119017). Etymology: named for the town in Tasmania where the registrant lives and the colour of the flowers)

(r) **'Winter Song'**

cv. *Elepidote rhododendron*: 'Lemon Embers' (s) × 'Windwalker'

**H:** J. Barlup (2003), **G:** J. Barlup (2009), **N:** J. Barlup (2012), **REG:** J. Barlup (2013)

Fls 23/ball truss, broadly funnel-shaped, 57 × 89mm, with 7 wavy-edged lobes, strong yellowish pink (37B) in bud, opening inside pale greenish yellow (2D), fading to white (155D) and with moderate red (185B), twin flares extending 13mm, then becoming spots for 25mm; outside pale greenish yellow (2D), fading to white (155D). Calyx insignificant. Lvs elliptic, 146 × 54mm, rounded at base, broadly acute at apex, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.5 × 1.8m in 10 years. Midseason (May, in the Seattle area, Washington, USA).

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY))

# Names and addresses

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

\* = deceased

Ahern, T.  
Bethlehem, Pennsylvania, USA

Amherstburg Rhododendron Committee Inc.  
Amherstburg, Ontario, Canada

Barlup, J.  
Bellevue, Washington, USA

Bauer, R. & S.  
Yorktown, Virginia, USA

Blackmore, R.  
Victoria, British Columbia, Canada

Botanical Garden, University of Latvia  
Riga, Latvia

Briarwood Gardens.  
East Sandwich, Massachusetts, USA

Burnham, T.  
Washington, D.C., USA

Clagett, W.L.  
Derwood, Maryland, USA

Collins, E.W.  
Hendersonville, North Carolina, USA

\*Cowles, J.C.  
Stow, Massachusetts, USA

Crown Estate Commissioners  
Windsor Great Park, Windsor, Berkshire, England

\*Dexter, C.O.  
Sandwich, Massachusetts, USA

Doppel, J.  
Lenhartsville, Pennsylvania, USA

Dosser, D.J.  
Lower Wilmot, Tasmania, Australia

Forest Lane Nursery  
Fort Bragg, California, USA

Fujioka, F.  
Freeland, Washington, USA

Griswold, D.A.  
Kirkland, Washington, USA

Griswold Nursery  
Kirkland, Washington, USA

Gutierrez, J.E.  
McLean, Virginia, USA

Hachmann, H.  
(Hans Hachmann), Baumschule H. Hachmann,  
Barmstedt, Germany

Hamilton, S.J.  
c/o Heritage Museums and Gardens, Sandwich,  
Massachusetts, USA

Heritage Museums and Gardens  
Sandwich, Massachusetts, USA

Hesse, H.  
Hamersen, Germany

Hübbers, K.-H. & Mrs L.  
Kranenburg, Germany

\*Hyde, W.G.T.  
Ferndown, Dorset, England

Jones, K.D.  
Fort Bragg, California, USA

Jørgensen, P.M.  
Det Norske Arboret, Hjeljestad, Norway

Klimavicz, J.F.  
Vienna, Virginia, USA

Kondratovičs, R.K.  
Riga, Latvia

Legrand, Mrs R.  
West Moors, Dorset, England

Looye, J.  
Rhodo Land Nursery, Niagara on the Lake,  
Ontario, Canada

Lyon, E.  
New Plymouth, New Zealand

Mac Mullan, D.  
Hamburg, Pennsylvania, USA

McDonald, S.F.  
Hampton, Virginia, USA

McGraw, D.  
c/o Heritage Museums and Gardens, Sandwich,  
Massachusetts, USA

Meerkerk Rhododendron Gardens  
Greenbank, Washington, USA

Morrissey, A.  
c/o Heritage Museums and Gardens, Sandwich,  
Massachusetts, USA

Muras, P. & Mrs M.  
Wieliczka, Poland

Nanney, L. & D.  
Springfield, Virginia, USA

O'Connor, Mrs J.M.  
Osterville, Massachusetts, USA

Player, L.M.  
Fayetteville, North Carolina, USA

Puddey, N.  
Woolgoolga, New South Wales, Australia

Rhododendron Nursery Babīte, University of Latvia  
Rīga, Latvia

Rhodo Land Nursery,  
Niagara on the Lake, Ontario, Canada

Ringstad, A.  
Gjøvik, Norway

Rouse, A.  
Hawthorn East, Victoria, Australia

Saint-Jalm, J.-F.  
Purit Beau Vallon, Séglien, France

Sandwich Club  
Sandwich, Massachusetts, USA

Saperstein, S.  
Mullumbimby, New South Wales, Australia

Schnupper, F.  
Westerstede, Germany

Snell, \*G.L.S. & Mrs W.  
Highfields, Queensland, Australia

Stewart, R.  
Springfield, Virginia, USA

Stoneman, J. & T.  
c/o Heritage Museums and Gardens,  
Massachusetts, USA

Visner, S.  
Washington, D.C., USA

Westhoff, J.  
Bremen, Germany

White's Nursery  
Germantown, Maryland, USA

Whitney Gardens  
Brinnon, Washington, USA

Wüstemeyer, W.  
Schermbek, Germany

Yamamoto, Y.  
Gamou district, Shiga-prefecture, Japan

# Corrections to the International Rhododendron Register and Checklist

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to the eight subsequent Supplements. The Registrar is grateful to all those who have brought such errors to his notice and welcomes knowledge of further amendments that may need to be made. Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version. The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the *Register and Checklist* (or one of its Supplements) on which the entry concerned is to be found.

(a) **'Aldis-ieva'** (p.35)

This should be written as 'Aldis-Ieva', as it represents a combination of two Latvian Christian names. It was raised in Latvia by K. Bauers & D. Getliņa in 1967. The description should read: Fls double, 2–3/truss, broadly funnel-shaped, 70–80mm wide, with 5 wavy-edged lobes, light pink (19/3), with an irregular white margin and strong cherry red spots. Calyx 3–5mm long. Lvs oblong or elliptic, 40 × 20mm, dark green. February to March.

(a) **'Bonbouche'** (p.162)

This should be 'Bon Bouche'

(a) **'Girard's Variegated Hot Shot'** (p.510)

Add the synonym 'Hot Shot Variegated' (p.618)

(a) **'Hot Shot Variegated'** (p.618)

This should be treated as a synonym of 'Girard's Variegated Hot Shot' (p.510)

**'Lil Clark'** (Seventh Supplement, p.18)

The following is a revised description based on further information provided by the registrant: Fls 10/ball truss, very broadly funnel-shaped, 100mm wide, with 5 wavy-edged lobes, deep reddish pink in bud, opening pale orange-yellow (19D) towards the base, broadly edged deep pink (185D), the latter colour fading with age; outside deep pink (185D); strong purplish red (59D) markings all around the base of the corolla; slightly scented. Calyx partially petaloid, 13mm long, similarly coloured. Lvs oblanceolate, 120 × 40mm, acute at apex, cuneate at base, margins decurved, matt green above, hairless. Shrub 1.0 × 0.75m in 6 years. November (Australia).

(a) **'Pissaro'** (p.1082)

This should be spelt 'Pissarro'


(a) **'Sārtā Niobe'** (p.1224)

The raiser listed as Kūrčevskis should read Karčevskis. The description should read: Fls double, 1–2/truss, broadly funnel-shaped, 80mm wide, with 5 wavy-edged lobes, carmine rose (621), with cherry red spots. Calyx 3–6mm long. Lvs oblong or elliptic, 30–40 × 15–20mm, dark green. February to March.

(r) **'Tukuma Pērle'** (p.1393)

This should be noted as an evergreen azalea, a sport from 'Niobe' and with the following description: Fls double, 1–2/truss, broadly funnel-shaped, 70mm wide, with 5 narrowly, but strongly wavy-edged lobes, white. Calyx 3–4mm long. Lvs oblong or elliptic, 30–50 × 13–18mm, green. February to March.

ISBN 9781907057519



9 781907 057519